

# Exam 2018-08-27

Maximilian Pfundstein

2019-03-17

## Contents

<b>1 Assignment 1 (3 points)</b>	<b>1</b>
1.1 Question 1.1 . . . . .	1
1.2 Question 1.2 . . . . .	1
1.3 Question 1.3 . . . . .	3
<b>2 Assignment 2</b>	<b>5</b>
2.1 Question 2.1 . . . . .	5
2.2 Question 2.2 . . . . .	6
2.3 Question 2.3 . . . . .	6





## 1 Assignment 1 (3 points)

### 1.1 Question 1.1

```
rng_runif = function(a, m, x_zero, nmax) {  
  
  if (!(a >= 0 && a < m)) stop("a in [0, m) is required")  
  if (nmax%%1 != 0) stop("nmax has to be an integer")  
  
  storage = vector(mode = "numeric", length = nmax+1)  
  storage[1] = x_zero  
  
  for (i in 1:(nmax-1)) {  
    storage[i+1] = ((a * storage[i]) %% m)  
  }  
  
  return(storage[2:length(storage)]/m)  
}
```

### 1.2 Question 1.2

```
prs_1 = rng_runif(69069, 2^32, 9999, 10000)  
prs_2 = rng_runif(630360016, (2^32-1), 690690, 10000)  
prs_3 = rng_runif(742938285, (2^32-1), 690690, 10000)  
prs_4 = rng_runif(1226874153, (2^32-1), 690690, 10000)  
  
unif_samples = runif(10000)
```



```
print(ks.test(x = prs_1, y = "punif"))
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: prs_1
## D = 0.010645, p-value = 0.2072
## alternative hypothesis: two-sided
```

```
print(ks.test(x = prs_2, y = "punif"))
```

```
## Warning in ks.test(x = prs_2, y = "punif"): ties should not be present for
## the Kolmogorov-Smirnov test
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: prs_2
## D = 0.018419, p-value = 0.002262
## alternative hypothesis: two-sided
```

```
print(ks.test(x = prs_3, y = "punif"))
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: prs_3
## D = 0.0063916, p-value = 0.8086
## alternative hypothesis: two-sided
```

```
print(ks.test(x = prs_4, y = "punif"))
```

```
## Warning in ks.test(x = prs_4, y = "punif"): ties should not be present for
## the Kolmogorov-Smirnov test
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: prs_4
## D = 0.013235, p-value = 0.06018
## alternative hypothesis: two-sided
```

```
print(ks.test(x = unif_samples, y = "punif"))
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: unif_samples
## D = 0.011378, p-value = 0.1501
## alternative hypothesis: two-sided
```

**Answer:** We see that the second and the last settings have a low-p value and have to be rejected at a significance level of  $\alpha = 10\%$

- $a \equiv 1 \pmod{p}$  for every prime divisor  $p$  of  $m$
- $a \equiv 1 \pmod{4}$  if 4 divides  $m$
- $c$  and  $m$  have to be relatively prime (no common divisors but 2)

In our case  $c = 1$ .

### 1.3 Question 1.3

```
rng_rnorm = function(n, a, m) {
  if (!(a >= 0 && a < m)) stop("a in [0, m) is required")

  n_half = ceiling(n/2)

  storage = vector(mode = "numeric", length = n)

  storage_theta = rng_runif(a = a, m = m, x_zero = 123456789, nmax = n_half) * 2 * pi
  storage_d = rng_runif(a = a, m = m, x_zero = 12345, nmax = n_half)

  for (i in 1:n_half) {
    storage[2*i] = sqrt(-2 * log(storage_d[i])) * cos(storage_theta[i])
    storage[2*i+1] = sqrt(-2 * log(storage_d[i])) * sin(storage_theta[i])
  }

  return(storage)
}
```

```
rng_normal_1 = rng_rnorm(10000, a = 69069, m = 2^32)
rng_normal_2 = rng_rnorm(10000, a = 630360016, m = 2^32-1)
rng_normal_3 = rng_rnorm(10000, a = 742938285, m = 2^32-1)
rng_normal_4 = rng_rnorm(10000, a = 1226874153, m = 2^32-1)
```

```
norm_samples = rnorm(10000)
```


```
## Warning: Removed 2 rows containing non-finite values (stat_bin).
```

```
## Warning: Removed 2 rows containing non-finite values (stat_bin).
```


```
## Warning: Removed 2 rows containing non-finite values (stat_bin).
```

```
## Warning: Removed 2 rows containing non-finite values (stat_bin).
```


a = 69069 and b =  $2^{32}$




a = 630360016 and b =  $2^{32} - 1$



a = 742938285 and b =  $2^{32} - 1$



a = 1226874153 and b =  $2^{32} - 1$



```
print(ks.test(x = rng_normal_1, y = "pnorm"))
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data:  rng_normal_1
## D = 0.0047014, p-value = 0.9799
## alternative hypothesis: two-sided
```

```
print(ks.test(x = rng_normal_2, y = "pnorm"))
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data:  rng_normal_2
## D = 0.015241, p-value = 0.01921
## alternative hypothesis: two-sided
```

```
print(ks.test(x = rng_normal_3, y = "pnorm"))
```

```
##  
## One-sample Kolmogorov-Smirnov test  
##  
## data: rng_normal_3  
## D = 0.006856, p-value = 0.7351  
## alternative hypothesis: two-sided
```

```
print(ks.test(x = rng_normal_4, y = "pnorm"))
```

```
##  
## One-sample Kolmogorov-Smirnov test  
##  
## data: rng_normal_4  
## D = 0.0094938, p-value = 0.3283  
## alternative hypothesis: two-sided
```

```
print(ks.test(x = norm_samples, y = "pnorm"))
```

```
##  
## One-sample Kolmogorov-Smirnov test  
##  
## data: norm_samples  
## D = 0.0097673, p-value = 0.2958  
## alternative hypothesis: two-sided
```

## 2 Assignment 2

### 2.1 Question 2.1

```
interpolate = function(A, X, func, gradient = NULL) {  
  
  # Interpolation function which is so be used  
  f_tilde = function(x, a0, a1, a2) a0 + a1 * x + a2 * x^2  
  
  # Error function, here MSE  
  f_error = function(A., X. = X, func. = func) {  
    return(sum((func.(X.) - f_tilde(X., A.[1], A.[2], A.[3]))^2))  
  }  
  
  #f_diff = function(A, func) {  
    # interpolation = a0 + a1 * x + a2 * x^2  
    # error = sum((interpolation - func(X))^2)  
    # return(error)  
  }  
  
  # Optimize f_error for parameters in A  
  res = optim(A, f_error, gradient, method = "BFGS")  
  
  # Now define with optimized parameters  
  f_tilde = function(x) res$par[1] + res$par[2] * x + res$par[3] * x^2
```

```

# Return parameters and function
return(list(A = res$par, f_tilde = f_tilde))
}

```

## 2.2 Question 2.2

```

f_approximate = function(func_target, bins, A_init, func_target_gradient = NULL) {

  # Initialize interval length and return matrix
  upper_boundary = 0
  interval_length = 1/bins
  res = matrix(0, nrow = bins, ncol = 5)
  colnames(res) = c("lower_boundary", "upper_boundary", "a0", "a1", "a2")

  # Approximate for each bin
  for (i in 1:bins) {
    lower_boundary = upper_boundary
    upper_boundary = lower_boundary + interval_length

    # We rely on three known points
    #known_values = c(func_target(lower_boundary),
    #                  func_target((lower_boundary + upper_boundary)/2),
    #                  func_target(upper_boundary))

    known_values = c(lower_boundary, (lower_boundary + upper_boundary)/2,
                     upper_boundary)

    # Now we optimize for this interval using the previous function
    interpolated = interpolate(func = func_target, X = known_values, A = A_init,
                              gradient = func_target_gradient)$A

    # And fill in the matrix with all necessary values
    res[i, 1] = lower_boundary
    res[i, 2] = upper_boundary
    res[i, 3:5] = interpolated
  }

  return(res)
}

```

## 2.3 Question 2.3


Definition of the original functions.

```

f1 = function(x) -x * (1 - x)
f2 = function(x) -x * sin(10 * pi * x)

```

The look like the following.



Now we perform the interpolation.

```
f1_res = f_approximate(f1, bins = 1000, rep(0.000, 3))
head(f1_res)
```

```
##      lower_boundary upper_boundary      a0      a1      a2
## [1,]          0.000          0.001 -7.697415e-08 -0.9990125 -0.0009988493
## [2,]          0.001          0.002  5.757218e-06 -1.0022222 -0.0030065828
## [3,]          0.002          0.003 -5.377707e-06 -0.9952695 -0.0049762640
## [4,]          0.003          0.004 -1.209750e-05 -0.9929714 -0.0069507160
## [5,]          0.004          0.005 -2.025462e-05 -0.9909215 -0.0089182095
## [6,]          0.005          0.006 -3.040951e-05 -0.9888805 -0.0108776017
```

```
f2_res = f_approximate(f2, bins = 1000, rep(0.005, 3))
head(f2_res)
```

```
##      lower_boundary upper_boundary      a0      a1      a2
## [1,]          0.000          0.001 2.617655e-06 -0.03141572 0.004960967
## [2,]          0.001          0.002 6.538140e-05 -0.09418439 0.004699845
## [3,]          0.002          0.003 1.905340e-04 -0.15676486 0.004188603
## [4,]          0.003          0.004 3.773268e-04 -0.21903202 0.003429245
## [5,]          0.004          0.005 6.246391e-04 -0.28086124 0.002424771
## [6,]          0.005          0.006 9.309811e-04 -0.34212856 0.001179168
```

```
f_tilde = function(x, a0, a1, a2) a0 + a1 * x + a2 * x^2
```

```
df$f1_y_interpolated = sapply(sequence, FUN = function(x) {
  target_row = f1_res[x >= f1_res[,1] & x < f1_res[,2]]
  return(f_tilde(x, target_row[3], target_row[4], target_row[5]))
})
```

```

})

df$f2_y_interpolated = sapply(sequence, FUN = function(x) {
  target_row = f2_res[x >= f2_res[,1] & x < f2_res[,2]]
  return(f_tilde(x, target_row[3], target_row[4], target_row[5]))
})

```

