

CHAPTER OVERVIEW

6: Chemical Bonding - Electron Pairs and Octets

Theories of chemical bonding invariably involve electrons. When one atom approaches another, the valence electrons, found in the outermost regions of the atoms, interact long before the nuclei can come close together. Electrons are the least massive components of an atom, and so they can relocate to produce electrostatic forces which hold atoms together. According to Coulomb's law, such electrostatic or Coulombic forces are quite large when charges are separated by distances of a few hundred picometers—the size of an atom. Coulombic forces, then, are quite capable of explaining the strengths of the bonds by which atoms are held together.

- 6.1: Prelude to Chemical Bonding
- 6.2: Ionic Bonding
- 6.3: Energy and the Formation of Ions
- 6.4: The Ionic Crystal Lattice
- 6.5: Ions and Noble-Gas Electron Configurations
- 6.6: Ionization Energies
- 6.7: Ionization of Transition and Inner Transition Elements
- 6.8: Electron Affinities
- 6.9: Binary Ionic Compounds and Their Properties
- 6.10: The Octet Rule
- 6.11: Physical Properties
- 6.11.1: Lecture Demonstrations
- 6.12: Chemical Properties
- 6.13: The Covalent Bond
- 6.14: Covalent Molecules and the Octet Rule
- 6.15: Writing Lewis Structures for Molecules
- 6.15.1: An Excess of Bonds
- 6.15.2: Deciding on a Skeleton Structure
- 6.15.3: Multiple Bonds
- 6.16: Examples of Lewis Structures
- 6.17: Polyatomic Ions
- 6.18: Ionic Compounds Containing Polyatomic Ions
- 6.19: Atomic Sizes
- 6.20: Ionic Sizes
- 6.21: Periodic Variation of IE and EA
- 6.21.1: Lecture Demonstrations

This page titled 6: Chemical Bonding - Electron Pairs and Octets is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Ed Vitz, John W. Moore, Justin Shorb, Xavier Prat-Resina, Tim Wendorff, & Adam Hahn.