


Project 1 - TMA4320

Modelling motor proteins as particles diffusing in parallel potentials

Haakon T. Simensen

January 22, 2020



This project description consists of 6 numbered sections. Of these sections, two sections [1 and 2] are pure theory sections where you get some useful background information. Sec. 3 contains some theory, as well as a few analytical exercises which are meant to give you insight into diffusion processes. The remaining three sections [4, 5 and 6] consist of programming exercises where you will do random walk simulations. The overall goal is to be able to model a complex biological process using a simple theoretical model which reflects some important aspects of the real process. I suggest to do the programming in python, as this is the only programming language in which you are guaranteed to get help from the student assistants. However, you are welcome to use e.g. C++ if you want. Some of the calculations may require some computing time, depending upon how efficiently you write the code (and what programming language you use). Therefore, I urge you to do the debugging phase of the coding with decreased numerical accuracy (e.g. number of particles, number of time steps, etc.), so that you do not waste time on waiting for the computer to produce rubbish results.

You should submit a report where you answer each exercise in the same order as given in this project description. The report should either be a .pdf-file written in L^AT_EX, or handed in as a Jupyter notebook accompanied with a .html- or -.pdf-file which is easily printable for me. Use the exercise numbers given here to mark your answers. You should only give explicit answers and discussions to the exercises. Do not include an introduction, methods, conclusion or any additional sections which you might be used to include in lab reports etc.. Do not write your names on the report, as we want the grading to be done anonymously. You are free to write the report in either English or Norwegian.

Your grade will be determined by the correctness of your results, your discussion of the results, and how you relate the computational results to real world physics. In general, the last exercises will weigh more on your final grade than the first exercises. Your code in itself will generally not be subject to grading. Since tiny syntax errors may have huge qualitative impact on your results, a certain degree of mercy will be given to those with erroneous results. That is, you will not be given the maximum score if your results are obviously physically incorrect, but your discussion viewed in light of the results will be judged in your favour. If you understand that the results are physically incorrect, you will be awarded for pointing out what is wrong, and why this is wrong, as this shows physical intuition and insight into the involved physics. That said, the final grade will be based on the overall impression of your report. In rare cases, your code might be used in the evaluation, so you should submit your code alongside the report (if you hand in a Jupyter notebook with all your code, you need not hand in a separate file containing your code).

1 Introduction

Motor proteins are, in short terms, proteins which are able to turn chemical energy into mechanical energy. Take for instance myosin. Myosin gets energy from ATP molecules, and turns this energy into mechanical movement by dragging a thin filament parallel to the myofilament they are attached to. The collective action of many such myosin proteins is what we know as muscle contractions! See Fig. 1 for an illustration of this process.




Figure 1: An illustration of myosin-ATPase, retrieved from [here](#).

A common way to model such processes theoretically is to model the motor protein as particles moving in several, parallel and periodic potentials. Transitions between the potentials (or energy levels) are triggered by catching an ATP-molecule, or releasing $ADP + P$. If the periodic potentials are symmetric, one can easily form an argument for why the net particle current in this case must average to zero. The key to motor protein transport is therefore the (asymmetric) shape of the periodic potentials.

In this exercise, we will model the motor protein transport as a current of particles moving in a 1-dimensional, two-level potential landscape, using random walk simulations. Before digging into the full-scale problem, we will start with some theory and warm-up exercises to get you going. In the end, we will compare our results with one of the few analytic solutions which exists to this rather complex problem.

2 Theory

Mathematical introduction

Diffusion is a process where particles move down a concentration gradient. That is, particles move from areas with high concentration to areas with low concentration. The physical driving force of diffusion is simply random movement of each particle. Diffusion of a substance, of which distribution is given by $\phi(x, t)$, is described by the diffusion equation. In one dimension (1D) the diffusion equation takes the form

$$\frac{\partial \phi(x, t)}{\partial t} = \frac{\partial}{\partial x} \left(D(x) \frac{\partial \phi(x, t)}{\partial x} \right), \quad (1)$$

where $D(x)$ is the (position dependent) diffusion coefficient.

We will now show the correspondence between random walk in 1D and the diffusion equation. Random walk is defined as follows: let $X(t)$ be a stochastic process in time giving the position of a particle after a time t . At time $t = 0$, $X(0) = 0$. The particle makes a step in every time interval Δt . The particle can move left and right along an axis, or stay put, with the following probabilities

$$X(t + \Delta t) = \begin{cases} X(t) + h & \text{with probability } \frac{d}{2}, \\ X(t) & \text{with probability } 1 - d, \\ X(t) - h & \text{with probability } \frac{d}{2}. \end{cases} \quad (2)$$

We now define the function $\phi(x, t) = P\{X(t) = x\}$. By conditional probabilities, we may straightforwardly write

$$\phi(x, t + \Delta t) = \frac{d}{2} \phi(x + h, t) + (1 - d) \phi(x, t) + \frac{d}{2} \phi(x - h, t). \quad (3)$$

Slightly rewrite this to

$$\frac{\phi(x, t + \Delta t) - \phi(x, t)}{\Delta t} = \frac{dh^2}{2\Delta t} \frac{\phi(x + h, t) - 2\phi(x, t) + \phi(x - h, t)}{h^2}. \quad (4)$$

DEF. DERIVATIVES

Now suppose that $D = \frac{dh^2}{2\Delta t}$ is a constant. If we then let Δt and h^2 go simultaneously to zero, we find

$$\frac{\partial \phi(x, t)}{\partial t} = D \frac{\partial^2 \phi(x, t)}{\partial x^2}. \quad (5)$$

Random assumption? or is this correct?

This is the diffusion equation with constant diffusion coefficient. We have therefore shown that random walk in the limit of infinitesimal steps in time and space corresponds to diffusion.

3 Exercises

The diffusion equation

Exercise 3.1

□ Show that

$$\phi(x, t) = \frac{1}{\sqrt{4\pi Dt}} \int_{-\infty}^{\infty} dy h(y) e^{-\frac{(x-y)^2}{4Dt}} \quad (6)$$

D(x) or D = const.?

is a solution of the diffusion equation.

□ Let $\phi(x, t)$ represent a distribution of particles along the x -axis at time $t \geq 0$. Give a physical interpretation of $h(y)$ (Hint: let $t \rightarrow 0$ in Eq. (6)).

Exercise 3.2

Assume that the probability of finding a particle at position x in a potential landscape $V(x)$ is given by¹

$$P(x) = Ze^{-\beta V(x)}, \quad (7)$$

where $V(x)$ is the particle's potential energy at position x , $\beta = (k_B T)^{-1}$ where k_B is the Boltzmann's constant and T is the temperature, and Z is the canonical partition function of the system.

We now want to perform a random walk using these probabilities to determine the movement of the particles. At each step in the random walk, we allow the particle only to move one step to the right, one step to the left, or staying put. We define the probability of moving one step to the right as p^+ , the probability of moving one step to the left as p^- , and the probability of staying put as p^0 . Let the probability of a particle moving to position x from a position $x_0 \in \{x-1, x, x+1\}$ during one time step be proportional to $P(x)$, and assume that the proportionality constant is equal for all step options.

- Show that, if the position of the particle before a random walk step is x_0 , the probabilities $\{p^+, p^0, p^-\}$ are

$$\left. \begin{aligned} p^+ &= \frac{1}{1 + e^{-\beta[V(x_0-1)-V(x_0+1)]} + e^{-\beta[V(x_0)-V(x_0+1)]}}, \\ p^0 &= \frac{1}{1 + e^{-\beta[V(x_0-1)-V(x_0)]} + e^{-\beta[V(x_0+1)-V(x_0)]}}, \\ p^- &= \frac{1}{1 + e^{-\beta[V(x_0+1)-V(x_0-1)]} + e^{-\beta[V(x_0)-V(x_0-1)]}}. \end{aligned} \right\} \begin{array}{l} \text{from} \\ (7) \end{array} \quad (8)$$

$p^+ = \frac{P(x_0+1)}{P(x_0+1) + P(x_0) + P(x_0-1)}$

Exercise 3.3

- What happens to the probabilities $\{p^+, p^0, p^-\}$ given by Eq. (8) when the temperature T satisfies the following conditions:

- $k_B T \gg |V(x+1) - V(x)|$ for any x ?
 - $k_B T \ll |V(x+1) - V(x)|$ for any x ?
- $\left. \begin{array}{l} \bullet k_B T \gg |V(x+1) - V(x)| \text{ for any } x? \\ \bullet k_B T \ll |V(x+1) - V(x)| \text{ for any } x? \end{array} \right\} \text{SE PÅ } \beta \quad e^{-\beta[V(x_0-1) - V(x_0)]} = 0$

An important tool for physicists is identifying the energy scales relevant to what we want to compute. In certain situations, we may ignore some interactions in a problem because the corresponding energy scale can be neglected compared to some other interaction. For example, when you compute the driving time from Trondheim to Oslo, you can with great precision ignore quantum effects and relativistic effects such as time dilation, as these effects are small on the relevant energy scale.

- How can we simplify the random walk problem using an energy scale argument in the two limits given above?

4 Warm-up exercises

Random walk in a potential

We will now work through some examples of random walk in one dimension. In all examples, start with 10000 particles at $x = 0$. Let each of these particles be subject to a random walk for 200 time steps, where the step length is $h = 1$, and the time step is $\Delta t = 1$ (we use dimensionless time and space axes). Simulate the random walk by picking a random number between 0 and 1, and walk to

¹This is not a totally random probability distribution, but a well known (although slightly simplified) result from statistical physics.

$$k_B T \ll [V(x+1) - V(x)]$$

$$p^+ = \frac{1}{1 + e^{-\beta[V(x_0-1) - V(x_0+1)]} + e^{-\beta[V(x_0) - V(x_0+1)]}},$$

$$p^0 = \frac{1}{1 + e^{-\beta[V(x_0-1) - V(x_0)]} + e^{-\beta[V(x_0+1) - V(x_0)]}},$$

$$p^- = \frac{1}{1 + e^{-\beta[V(x_0+1) - V(x_0-1)]} + e^{-\beta[V(x_0) - V(x_0-1)]}}.$$

$$e^{-\beta V(x_0)}$$

Exercise 3.3

What happens to the probabilities p^+ , p^0 , p^- given by Eq. (9) when the

$$p^0 = \frac{1}{1 + e^{-\beta[V(x_0-1) - V(x_0+1)]}} \cdot \frac{e^{-\beta V(x_0)}}{e^{-\beta V(x_0)}}$$

$$= \frac{e^{-\beta V(x_0)}}{e^{-\beta V(x_0)} + e^{-\beta[V(x_0-1) + V(x_0) - V(x_0+1)]}} \cdot \frac{e^{\beta V(x_0-1)}}{e^{\beta V(x_0-1)}}$$

$$= \frac{e^{-\beta[V(x_0) - V(x_0-1)]}}{e^{-\beta[V(x_0) - V(x_0-1)]} + e^{-\beta[V(x_0) - V(x_0+1)]}}$$

$$= \frac{e^{-\beta[V(x_0) - V(x_0-1)]}}{e^{-\beta[V(x_0) - V(x_0-1)]} + e^{\beta[V(x_0+1) - V(x_0)]}} = \frac{1}{\infty} = 0$$

$$k_B T \gg [V(x+1) - V(x_0)]$$

$$p^+ = \frac{1}{1 + e^{-\beta[V(x_0-1) - V(x_0+1)]} + e^{-\beta[V(x_0) - V(x_0+1)]}},$$

$$p^0 = \frac{1}{1 + e^{-\beta[V(x_0-1) - V(x_0)]} + e^{-\beta[V(x_0+1) - V(x_0)]}},$$

$$p^- = \frac{1}{1 + e^{-\beta[V(x_0+1) - V(x_0-1)]} + e^{-\beta[V(x_0) - V(x_0-1)]}}.$$

$$p^0 = \frac{1}{3}$$

$$p^+ = \frac{1}{2 + e^{-\beta[V(x_0-1) - V(x_0+1)]}} \cdot \frac{e^{-\beta V(x_0)}}{e^{-\beta V(x_0)}} = \frac{e^{-\beta V(x_0)}}{2e^{-\beta V(x_0)} + e^{-\beta[V(x_0-1) + V(x_0) - V(x_0+1)]}}$$

$$= \frac{e^{-\beta[V(x_0) - V(x_0-1)]}}{2e^{-\beta[V(x_0) - V(x_0-1)]} + e^{-\beta[V(x_0) - V(x_0+1)]}}$$

$$= \frac{e^{-\beta[V(x_0) - V(x_0-1)]}}{2e^{-\beta[V(x_0) - V(x_0-1)]} + e^{\beta[V(x_0+1) - V(x_0)]}} = \frac{1}{3}$$

the left if this number is equal to or less than p^- , walk to the right if it is greater than $(1 - p^+)$, and stay put otherwise. In each example a potential is defined, and you should use Eq. (8) to determine $\{p^+, p^0, p^-\}$ for each particle at each time step.

The quantity βk can to some extent be interpreted as the ratio between the potential energy scale and thermal energy scale relevant to a particle diffusing through the potentials below². Plot the particle distribution along the x -axis after the random walk simulation for $\beta k \in \{0.01, 1, 100\}$ for each potential. Explain the different results for the different values of βk using your knowledge of the involved physics.

Programming exercise 4.1

$$V(x) = k$$

Fit a normal distribution to the resulting particle distribution (use for instance the function `fit` in `scipy.norm`), and explain the results using Sec. 2.

Programming exercise 4.2

$$V(x) = kx$$

Fit a normal distribution to the resulting particle distribution.

Programming exercise 4.3

$$V(x) = k \left(\frac{x}{20} + \sin \left(\frac{x}{5} \right) \right)$$

As an additional task for this potential: try running the simulation at 200 time steps using $\beta k = 0.01$, and then set $\beta k = 100$ and run the simulation for an additional 200 time steps. What do you observe?

Programming exercise 4.4

$$V(x) = kx^2$$

Can βk alone be used as a ratio to determine whether the thermal energy or potential energy dominates the random walk behavior in this potential? Why, or why not?

5 Random walk in a ratchet potential

We will now create a toy model for motor protein motion by modelling them as particles moving in two parallel and periodic potentials. Although the model is simplified, this is in fact one of the main theoretical principles used to model motor protein motion!

Define two potentials, $V_1(x)$ and $V_2(x)$. The particles can move inside the potential landscapes defined by $V_1(x)$ and $V_2(x)$, and they can be excited (de-excited) between these energy levels. $V_2(x)$ is a constant function of x , and its magnitude is not important. $V_1(x)$ is a periodic potential. Let the x -axis be discretized with step length $h = 1$, and let the spatial period of $V_1(x)$ be N_x steps. We then have $V_1(x + nN_x) = V_1(x)$ for $n \in \mathbb{Z}$. $V_1(x)$ is defined on the interval $x \in \langle -(1 - \alpha)N_x, \alpha N_x \rangle$ as

$$V_1(x) = \begin{cases} k \frac{x}{\alpha N_x} & 0 < x \leq \alpha N_x, \\ -k \frac{x}{(1-\alpha)N_x} & -(1-\alpha)N_x < x \leq 0, \end{cases} \quad (9)$$

²Remember that we operate with dimensionless position x , so that k is given in units of energy.

where $0 \leq \alpha \leq 1$. The shape of this potential is a so-called **sawtooth wave**. α defines the x -positions of the teeth's peaks. In particular, $\alpha = 0.5$ results in a symmetric potential about $x = 0$. k determines the strength of the potential.

The particles living in this system are excited and de-excited between the energy levels 1 and 2 with respective potential landscapes $V_1(x)$ and $V_2(x)$ at even intervals T_p . That is, if the particles start in **energy level 2 at $t = 0$, the particles are de-excited to energy level 1 at $t = T_p$** . They live in the potential landscape defined by $V_1(x)$ in the time interval $T_p \leq t < 2T_p$, before the particles are excited to live in the potential landscape defined by $V_2(x)$ in the time interval $2T_p \leq t < 3T_p$, etc. In the **(de-)excitation process, the x -coordinate is conserved**, so that a particle situated at x_j before an excitation moves from $V_1(x_j) \rightarrow V_2(x_j)$.

Define the **normalized particle current $J(t_i)$** as

$$J(t_i) = \frac{n^+(t_i) - n^-(t_i)}{N_p}, \quad (10)$$

where $n^+(t_i)$ is the number of particles moving to the right on the time interval $(t_i - t_{i-1})$, $n^-(t_i)$ is the number of particles moving to the left, and N_p is the total number of particles in the system.

5.1 Programming exercise 5.1

Let $\alpha = 0.8$ and $T_p = 500$. Use step length $h = 1$, $N_x = 90$ steps per potential period, time step $\Delta t = 1$ and $\beta k = 1000$. Define an x -axis that fits two full sawtooth peaks (in total $2N_x = 180$ discrete points), and use periodic boundary conditions. That is, if a particle moves to the left from the leftmost point x_0 , it arrives at x_{179} (if we use zero-based numbering). Let the number of particles in the system be $N_p = 12N_x$. Let the particles start in $V_2(x)$ and be evenly distributed along the x -axis at $t = 0$, so that there are six particles per discrete x -point to start with. They de-excite to $V_1(x)$ after $T_p = 500$ time steps, and stay there for $T_p = 500$ time steps. We will further on refer to two such periods as a cycle.

Run the random walk simulation for $20T_p = 10000$ time steps, that is 10 cycles. Calculate the cycle-averaged current for each of the 10 cycles, $J_{\text{avg}}(n) = \frac{1}{2T_p} \sum_{i=1+2T_p n}^{2T_p(n+1)} J(t_i)$, where n labels the cycles. You should obtain negative average currents.

- Write down the 10 cycle-averaged currents you obtain.
- Why do we get non-zero currents? Give a physical interpretation of this.
- Compare $J_{\text{avg}}(0)$ with $J_{\text{avg}}(n)$ for $n > 0$. Give a physical interpretation of this difference.

Repeat this process, only this time with $\alpha = 0.2$. The cycle-averaged currents should now be positive. Give a physical interpretation of this result compared to what you obtained with $\alpha = 0.8$.

Programming exercise 5.2

From now on and in the rest of this project, start each simulation with half of the particles at each of the two x -coordinates which coincide with the minima of $V_1(x)$ (however, let the particles start in energy level 2 defined by $V_2(x)$). Reset the system into this state for every new set of parameters T_p or α .

Choose 50 values of T_p equally spaced between 1 and 500, and run the random walk simulation for one cycle for each of these values. Let the number of particles in the system be $N_p = 40N_x$, and use $\alpha = 0.8$. Calculate one cycle-averaged current J_{avg} for all these values of T_p . Plot J_{avg} as function of T_p . What do you observe? Give a physical interpretation of the results.

Programming exercise 5.3

Now let $T_p = 500$, choose 50 equally spaced values of α between 0 and 1, and run the simulation for one cycle for each of these values of α . Let the number of particles in the system be $N_p = 12N_x$. Calculate the cycle-averaged current J_{avg} for all these values of α . Plot J_{avg} as function of α . What do you observe? Give a physical interpretation of the results.

6 Analytical solution

In the simple sawtooth-shaped potential which we encountered in the last section, there exists in fact an analytical solution to the current in certain parameter limits. Let us briefly go through the argument which leads to the solution:

When the particles find themselves in the sawtooth-shaped potential $V_1(x)$, they will relax towards the potential minima. Hence, as the particles are excited to the second energy level defined by $V_2(x)$, they start out localized at positions $x_n = nN_x$, $n \in \mathbb{Z}$. Since $V_2(x)$ is a constant potential, the particles will be subject to isotropic diffusion once excited to energy level 2. We learned in Sec. 2 that random walk in a flat potential corresponds to diffusion with constant diffusion coefficient. If the particle distribution at $t = 0$ is $\phi(x, t = 0) \sim \delta(x)$, where $\delta(x)$ is the Dirac delta function, one can show that the normalized solution of the diffusion equation at $t > 0$ is

$$\phi(x, t) = \frac{1}{\sqrt{4\pi Dt}} e^{-\frac{x^2}{4Dt}}, \quad (11)$$

where $D = \frac{dh^2}{2\Delta t}$ (all these quantities are defined in Sec. 2). Random walk in a flat potential corresponds to $d = 2/3$. We moreover have $h^2/\Delta t = 1$, so that $D = 1/3$. After diffusion with duration $t = T_p$, the particles are de-excited into $V_1(x)$, where the particles once again move down the potential gradient into one of the minima. The particles on the interval $-(1 - \alpha)N_x \leq x < \alpha N_x$ will therefore move towards $x = 0$. The particles having diffused to the right of αN_x will be caught in a minimum to the right, and the particles to the left of $-(1 - \alpha)N_x$ will be caught in a minimum to the left.

The number of particles having moved to the right after a cycle of length $2T_p$ can be found by using Eq. (11),

$$n^+ = N_p \sqrt{\frac{3}{4\pi T_p}} \int_{\alpha N_x}^{\infty} dx e^{-\frac{3x^2}{4T_p}}, \quad (12)$$

and the number of particles having moved to the left after the cycle of length $2T_p$ follows as

$$n^- = N_p \sqrt{\frac{3}{4\pi T_p}} \int_{-\infty}^{-(1-\alpha)N_x} dx e^{-\frac{3x^2}{4T_p}}. \quad (13)$$

The normalized, cycle-averaged particle current can therefore be expressed as

$$J_{\text{avg}} = \frac{N_x}{2T_p} \sqrt{\frac{3}{4\pi T_p}} \left(\int_{\alpha N_x}^{\infty} dx e^{-\frac{3x^2}{4T_p}} - \int_{-\infty}^{-(1-\alpha)N_x} dx e^{-\frac{3x^2}{4T_p}} \right). \quad (14)$$

Some algebra later, we arrive at

$$J_{\text{avg}} = \frac{N_x}{4T_p} \left[\text{erfc} \left(\frac{\alpha N_x}{2} \sqrt{\frac{3}{T_p}} \right) - \text{erfc} \left(\frac{(1-\alpha)N_x}{2} \sqrt{\frac{3}{T_p}} \right) \right], \quad (15)$$

where $\text{erfc}(x) = \frac{2}{\sqrt{\pi}} \int_x^{\infty} dz e^{-z^2}$ is the "complimentary error function".

Programming exercise 6.1

Compare the results of Ex. 5.3 with the analytical current [Eq. (15)] as function of α when $T_p = 500$. Plot both the numerical and analytical J_{avg} as function of α in the same figure. Compare and comment the graphs!

Programming exercise 6.2

Repeat Ex. 5.2, but this time with 20 values of T_p equally space between 80 and 1500, and use $\alpha = 0.8$ and $N_x = 10$ (so that the system is discretized into 20 spatial steps). We use these parameters in order to avoid a very long computation time. Compare the numerical results with the analytical current [Eq. (15)]. Plot both the numerical and analytical J_{avg} as function of T_p in the same figure. Compare and comment the graphs!

Programming exercise 6.3

Repeat programming exercise 6.1, but this time with $\beta k \in \{0.01, 1, 2, 3, 5, 10\}$. Compare the analytical and numerical results!

Exercise 6.4

Your results in 6.2 and 6.3 should indicate that the "analytical argument" presented above has some limitations. That is, the argument is not waterproof. Your last task in this exercise is to identify which parts of the argument contains implicit assumptions, and under what circumstances these assumptions do not hold!

Good luck!