

QCQI Chapter 2

Zhou Huanhai

February 3, 2021

Exercise 1

The exercises are automatically numbered, starting from one. Packages such as `amsmath` and `hyperref` are included by default.

Paragraphs are not indented, but are instead separated by some vertical space.

As an example: the *standard inner product* on \mathbb{R}^n is defined as

$$\vec{a} \cdot \vec{b} := x_1 y_1 + \cdots + x_n y_n \quad \text{for } \vec{a}, \vec{b} \in \mathbb{R}^n.$$

Note that `*` can be used instead of `\cdot`, and `\R` instead of `\mathbb{R}`. (For a normal asterisk, use `\ast`.) Of course, there are macros for the natural numbers etc. too. Commands such as `\abs{}` and `\Set{}` can be used to easily create (scaled) delimiters. For example,

$$\left| \frac{1}{1 - \lambda h} \right| \leq 1 \quad \text{and} \quad \left\{ x \in \mathbb{R} \mid 1 < \sqrt{x^3 + 2} < \frac{3}{2} \right\}.$$

The starred version of these commands disables the auto-scaling.

Exercise 2

A matrix representation of A is:

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

where we take $\{|0\rangle, |1\rangle\}$ as the input and output bases.

If we change the output bases to $\{|+\rangle, |-\rangle\}$ (input bases remains), note that $A|0\rangle = |1\rangle = \frac{|+\rangle - |-\rangle}{\sqrt{2}}$ and $A|1\rangle = |0\rangle = \frac{|+\rangle + |-\rangle}{\sqrt{2}}$, there is a different matrix representation:

$$\begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{bmatrix}$$

Exercise 3

For simplicity, take $\{|v_i\rangle\}$, $\{|w_j\rangle\}$ and $\{|x_k\rangle\}$ as the corresponding input and output bases.

Given that

$$\forall i. A|v_i\rangle = \sum_j A_{ji}|w_j\rangle$$

and

$$\forall j. B|w_j\rangle = \sum_k B_{kj}|x_k\rangle,$$

we have

$$\begin{aligned}\forall i. BA|v_i\rangle &= B \sum_j A_{ji}|w_j\rangle \\ &= \sum_j A_{ji}B|w_j\rangle \\ &= \sum_j \sum_k A_{ji}B_{kj}|x_k\rangle \\ &= \sum_k \sum_j B_{kj}A_{ji}|x_k\rangle\end{aligned}$$

Hence for the new operator BA , there is a matrix representation denoted as $(BA)_{\text{rank}(X) \times \text{rank}(V)}$, which satisfies:

$$\forall k, i. (BA)_{k,i} = \sum_k \sum_j B_{kj}A_{ji}$$

i.e., $(BA)_{\text{rank}(X) \times \text{rank}(V)}$ is the matrix product of $B_{\text{rank}(X) \times \text{rank}(W)}$ and $A_{\text{rank}(W) \times \text{rank}(V)}$.

Exercise 4

Let n denotes the rank of V and $\{|v_i\rangle\}$ as a set of bases of V .

Have the identity operator I written in a matrix form as follows:

$$\forall j = 0, 1, 2, \dots, n. I|v_j\rangle = |v_j\rangle = \sum_{i=0,1,\dots,n} k_{ij}|v_i\rangle$$

Assume that

$$\exists m \neq n. k_{mn} \neq 0,$$

There are only two possible cases: Either $\{|v_i\rangle\}$ are linearly dependent, or $|v_m\rangle = \vec{0}$,

which leads to contradiction with the fact that $\{|v_i\rangle\}$ forms a set of bases of V .

Therefore we have

$$\forall i \neq j. k_{ij} = 0$$

and hence

$$\forall i. k_{ii} = 1$$

So the one and only matrix representation of I is $\text{diag}(\underbrace{1, 1, 1, \dots, 1}_n)$.

Exercise 5

Let $\vec{x} = (x_1, \dots, x_n) \in \mathbb{C}^n$, $\vec{y} = (y_1, \dots, y_n) \in \mathbb{C}^n$ and $\vec{z} = (z_1, \dots, z_n) \in \mathbb{C}^n$.

Check the requirements of a inner product:

- linear in the second argument:

$$\begin{aligned} (\vec{x}, \lambda_y \vec{y} + \lambda_z \vec{z}) &= (\vec{x}, (\lambda_y y_1 + \lambda_z z_1, \dots, \lambda_y y_n + \lambda_z z_n)) \\ &= \sum_i x_i^* (\lambda_y y_i + \lambda_z z_i) \\ &= \lambda_y \sum_i x_i^* y_i + \lambda_z \sum_i x_i^* z_i \\ &= \lambda_y (\vec{x}, \vec{y}) + \lambda_z (\vec{x}, \vec{z}) \end{aligned}$$

- Note that $\forall a, b \in \mathbb{C}. (ab)^* = a^* b^* = b^* a^*$ and $a^* + b^* = (a + b)^*$,

then we have:

$$(\vec{x}, \vec{y}) = \sum_i x_i^* y_i = \sum_i (y_i^* x_i)^* = (\sum_i y_i^* x_i)^* = (\vec{y}, \vec{x})^*$$

- Note that $\forall a \in \mathbb{C}. a^* a \geq 0$ with equality if and only if $a = 0$,

then we have

$$(\vec{x}, \vec{x}) = \sum_i x_i^* x_i \geq 0,$$

with equality if and only if all $x_i = 0$, i.e., $\vec{x} = \vec{0}$.

So (\cdot, \cdot) is an inner product on \mathbb{C}^n .

Exercise 6

Let (\cdot, \cdot) be inner product from $V \times V$ to \mathbb{C} , and $|x\rangle, |y\rangle, |z\rangle \in V$.

Based on the second property (conjugate-symmetry) and apply linearity in the second argument, we have

$$\begin{aligned} (\lambda_x |x\rangle + \lambda_y |y\rangle, |z\rangle) &= (|z\rangle, \lambda_x |x\rangle + \lambda_y |y\rangle)^* \\ &= (\lambda_x (|z\rangle, |x\rangle) + \lambda_y (|z\rangle, |y\rangle))^* \\ &= \lambda_x^* (|z\rangle, |x\rangle)^* + \lambda_y^* (|z\rangle, |y\rangle)^* \\ &= \lambda_x^* (|x\rangle, |z\rangle) + \lambda_y^* (|y\rangle, |z\rangle) \end{aligned}$$

i.e., an inner product is conjugate-linear in the first argument.

Exercise 7

Let $\{|0\rangle, |1\rangle\}$ denotes the orthonormal bases of the vector representation.

We have

$$\begin{aligned} \langle w|v\rangle &= (\langle 0| + \langle 1|)(|0\rangle - |1\rangle) \\ &= \langle 0|0\rangle - \langle 1|1\rangle + \langle 1|0\rangle - \langle 0|1\rangle \\ &= 1 - 1 + 0 - 0 \\ &= 0 \end{aligned}$$

So $|w\rangle$ and $|v\rangle$ are orthogonal.

Their normalized forms are $|w'\rangle = \frac{|w\rangle}{\|w\rangle} = (\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$ and $|v'\rangle = \frac{|v\rangle}{\|v\rangle} = (\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2})$.

Exercise 8

Let $\{|w_i\rangle\}$ denotes the original basis and $\{|v_i\rangle\}$ as the basis given by the Gram-Schmidt procedure.

According to definition, it is clear that they are normalized, i.e.,

$$\forall i. \langle v_i|v_i\rangle = \|v_i\| = 1$$

Note that the inner product is conjugate-symmetry, we just need to check whether for all $1 \leq k \leq d-1$ and $1 \leq l \leq k$, $|v_{k+1}\rangle$ and $|v_l\rangle$ are orthogonal. And for simplicity we check numerators of the Gram-Schmidt form.

- $|v_2\rangle$ and $|v_1\rangle$ are orthogonal because

$$(\langle w_2| - \langle v_1|w_2\rangle \langle v_1|)|v_1\rangle = \langle w_2|v_1\rangle - \langle v_1|w_2\rangle \langle v_1|v_1\rangle = 0$$

- Assume that for all $1 \leq i \leq k$ ($2 \leq k \leq d-1$), $\{|v_i\rangle\}$ are orthonormal, then we have

$$\begin{aligned} \forall 1 \leq i \leq k. (\langle w_{k+1}| - \sum_{j=1}^k \langle v_j|w_{k+1}\rangle \langle v_j|)|v_i\rangle &= \langle w_{k+1}|v_i\rangle - \sum_{j=1}^k \langle v_j|w_{k+1}\rangle \delta_{ij} \\ &= \langle w_{k+1}|v_i\rangle - \langle v_i|w_{k+1}\rangle \\ &= 0 \end{aligned}$$

i.e., $\{|v_i\rangle\} \cup \{|v_{k+1}\rangle\}$ are orthonormal.

By induction we prove that $\{|v_i\rangle\}$ is an orthonormal basis.

Exercise 9

- $X = |0\rangle\langle 1| + |1\rangle\langle 0|$
- $Y = -i|0\rangle\langle 1| + i|1\rangle\langle 0|$
- $Z = |0\rangle\langle 0| - |1\rangle\langle 1|$

Exercise 10

Let A be the matrix representation for the operator $|v_j\rangle\langle v_k|$, we have

$$\forall m \neq j \text{ or } n \neq k. \langle v_m|A|v_n\rangle = 0$$

Note that $A|v_n\rangle$ is a linear combination of $\{|v_i\rangle\}$, hence $\langle v_m|A|v_n\rangle = 0$ if and only if $A_{mn} = 0$.

Furthermore,

$$\langle v_j|A|v_k\rangle = A_{jk} \langle v_j|v_k\rangle = A_{jk} = 1$$

So A is a $\text{rank}(V) \times \text{rank}(V)$ matrix, with $A_{jk} = 1$ and other entries set to 0.

Exercise 11

- $\det|X - \lambda I| = \lambda^2 - 1 = 0 \Rightarrow \lambda_{1,2} = \pm 1$
 $\lambda_1 = 1: (X - I)|x\rangle = 0 \Rightarrow$ a normalized eigenvector $|x_1\rangle = \frac{\sqrt{2}}{2}(1, 1)$
 $\lambda_2 = -1: (X + I)|x\rangle = 0 \Rightarrow$ a normalized eigenvector $|x_2\rangle = \frac{\sqrt{2}}{2}(1, -1)$

Hence a diagonal representation is $X = |x_1\rangle\langle x_1| - |x_2\rangle\langle x_2|$

- $\det|Y - \lambda I| = \lambda^2 - 1 = 0 \Rightarrow \lambda_{1,2} = \pm 1$
 $\lambda_1 = 1: (Y - I)|y\rangle = 0 \Rightarrow$ a normalized eigenvector $|y_1\rangle = \frac{\sqrt{2}}{2}(1, i)$
 $\lambda_2 = -1: (Y + I)|y\rangle = 0 \Rightarrow$ a normalized eigenvector $|y_2\rangle = \frac{\sqrt{2}}{2}(1, -i)$

Hence a diagonal representation is $Y = |y_1\rangle\langle y_1| - |y_2\rangle\langle y_2|$

- $\det|Z - \lambda I| = \lambda^2 - 1 = 0 \Rightarrow \lambda_{1,2} = \pm 1$
 $\lambda_1 = 1: (Z - I)|z\rangle = 0 \Rightarrow$ a normalized eigenvector $|z_1\rangle = (1, 0)$
 $\lambda_2 = -1: (Z + I)|z\rangle = 0 \Rightarrow$ a normalized eigenvector $|z_2\rangle = (0, 1)$

Hence a diagonal representation is $Z = |z_1\rangle\langle z_1| - |z_2\rangle\langle z_2|$

Exercise 12

$$\det\left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} - \lambda I\right) = (1 - \lambda)^2 = 0$$

We get the only eigenvalue $\lambda = 1$.

Let $(A - I)|x\rangle = 0$, then get an eigenvector $|x\rangle = (0, 1)$ and the rank of the eigenspace w.r.t. $\lambda = 1$ is 1.

Note that

$$k|x\rangle\langle x| = \begin{bmatrix} 0 & 0 \\ 0 & k \end{bmatrix} \neq \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Hence the matrix is not diagonalizable.

Exercise 13

Suppose $|w\rangle \in W$ and $|v\rangle \in V$.

Let operator $A = |w\rangle\langle v|$ and $B = |v\rangle\langle w|$. It is not difficult to see that A maps from V to W and B maps from W to V .

$$\forall |i\rangle \in V. (A|i\rangle)^\dagger = \langle v|i\rangle^* \langle w| = \langle i|v\rangle \langle w| = \langle i|A^\dagger = \langle i|B$$

Hence $B = A^\dagger$, i.e., $(|w\rangle\langle v|)^\dagger = |v\rangle\langle w|$.

Exercise 14

Suppose A_i maps from V to W .

For all $|v\rangle \in V, |w\rangle \in W$,

$$\begin{aligned} ((\sum_i a_i A_i)^\dagger |v\rangle, |w\rangle) &= (|v\rangle, \sum_i a_i A_i |w\rangle) \\ &= \sum_i a_i (|v\rangle, A_i |w\rangle) \\ &= \sum_i a_i (A_i^\dagger |v\rangle, |w\rangle) \\ &= ((\sum_i a_i^* A_i^\dagger) |v\rangle, |w\rangle) \end{aligned}$$

Hence $(\sum_i a_i A_i)^\dagger = \sum_i a_i^* A_i^\dagger$.

Exercise 15

Suppose A maps from V to W .

For all $|v\rangle \in V, |w\rangle \in W$,

$$((A^\dagger)^\dagger |v\rangle, |w\rangle) = (|v\rangle, A^\dagger |w\rangle) = (A^\dagger |w\rangle, |v\rangle)^* = (|w\rangle, A |v\rangle)^* = (A |v\rangle, |w\rangle)$$

Hence $(A^\dagger)^\dagger = A$.

Exercise 16

Suppose P maps from V to its subspace W :

$$P = \sum_{i=1}^k |i\rangle \langle i|,$$

where $\{|i\rangle\}$ is an orthonormal basis for W .

$$\begin{aligned} \forall |v\rangle \in V, P^2 &= \sum_i |i\rangle \langle i| \sum_j |j\rangle \langle j| \\ &= \sum_{i,j} |i\rangle \langle i|j\rangle \langle j| \\ &= \sum_{i,j} \delta_{ij} |i\rangle \langle j| \\ &= \sum_i |i\rangle \langle i| \\ &= P \end{aligned}$$

Exercise 17

• \Rightarrow :

For any Hermitian matrix A , suppose it has an eigenvalue λ and a corresponding eigenvector $|x\rangle$.

We have

$$\langle x| A |x\rangle = \langle x| \lambda |x\rangle = \lambda \langle x|x\rangle$$

On the other side,

$$\langle x| A |x\rangle = \langle x| A^\dagger |x\rangle = \langle x| A |x\rangle^* = \lambda^* \langle x|x\rangle$$

Note that $\langle x|x\rangle > 0$, so $\lambda = \lambda^*$. i.e., λ is real.

• \Leftarrow :

Let A be any normal matrix with real eigenvalues. As it is normal, it can be diagonalized as follows:

$$A = \sum_i \lambda_i |i\rangle \langle i|$$

where $\{|i\rangle\}$ are eigenvectors corresponding to eigenvalues $\{\lambda_i\}$.

(Notice that the rank of an eigenspace may be greater than 1, but it does not matter.)

Given that λ_i is real,

$$A^\dagger = (A^*)^T = \sum_i \lambda_i^* |i\rangle \langle i| = \sum_i \lambda_i |i\rangle \langle i| = A$$

i.e., A is Hermitian.

Combining the two parts above, a normal matrix is Hermitian if and only if it has real eigenvalues.

Exercise 18

Let U be a unitary matrix, and λ as any of its eigenvalues. Then we have

$$(U|v\rangle, U|w\rangle) = (\lambda|v\rangle, \lambda|w\rangle) = \lambda\lambda^*(|v\rangle, |w\rangle)$$

Also by definition

$$(U|v\rangle, U|w\rangle) = (|v\rangle, |w\rangle)$$

Hence

$$\lambda\lambda^* = 1$$

Note that $\|\lambda\| \|\lambda^*\| = 1$ and $\|\lambda\| = \|\lambda^*\| \geq 0$, so $\|\lambda\| = 1$.

In conclusion, all eigenvalues of U have modulus 1.

Exercise 19

- I : It is clear that I is Hermitian and unitary.
- X :

$$X^\dagger = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = X$$

$$X^\dagger X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

- Y :

$$Y^\dagger = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} = Y$$

$$Y^\dagger Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

- Z :

$$Z^\dagger = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = Z$$

$$Z^\dagger Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Exercise 20

Let $U = \sum_i |w_i\rangle\langle v_i|$.

Note that $|v_i\rangle$ and $|w_i\rangle$ are both orthonormal bases, hence the operator U is unitary.

$$\begin{aligned}
A'_{ij} &= \langle v_i | A | v_j \rangle = \langle v_i | U U^\dagger A U U^\dagger | v_j \rangle \\
&= \sum_k \langle v_i | w_k \rangle \langle v_k | U^\dagger A U \sum_l \langle w_l | v_j \rangle | v_l \rangle \\
&= \sum_k \langle v_i | w_k \rangle \langle v_k | \sum_m | v_m \rangle \langle w_m | A \sum_n | w_n \rangle \langle v_n | \sum_l \langle w_l | v_j \rangle | v_l \rangle \\
&= \sum_{k,l} \langle v_i | w_k \rangle \langle w_l | v_j \rangle \sum_{m,n} \delta_{km} A''_{mn} \delta_{nl} \\
&= \sum_{k,l} \langle v_i | w_k \rangle \langle w_l | v_j \rangle A''_{kl}
\end{aligned}$$

which characterizes the relationship between A' and A'' .

Exercise 21

- \Rightarrow : We also prove by induction on the dimension d of V . The case $d = 1$ is also trivial.

Let λ be an eigenvalue of M , P the projector onto the λ eigenspace, and Q the projector onto the orthogonal complement. Then

$$M = (P + Q)M(P + Q) = PMP + QMP + PMQ + QMQ$$

Note that $QMP = 0$ as M takes the subspace P into itself. Also we have

$$PMQ = (QM^\dagger P)^\star = (QMP)^\star = 0$$

Next we prove that QMQ is Hermitian:

$$(QMQ)^\dagger = QM^\dagger Q = QMQ$$

By induction, QMQ is diagonal w.r.t. some orthonormal basis for the subspace Q , and PMP is already diagonal w.r.t. some orthonormal basis for the λ eigenspace.

It follows that $M = PMP + QMQ$ is also diagonal, w.r.t. some orthonormal basis for V .

- \Leftarrow : It turns out that the statement does not hold conversely, in the Hermitian case.

Exercise 22

Let A be any Hermitian operator with eigenvalues λ_1 and λ_2 , and corresponding eigenvectors $|\lambda_1\rangle$ and $|\lambda_2\rangle$. (λ_1 and λ_2 are real as A is Hermitian.)

Note that $A|\lambda_2\rangle = \lambda_2|\lambda_2\rangle$ and $\langle\lambda_1|A^\dagger = \lambda_1^\star\langle\lambda_1| = \lambda_1\langle\lambda_1|$,

$$\langle\lambda_1|A|\lambda_2\rangle = \lambda_2\langle\lambda_1|\lambda_2\rangle$$

Also we have

$$\langle\lambda_1|A|\lambda_2\rangle = \langle\lambda_1|A^\dagger|\lambda_2\rangle = \lambda_1\langle\lambda_1|\lambda_2\rangle$$

Combine the two equations above we get

$$(\lambda_1 - \lambda_2)\langle\lambda_1|\lambda_2\rangle = 0$$

As $\lambda_1 \neq \lambda_2$, $|\lambda_1\rangle$ and $|\lambda_2\rangle$ are necessarily orthogonal.

Exercise 23

For simplicity we use P to denote both the projector and the associated subspace.

Let V be the total vector space and Q the orthogonal complement of P .

For any non-zero vector $|v\rangle \in V$, it could be written in the following form:

$$|v\rangle = |v_P\rangle + |v_Q\rangle$$

where $|v_P\rangle \in P$ and $|v_Q\rangle \in Q$.

If $P|v\rangle = \lambda|v\rangle$ holds for some λ , we have

$$\begin{aligned} P|v\rangle &= P|v_P\rangle + P|v_Q\rangle \\ &= |v_P\rangle \\ &= \lambda|v_P\rangle + \lambda|v_Q\rangle \end{aligned}$$

Hence

$$(\lambda - 1)|v_P\rangle + \lambda|v_Q\rangle = \vec{0}$$

From exercise 2.22 we know that $|v_P\rangle$ and $|v_Q\rangle$ are orthonormal and they cannot be zero vectors at the same time, so there are only two possible cases:

- $\lambda = 0$: In this case $|v\rangle \in Q$
- $\lambda = 1$: In this case $|v\rangle \in P$

In conclusion, the eigenvalues of P are all either 0 or 1.

Exercise 24

For any positive operator A , we have

$$A = \frac{A + A^\dagger}{2} + i \frac{A - A^\dagger}{2i}$$

Let $B = \frac{A + A^\dagger}{2}$ and $C = \frac{A - A^\dagger}{2i}$, and it is clear that they are both Hermitian. Hence we have

$$\forall |v\rangle. \langle v|A|v\rangle = \langle v|B + iC|v\rangle = \langle v|B|v\rangle + i\langle v|C|v\rangle \geq 0$$

Note that B and C are Hermitian, $\langle v|B|v\rangle$ and $\langle v|C|v\rangle$ should both be real. Since the equation above holds for all $|v\rangle$, C must be a zero operator. Therefore

$$A = A^\dagger$$

i.e., A is Hermitian.

Exercise 25

For any $|v\rangle$, suppose A maps from $|v\rangle$ to $|w\rangle$.

We have

$$(|v\rangle, A^\dagger A|v\rangle) = \langle v|A^\dagger A|v\rangle = \langle w|w\rangle \geq 0$$

By definition $A^\dagger A$ is positive.

Exercise 26

•

$$|\psi\rangle^{\otimes 2} = \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right) \otimes \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right) = \frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle + \frac{1}{2} |10\rangle + \frac{1}{2} |11\rangle$$

$$|\psi\rangle^{\otimes 2} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \otimes \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

•

$$\begin{aligned} |\psi\rangle^{\otimes 3} &= \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right) \otimes \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right) \otimes \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}}\right) \\ &= \frac{\sqrt{2}}{4} |000\rangle + \frac{\sqrt{2}}{4} |001\rangle + \frac{\sqrt{2}}{4} |010\rangle + \frac{\sqrt{2}}{4} |011\rangle + \frac{\sqrt{2}}{4} |100\rangle + \frac{\sqrt{2}}{4} |101\rangle \\ &\quad + \frac{\sqrt{2}}{4} |110\rangle + \frac{\sqrt{2}}{4} |111\rangle \end{aligned}$$

$$|\psi\rangle^{\otimes 3} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \otimes \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \otimes \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} = \frac{\sqrt{2}}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

Exercise 27

•

$$X \otimes Z = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}$$

•

$$I \otimes X = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

•

$$X \otimes I = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

Note that $I \otimes X \neq X \otimes I$, hence the tensor product is not commutative, in general.

Exercise 28

- Distribution of transposition:

$$(A \otimes B)^T = \begin{bmatrix} A_{11}B^T & A_{21}B^T & \cdots & A_{m1}B^T \\ A_{12}B^T & A_{22}B^T & \cdots & A_{m2}B^T \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n}B^T & A_{2n}B^T & \cdots & A_{mn}B^T \end{bmatrix} = \begin{bmatrix} A_{11} & A_{21} & \cdots & A_{m1} \\ A_{12} & A_{22} & \cdots & A_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \cdots & A_{mn} \end{bmatrix} \otimes B^T = A^T \otimes B^T$$

- Distribution of complex conjugation:

$$(A \otimes B)^* = \begin{bmatrix} A_{11}^*B^* & A_{12}^*B^* & \cdots & A_{1n}^*B^* \\ A_{21}^*B^* & A_{22}^*B^* & \cdots & A_{2n}^*B^* \\ \vdots & \vdots & \ddots & \vdots \\ A_{m1}^*B^* & A_{m2}^*B^* & \cdots & A_{mn}^*B^* \end{bmatrix} = \begin{bmatrix} A_{11}^* & A_{12}^* & \cdots & A_{1n}^* \\ A_{21}^* & A_{22}^* & \cdots & A_{2n}^* \\ \vdots & \vdots & \ddots & \vdots \\ A_{m1}^* & A_{m2}^* & \cdots & A_{mn}^* \end{bmatrix} \otimes B^* = A^* \otimes B^*$$

- Distribution of adjoint operation:

Based on the two distribution laws above, we have

$$(A \otimes B)^\dagger = ((A \otimes B)^*)^T = ((A^* \otimes B^*)^T = A^\dagger \otimes B^\dagger$$

Exercise 29

Let $U_1 : V \rightarrow V'$ be a unitary operator and $U_2 : W \rightarrow W'$ as another unitary operator.

Based on exercise 2.28 we know that $(U_1 \otimes U_2)^\dagger = U_1^\dagger \otimes U_2^\dagger$. So

$$(U_1 \otimes U_2)^\dagger (U_1 \otimes U_2) = U_1^\dagger U_1 \otimes U_2^\dagger U_2 = I_1 \otimes I_2 = I_{V \otimes W}$$

Similarly we have $(U_1 \otimes U_2)(U_1 \otimes U_2)^\dagger = I_{V' \otimes W'}$, i.e., $U_1 \otimes U_2$ is also unitary.

Exercise 30

Let A be a Hermitian operator on V and B as another Hermitian operator on W .

Based on exercise 2.28 we know that

$$(A \otimes B)^\dagger = A^\dagger \otimes B^\dagger = A \otimes B$$

i.e., $A \otimes B$ is also Hermitian.

Exercise 31

Let A be a positive operator on V and B as another positive operator on W .

Note that $|v_i\rangle \otimes |w_j\rangle$ forms an orthonormal basis for $V \otimes W$, hence for any $|u\rangle$ in $V \otimes W$, it is a linear combination of $\{|v_i\rangle \otimes |w_j\rangle\}$:

$$|u\rangle = \sum_{k,l} a_{kl} |v_k\rangle \otimes |w_l\rangle$$

Consider the natural inner product on $V \otimes W$ and apply linearity:

$$\begin{aligned}(|u\rangle, (A \otimes B)|u\rangle) &= \left(\sum_{k,l} a_{kl} |v_k\rangle \otimes |w_l\rangle, \sum_{k,l} a_{kl} A|v_k\rangle \otimes B|w_l\rangle\right) \\ &= \sum_{k,l} \|a_{kl}\|^2 \langle v_k| A|v_k\rangle \langle w_l| B|w_l\rangle \\ &\geq 0\end{aligned}$$

Hence $A \otimes B$ is also positive.

Exercise 32

Let $P_V = \sum_i |v_i\rangle\langle v_i|$ be a projector on V and $P_W = \sum_j |w_j\rangle\langle w_j|$ as another projector on W . For simplicity we have P_V and P_W also denote the corresponding subspace.

It is clear that

$$P_V \otimes P_W = \sum_{i,j} |v_i\rangle \otimes |w_j\rangle \langle v_i| \otimes \langle w_j|$$

Note that $|v_i\rangle \otimes |w_j\rangle$ forms a orthonormal basis for $V \otimes W$, therefore $P_V \otimes P_W$ is also a projector on $V \otimes W$, w.r.t. the subspace $P_V \otimes P_W$.

Exercise 33

We prove the statement inductively.

- When $n = 1$, it holds that

$$H^{\otimes 1} = \frac{1}{\sqrt{2}}(|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| - |1\rangle\langle 1|) = \frac{1}{\sqrt{2}} \sum_{x_1, y_1=0,1} (-1)^{x_1 \cdot y_1} |x_1\rangle\langle y_1|$$

- Assume that for any $n \geq 1$,

$$H^{\otimes n} = \frac{1}{\sqrt{2^n}} \sum_{x_n, y_n=0,1,\dots,2^n-1} (-1)^{x_n \cdot y_n} |x_n\rangle\langle y_n|$$

Note that every time we tensor with H , the basis changes from $|x_n\rangle$ to $|x_{n+1}\rangle$, with $|i\rangle \leftarrow |i\rangle \otimes |0\rangle$ and $|i + 2^n\rangle \leftarrow |i\rangle \otimes |1\rangle$ for all $0 \leq i < 2^n$.

Hence we have

$$H^{\otimes n+1} = H^{\otimes n} \otimes H = \frac{1}{\sqrt{2^{n+1}}} \sum_{x_{n+1}, y_{n+1}=0,1,\dots,2^{n+1}-1} (-1)^{x_{n+1} \cdot y_{n+1}} |x_{n+1}\rangle\langle y_{n+1}|$$

By induction we can prove that the equation of $H^{\otimes n}$ always holds.

The matrix representation for $H^{\otimes 2}$ is

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

Exercise 34

Let $A = \begin{bmatrix} 4 & 3 \\ 3 & 4 \end{bmatrix}$.

$$\det(A - \lambda I) = 0 \Rightarrow \lambda_1 = 1, \lambda_2 = 7$$

The corresponding normalized eigenvectors are

$$|\lambda_1\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$|\lambda_2\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Then

$$A = |\lambda_1\rangle\langle\lambda_1| + 7|\lambda_2\rangle\langle\lambda_2|$$

The square root of A is

$$\sqrt{A} = |\lambda_1\rangle\langle\lambda_1| + \sqrt{7}|\lambda_2\rangle\langle\lambda_2| = \frac{1}{2} \begin{bmatrix} 1 + \sqrt{7} & -1 + \sqrt{7} \\ -1 + \sqrt{7} & 1 + \sqrt{7} \end{bmatrix}$$

The logarithm of A is

$$\log(A) = \log(1)|\lambda_1\rangle\langle\lambda_1| + \log(7)|\lambda_2\rangle\langle\lambda_2| = \frac{\log(7)}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Exercise 35

By definition we get

$$\vec{v} \cdot \vec{\sigma} = \begin{bmatrix} v_3 & v_1 - iv_2 \\ v_1 + iv_2 & -v_3 \end{bmatrix}$$

Note that \vec{v} is a unit vector,

$$\begin{aligned} \det(\vec{v} \cdot \vec{\sigma} - \lambda I) &= 0 \Rightarrow \lambda^2 = v_1^2 + v_2^2 + v_3^2 = 1 \\ &\Rightarrow \lambda_{1,2} = \pm 1 \end{aligned}$$

Note that $\vec{v} \cdot \vec{\sigma}$ is Hermitian, hence it could be diagonalized as follows:

$$\vec{v} \cdot \vec{\sigma} = |\lambda_1\rangle\langle\lambda_1| - |\lambda_2\rangle\langle\lambda_2|$$

where $|\lambda_{1,2}\rangle$ are the corresponding normalized eigenvectors. (Do not need to explicitly solve them since we already know that $\vec{v} \cdot \vec{\sigma}$ is diagonalizable.)

Furthermore,

$$|\lambda_1\rangle\langle\lambda_1| - |\lambda_2\rangle\langle\lambda_2| = I$$

Based on the definition of operator functions and applying Euler's equation, we have

$$\begin{aligned} \exp(i\theta \vec{v} \cdot \vec{\sigma}) &= e^{i\theta} |\lambda_1\rangle\langle\lambda_1| + e^{-i\theta} |\lambda_2\rangle\langle\lambda_2| \\ &= (\cos \theta + i \sin \theta) |\lambda_1\rangle\langle\lambda_1| + (\cos \theta - i \sin \theta) |\lambda_2\rangle\langle\lambda_2| \\ &= \cos \theta (|\lambda_1\rangle\langle\lambda_1| + |\lambda_2\rangle\langle\lambda_2|) + i \sin \theta (|\lambda_1\rangle\langle\lambda_1| - |\lambda_2\rangle\langle\lambda_2|) \\ &= \cos \theta I + i \sin \theta \vec{v} \cdot \vec{\sigma} \end{aligned}$$

Exercise 36

- $\text{tr}(X) = 0$
- $\text{tr}(Y) = 0$
- $\text{tr}(Z) = 1 - 1 = 0$

Exercise 37

Suppose both A and B act on V which has an orthonormal basis $|v_i\rangle$.

$$\begin{aligned}\text{tr}(AB) &= \sum_i \langle v_i | AB | v_i \rangle \\ &= \sum_i \langle v_i | A B | v_i \rangle \\ &= \sum_{i,j} \langle v_i | A | v_j \rangle \langle v_j | B | v_i \rangle \\ &= \sum_{i,j} \langle v_j | B | v_i \rangle \langle v_i | A | v_j \rangle \\ &= \sum_j \langle v_j | BA | v_j \rangle \\ &= \text{tr}(BA)\end{aligned}$$

Exercise 38

Suppose both A and B act on V which has an orthonormal basis $|v_i\rangle$.

$$\begin{aligned}\text{tr}(A + B) &= \sum_i \langle v_i | A + B | v_i \rangle \\ &= \sum_i \langle v_i | A | v_i \rangle + \sum_i \langle v_i | B | v_i \rangle \\ &= \text{tr}(A) + \text{tr}(B)\end{aligned}$$

Note that an inner product is linear in its second argument,

$$\begin{aligned}\text{tr}(zA) &= \sum_i \langle v_i | zA | v_i \rangle \\ &= z \sum_i \langle v_i | A | v_i \rangle \\ &= z \text{tr}(A)\end{aligned}$$

Exercise 39

39.1

Let A, B, C be operators in L_V .

- Based on exercise 2.38 we have

$$\begin{aligned}
(A, z_b B + z_c C) &= \text{tr}(A^\dagger(z_b B + z_c C)) \\
&= \text{tr}(z_b A^\dagger B) + \text{tr}(z_c A^\dagger C) \\
&= z_b \text{tr}(A^\dagger B) + z_c \text{tr}(A^\dagger C) \\
&= z_b(A, B) + z_c(A, C)
\end{aligned}$$

where z_b and z_c are arbitrary complex numbers.

i.e., (\cdot, \cdot) is linear in its second argument.

- Conjugate-symmetry:

$$(A, B) = \text{tr}(A^\dagger B) = \sum_i \langle i | A^\dagger B | i \rangle = \left(\sum_i \langle i | B^\dagger A | i \rangle \right)^* = (B, A)^*$$

•

$$\begin{aligned}
(A, A) &= \text{tr}(A^\dagger A) = \sum_i \langle i | A^\dagger A | i \rangle \\
&= \sum_i \langle i | A^\dagger I A | i \rangle \\
&= \sum_i \langle i | A^\dagger \left(\sum_j | j \rangle \langle j | \right) A | i \rangle \\
&= \sum_{i,j} \langle i | A^\dagger | j \rangle \langle j | A | i \rangle \\
&= \sum_{i,j} \langle j | A | i \rangle^* \langle j | A | i \rangle \\
&= \sum_i \|\langle i | A | i \rangle\|^2 \\
&\geq 0
\end{aligned}$$

with equality if and only if A is a zero operator.

39.2

Let $|i\rangle$ be an orthonormal basis for V .

It is clear that $|i\rangle\langle j|$ are linearly independent. And for any operator in L_V , it could be written in the outer product form, i.e., linear combination of $|i\rangle\langle j|$.

Hence $|i\rangle\langle j|$ forms a basis for L_V . So

$$\text{rank}(L_V) = \|\{|i\rangle\langle j|\}\| = d^2$$

39.3

Note that the basis given above is countable, we get a set of basis namely $|l_k\rangle$, where

$$|l_k\rangle \equiv |i\rangle\langle j| \quad (i = \lfloor k/d \rfloor - 1, j = k \bmod n)$$

Apply the Gram-Schmidt procedure:

- Define $|l'_1\rangle = \frac{|l_1\rangle}{\|l_1\rangle}$
- For $1 \leq k \leq d^2 - 1$ define $|l'_{k+1}\rangle$ inductively by

$$|l'_{k+1}\rangle = \frac{|l_{k+1}\rangle - \sum_{i=1}^k \langle l'_i | l_{k+1} \rangle |l'_i\rangle}{\| |l_{k+1}\rangle - \sum_{i=1}^k \langle l'_i | l_{k+1} \rangle |l'_i\rangle \|}$$

The vectors $|l'_1\rangle, |l'_2\rangle, \dots, |l'_{d^2}\rangle$ form an orthonormal basis for L_V .