
Exercise 2.1 Prove the “homogeneous Nullstellensatz,” which says if $\mathfrak{a} \subseteq S$ is a homogeneous ideal, and if $f \in S$ is a homogeneous polynomial with $\deg f > 0$, such that $f(P) = 0$ for all $P \in Z(\mathfrak{a})$ in \mathbf{P}^n , then $f^q \in \mathfrak{a}$ for some $q > 0$.

Solution:

□

Exercise 2.2 For a homogeneous ideal $\mathfrak{a} \subseteq S$, show that the following conditions are equivalent:

- (i.) $Z(\mathfrak{a}) = \emptyset$ (the empty set);
 - (ii.) $\sqrt{\mathfrak{a}} =$ either S or the ideal $S_+ = \bigoplus_{d>0} S_d$;
 - (iii.) $\mathfrak{a} \supseteq S_d$ for some $d > 0$.
-

Solution:

□

Exercise 2.3

- (a) If $T_1 \subseteq T_2$ are subsets of S^h , then $Z(T_1) \supseteq Z(T_2)$.
 - (b) If $Y_1 \subseteq Y_2$ are subsets of \mathbf{P}^n , then $I(Y_1) \supseteq I(Y_2)$.
 - (c) For any two subsets Y_1, Y_2 of \mathbf{P}^n , $I(Y_1 \cup Y_2) = I(Y_1) \cap I(Y_2)$.
 - (d) If $\mathfrak{a} \subseteq S$ is a homogeneous ideal with $Z(\mathfrak{a}) \neq \emptyset$, then $I(Z(\mathfrak{a})) = \sqrt{\mathfrak{a}}$.
 - (e) For any subset $Y \subseteq \mathbf{P}^n$, $Z(I(Y)) = \overline{Y}$.
-

Solution:

□

Exercise 2.4

- (a) There is a one-to-one inclusion-reversing correspondence between algebraic sets in \mathbf{P}^n and homogeneous radical ideals of S not equal to S_+ given by $Y \mapsto I(Y)$ and $\mathfrak{a} \mapsto Z(\mathfrak{a})$. *Note:* Since S_+ does not occur in this correspondence, it is sometimes called the *irrelevant* maximal ideal of S .
 - (b) An algebraic set $Y \subseteq \mathbf{P}^n$ is irreducible if and only if $I(Y)$ is a prime ideal.
 - (c) Show that \mathbf{P}^n itself is irreducible.
-

Solution:

□

Exercise 2.5

- (a) \mathbf{P}^n is a noetherian topological space.
 - (b) Every algebraic set in \mathbf{P}^n can be written uniquely as a finite union of irreducible algebraic sets, no one containing another. These are called its *irreducible components*.
-

Solution:

□