

The Math of the Shortprint Application

Terence C.

This is a high-level overview of the techniques used in the shortprint application for the user who wants to know precisely what is being computed for the sake of accurate inferences. There are many hyperlinks that may not be displayed in a preview version of this pdf, but they should be visible and clickable upon download. The urls will also be listed at the end for the reader who does not want to download the pdf. Some statistical background will be assumed throughout.

Notation

For simplicity, when we say *cards*, we will only be referring to the cards pulled that are of the rarity that the user cares about. We will let n denote the total number of cards pulled (of that rarity) and k denote the different number of card names (in that rarity). For simplicity we will label these cards:

$$\{1, 2, 3, \dots, k\} \tag{1}$$

and write $[k]$ for this set of labels. For $i \in [k]$, we will write x_i for the number of card i pulled (which we will view as a random variable). We will also write p_i for the “true” (unknown to us) probability of seeing card i each time a card of that rarity is pulled. So if there is no shortprinting, then $p_i = \frac{1}{k}$ for all i . Finally, we will use the symbol $\mathbb{P}$ to denote probability.

Option 1

Computing the p -value

For this computation only, we will use the normal approximation to the binomial distribution. To calculate the p value, we assume that all $p_i = \frac{1}{k}$ and ask how likely it is that our observed ratios $\{x_i/k\}_{i \in [k]}$ are as different

from $\frac{1}{k}$ as we observe. Under this hypothesis, we have:

$$\sum_{i=1}^k \frac{(x_i - n/k)^2}{n/k} \sim \chi_{k-1}^2 \quad (2)$$

(Pearson's chi-square test); the p -value we display is accordingly the probability that a χ_{k-1}^2 random variable is at least as large as the left-hand side of (2).

The individual confidence intervals

For the individual confidence intervals, we will use the Clopper-Pearson method. Consider the following functions:

$$f^+(p, m) := \sum_{j=m}^n \binom{n}{j} p^j (1-p)^{n-j} \quad (3)$$

$$f^-(p, m) := \sum_{j=0}^m \binom{n}{j} p^j (1-p)^{n-j} \quad (4)$$

for p a real number in the closed interval $[0, 1]$ and m an integer between 0 and n (inclusive). So for each card i , and integer m between 0 and n , we may express:

$$\mathbb{P}[x_i \geq m] = f^+(p_i, x_i), \quad \mathbb{P}[x_i \leq m] = f^-(p_i, x_i). \quad (5)$$

If we hold p fixed, $f^+(p, m)$ is a strictly decreasing function of m , so

$$\mathbb{P}[f^+(p_i, x_i) \leq f^+(p_i, m)] = \mathbb{P}[x_i \geq m] = f^+(p_i, m). \quad (6)$$

And if β is such that $f^+(p_i, m) \leq \beta < f^+(p_i, m-1)$, then we also have

$$\mathbb{P}[f^+(p_i, x_i) \leq \beta] = \mathbb{P}[x_i \geq m] = f^+(p_i, m) \quad (7)$$

(which is at most β by assumption). Altogether, we see that for all β in the closed interval $[0, 1]$:

$$\mathbb{P}[f^+(p_i, x_i) \leq \beta] \leq \beta. \quad (8)$$

Using the fact that $f^-(p, m)$ is strictly increasing in m (when p is held fixed), we have

$$\mathbb{P}[f^-(p_i, x_i) \leq f^-(p_i, m)] = \mathbb{P}[x_i \leq m] = f^-(p_i, m) \quad (9)$$

which also the probability that $f^-(p_i, x_i) \leq \beta$ for β such that $f^-(p_i, m) \leq \beta < f^-(p_i, m+1)$. So for all β in the closed interval $[0, 1]$:

$$\mathbb{P}[f^-(p_i, x_i) \leq \beta] \leq \beta. \quad (10)$$

Take α to be the desired confidence (for example, $\alpha = .95$); for each card i , we have by (8) and (10):

$$\mathbb{P}\left[f^+(p_i, x_i) \leq \frac{1-\alpha}{2} \text{ or } f^-(p_i, x_i) \leq \frac{1-\alpha}{2}\right] \leq 1-\alpha \quad (11)$$

or in other words

$$\mathbb{P}\left[f^+(p_i, x_i) > \frac{1-\alpha}{2} \text{ and } f^-(p_i, x_i) > \frac{1-\alpha}{2}\right] \geq \alpha. \quad (12)$$

Combinatorially, one can see that $f^+(p_i, x_i)$ must be a strictly decreasing function of p_i and $f^-(p_i, x_i)$ must be a strictly increasing function of p_i (holding x_i fixed and letting p_i range from 0 to 1). So if we take $p_{\min}$ to be the (unique) solution to

$$f^-(p_{\min}, x_i) = \frac{1-\alpha}{2} \quad (13)$$

and $p_{\max}$ to be the (unique) solution to

$$f^+(p_{\max}, x_i) = \frac{1-\alpha}{2} \quad (14)$$

then by (12), we have:

$$\mathbb{P}[p_{\min} < p_i < p_{\max}] \geq \alpha. \quad (15)$$

The confidence interval provided by the application are these values of $p_{\min}$ and $p_{\max}$. Note that p_i is a fixed and unknown quantity and *not* a random variable (but the $p_{\min}$ and $p_{\max}$ are, since they are functions of the random variable x_i).

Simultaneous confidence intervals

Take $p_{\min,k}$ to be the unique solution to:

$$f^-(p_{\min,k}, x_i) = \frac{1-\alpha}{2k} \quad (16)$$

and $p_{\max,k}$ to be the unique solution to

$$f^+(p_{\max,k}, x_i) = \frac{1 - \alpha}{2k}. \quad (17)$$

For each card i , the argument of (15) shows:

$$\mathbb{P}[p_{\min,k} < p_i < p_{\max,k}] \geq 1 - \frac{1 - \alpha}{k}; \quad (18)$$

or in other words:

$$\mathbb{P}[p_{\min,k} \geq p_i \text{ or } p_{\max,k} \leq p_i] \leq \frac{1 - \alpha}{k}. \quad (19)$$

Taking the union bound, we have:

$$\mathbb{P}[p_{\min,k} \geq p_i \text{ or } p_{\max,k} \leq p_i \text{ for some } i] \leq 1 - \alpha; \quad (20)$$

in other words

$$\mathbb{P}[p_{\min,k} < p_i < p_{\max,k} \text{ for all } i] \geq \alpha. \quad (21)$$

The interval provided by the application are these values of $p_{\min,k}$ and $p_{\max,k}$.

While it is true that these simultaneous confidence intervals will contain all the true values of p_i with probability at least α , our confidence intervals may be slightly narrower than are necessary. But despite the naïveté of the union bound, “cleverer” approaches surprisingly lead to even wider such intervals (see for instance, this paper by Leo Goodman [Goo65]).

Option 2

Simultaneous confidence intervals

The simultaneous confidence intervals for Option 2 are computed in the same manner as those for Option 1, except that k is now smaller.

Option 3

Review of Dirichlet distributions

Up until this point, we treated the probabilities $\{p_i\}_{i \in [k]}$ as fixed unknowns; in Option 3, they are treated as random variables. For notational convenience, we will let:

$$\Delta := \{(p_1, p_2, \dots, p_k) \mid \sum_{i=1}^k p_i = 1 \text{ and all } 0 \leq p_i \leq 1\} \subset \mathbb{R}^k, \quad (22)$$

which is the space of possible values for the $\{p_i\}$. Δ is a simplex with volume

$$\int_{\vec{x} \in \Delta} 1 \prod_{i=1}^k dp_i = \frac{1}{k!} \quad (23)$$

(several proofs may be found here). Before we discuss the user-inputted prior probability that there is no shortprinting, suppose we were completely agnostic about what the true values of the $\{p_i\}_{i \in [k]}$ were. The prior distribution on Δ would then be:

$$f_{\text{prior}}(p_1, p_2, \dots, p_k) \prod_{i=1}^k dp_i = k! \prod_{i=1}^k dp_i \quad (24)$$

but for simplicity, it is better to view

$$f_{\text{prior}}(p_1, p_2, \dots, p_k) \prod_{i=1}^k dp_i \sim 1 \prod_{i=1}^k dp_i \quad (25)$$

(i.e. $f_{\text{prior}}(p_1, p_2, \dots, p_k)$ is some constant multiple of 1), and not worry about the exact constant of proportionality. How should we update after making our observations of the $\{x_i\}_{i \in [k]}$? Conditional on given values of the $\{p_i\}$, the probability of our observations are:

$$\prod_{i=1}^k (p_i)^{x_i} \quad (26)$$

so via Bayes' theorem, our posterior distribution is:

$$f_{\text{post}}(p_1, p_2, \dots, p_k) \prod_{i=1}^k dp_i \sim \prod_{i=1}^k (p_i)^{x_i} \prod_{i=1}^k dp_i. \quad (27)$$

What is the constant of proportionality? From Wikipedia:

$$\int_{\Delta} \prod_{i=1}^k (p_i)^{x_i} \prod_{i=1}^k dp_i = B(x_1 + 1, x_2 + 1, \dots, x_k + 1) \quad (28)$$

(where B is the multivariate Beta function, described here in terms of the Gamma function). Since the $\{x_i\}$ are all non-negative integers, we have:

$$B(x_1 + 1, x_2 + 1, \dots, x_k + 1) = \frac{(x_1)!(x_2)! \cdots x_k!}{n!} = 1 / \binom{n}{x_1, x_2, \dots, x_k}, \quad (29)$$

using the fact that $\Gamma(x) = (x-1)!$ for positive integers x . We may then express our posterior as:

$$f_{\text{post}}(p_1, p_2, \dots, p_k) \prod_{i=1}^k = \binom{n}{x_1, x_2, \dots, x_k} \prod_{i=1}^k (p_i)^{x_i} \prod_{i=1}^k dp_i. \quad (30)$$

These posteriors are known as *Dirichlet distributions*. When new observations come in, one simply updates the $\{x_i\}$ and recomputes the above distribution.

Averaging a discrete and continuous prior

We will take q to be the user-inputted prior probability that there is no shortprinting. So with probability q , we have

$$p_i = \frac{1}{k} \text{ for all } i \in [k]. \quad (31)$$

To simplify the application, we have assumed that the prior on the rest of Δ is uniform, but we will discuss in the next subsection how one can modify this to a more general prior. Our prior is now:

$$q\delta_{\frac{1}{k}} + (1-q) \prod_{i=1}^k dp_i \quad (32)$$

where $\delta_{\frac{1}{k}}$ is the Dirac measure with support at the point $(\frac{1}{k}, \frac{1}{k}, \dots, \frac{1}{k}) \in \mathbb{R}^k$. For the reader unfamiliar with measure theory, $\delta_{\frac{1}{k}}$ may be viewed as a probability distribution that is very, very narrowly concentrated at the point $(\frac{1}{k}, \frac{1}{k}, \dots, \frac{1}{k})$, but still has integral 1 (informally speaking, this distribution is “infinitely narrow” and “infinitely concentrated” at that point). We did not write f_{prior} because the Dirac measure *cannot* be expressed as the product of a measurable function and the uniform distribution $\prod_{i=1}^k dp_i$.

Updating is not too different than the previous case. The probability of our outcomes conditional on all $p_i = \frac{1}{k}$ is

$$\prod_{i=1}^k \frac{1}{k^{x_i}} = \frac{1}{k^n} \quad (33)$$

so directly applying Bayes’ theorem, the posterior must be proportional to

$$\frac{q}{k^n} \delta_{\frac{1}{k}} + (1-q) \prod_{i=1}^k (p_i)^{x_i} dp_i. \quad (34)$$

Integrating the above over Δ gives $\frac{q}{k^n} + (1 - q)/\binom{n}{x_1, x_2, \dots, x_k}$ (using (28)) so our posterior distribution is

$$\frac{\frac{q}{k^n} \delta_{\frac{1}{k}} + (1 - q) \prod_{i=1}^k (p_i)^{x_i} dp_i}{\frac{q}{k^n} + (1 - q)/\binom{n}{x_1, x_2, \dots, x_k}}. \quad (35)$$

More general priors (not included in application)

It might feel unnatural to use have the prior be uniform outside of the point $(\frac{1}{k}, \frac{1}{k}, \dots, \frac{1}{k})$. If you knew shortprints would occur, you would probably bet on specific cards being printed less than others. As long as one restricts oneself to priors of the form:

$$q \delta_{\frac{1}{k}} + (1 - q) \prod_{i=1}^k \frac{\prod_{i=1}^k (p_i)^{\alpha_i} dp_i}{B(\alpha_1 + 1, \alpha_2 + 1, \dots, \alpha_k + 1)}, \quad (36)$$

(where the $\{\alpha_i\}_{i \in [k]}$ are non-negative real numbers) this is still quite easy to work with. To pick the $\{\alpha_i\}_{i \in [k]}$, consider the following thought experiment: Suppose you knew for a fact that there would be shortprints in this set but also forgot everything you saw about the data. What would you expect to see in the first 100 cards pulled from that rarity? For instance, you could look at all past instances of shortprints and use the ratios that seem to appear the most often. For each $i \in [k]$, take α_i to be your guess for the number of card i that would appear in the first 100 pulls of that rarity. (The number 100 can of course be replaced with any convenient quantity.) If choosing the α_i in this manner, the Beta function may be rewritten via (29) since the arguments would all be positive integers. Analogously to (34), the posterior distribution would be proportional to

$$\frac{q}{k^n} \delta_{\frac{1}{k}} + \frac{(1 - q) \prod_{i=1}^k (p_i)^{x_i + \alpha_i} dp_i}{B(\alpha_1 + 1, \alpha_2 + 1, \dots, \alpha_k + 1)} \quad (37)$$

and the integral over this would be

$$\frac{q}{k^n} + \frac{(1 - q) B(\alpha_1 + x_1 + 1, \alpha_2 + x_2 + 1, \dots, \alpha_k + x_k + 1)}{B(\alpha_1 + 1, \alpha_2 + 1, \dots, \alpha_k + 1)}. \quad (38)$$

The posterior distribution would be (37) divided by (38). If the $\{\alpha_i\}$ were all taken to be integers that summed to 100, we could write:

$$\frac{B(\alpha_1 + x_1 + 1, \dots, \alpha_k + x_k + 1)}{B(\alpha_1 + 1, \dots, \alpha_k + 1)} = \frac{\binom{n}{\alpha_1, \dots, \alpha_k}}{\binom{n+100}{\alpha_1+x_1, \dots, \alpha_k+x_k}} \quad (39)$$

$$= \frac{n!}{(100+n)!} \prod_{i=1}^k \frac{(\alpha_i + x_i)!}{(x_i)!}. \quad (40)$$

We did not include this flexibility in the application as the explanation for its use would be too technical for a general audience. But the user may copy and suitably modify the code of the application to serve this purpose if they wish.

Credible intervals

Finally we come to what the application actually computes in Option 3: the credible intervals for the desired confidence, which we will call α (and we encourage the reader to set $\alpha = .95$ when reading the subsequent discussion). To do so, we begin by computing the marginal distribution of each p_i in the joint posterior distribution in (35). We will slightly rewrite this joint posterior as:

$$\frac{\frac{q}{k^n} \delta_{\frac{1}{k}} + \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_1, x_2, \dots, x_k} \prod_{i=1}^k (p_i)^{x_i} dp_i}{\frac{q}{k^n} + (1-q) / \binom{n}{x_1, x_2, \dots, x_k}}, \quad (41)$$

to view it as a weighted average of the Dirac measure $\delta_{\frac{1}{k}}$ and the Dirichlet distribution $\binom{n}{x_1, x_2, \dots, x_k} \prod_{i=1}^k (p_i)^{x_i} dp_i$. The marginal distribution of $\delta_{\frac{1}{k}}$ on p_i is itself a Dirac measure with support $\frac{1}{k}$ (which we'll denote by $\delta_{\frac{1}{k}, p_i}$). And the marginal distribution of the Dirichlet distribution $\binom{n}{x_1, x_2, \dots, x_k} \prod_{i=1}^k (p_i)^{x_i} dp_i$ is itself the Dirichlet distribution:

$$\binom{n}{x_i} (p_i)^{x_i} (1 - p_i)^{n-x_i} dp_i \quad (42)$$

(see again Wikipedia). Taking the appropriate weighted averages, the marginal posterior on each p_i is:

$$\frac{\frac{q}{k^n} \delta_{\frac{1}{k}, p_i} + \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i}{\frac{q}{k^n} + (1-q)/\binom{n}{x_1, x_2, \dots, x_k}}. \quad (43)$$

Let now α be the user-inputted desired credible level (for instance, $\alpha = .95$). We want to find a “good” choice of ℓ_i and r_i such that $0 \leq \ell_i \leq r_i \leq 1$ and:

$$\int_{p_i=\ell_i}^{r_i} \left(\frac{\frac{q}{k^n} \delta_{\frac{1}{k}, p_i} + \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i}{\frac{q}{k^n} + (1-q)/\binom{n}{x_1, x_2, \dots, x_k}} \right) \geq \alpha; \quad (44)$$

in other words, $\mathbb{P}[\ell_i \leq p_i \leq r_i] \geq \alpha$. A naïve approach might be to solve:

$$\mathbb{P}[p_i \leq \ell_i] = \frac{1-\alpha}{2}, \quad \mathbb{P}[p_i \geq r_i] = \frac{1-\alpha}{2} \quad (45)$$

for ℓ_i and r_i but the danger is that the Dirac measure might prevent solutions from occurring! Instead, we find the *maximum* ℓ_i and *minimum* r_i such that:

$$\mathbb{P}[p_i < \ell_i] \leq \frac{1-\alpha}{2}, \quad \mathbb{P}[p_i > r_i] \geq 1 - \frac{1-\alpha}{2}. \quad (46)$$

For notational simplicity, we will set

$$A := \left(\frac{q}{k^n} + (1-q)/\binom{n}{x_1, x_2, \dots, x_k} \right) \frac{1-\alpha}{2} \quad (47)$$

so we may write these requirements as:

$$\int_{p_i=0}^{p_i=\ell_i} \frac{q}{k^n} \delta_{\frac{1}{k}, p_i} + \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i \geq A \quad (48)$$

$$\int_{p_i=r_i}^{p_i=1} \frac{q}{k^n} \delta_{\frac{1}{k}, p_i} + \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i \geq A. \quad (49)$$

We begin with ℓ_i , and calculate:

$$L := \int_{p_i=0}^{p_i=1/k} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i. \quad (50)$$

If $L \geq A$, then our desired value of ℓ_i is the solution to:

$$\int_{p_i=0}^{p_i=\ell_i} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i = A \quad (51)$$

(which will be at most $\frac{1}{k}$). Rewriting this as:

$$\int_{p_i=0}^{p_i=\ell_i} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i = \frac{A \binom{n}{x_1, x_2, \dots, x_k}}{(1-q)}, \quad (52)$$

this value of ℓ_i may be obtained using the `scipy.stats.beta.ppf` function in Python's Scipy package. If on the other hand, $L < A$, we next check if

$$L + \frac{q}{k^n} \geq A. \quad (53)$$

If it is, then our desired value of ℓ_i is $\frac{1}{k}$. If we have $L + \frac{q}{k^n} < A$, then ℓ_i is the solution to:

$$\frac{q}{k^n} + \int_{p_i=0}^{p_i=\ell_i} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i = A \quad (54)$$

(which will be strictly greater than $\frac{1}{k}$.) This again may be solved via the `scipy.stats.beta.ppf` function by rewriting the expression in a similar manner to (52). Solving for r_i is very similar; we begin by calculating:

$$R := \int_{p_i=1/k}^{p_i=1} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i \quad (55)$$

If $R \geq A$, then the value of r_i is the solution to:

$$\int_{p_i=r_i}^{p_i=1} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i = A, \quad (56)$$

which will be at least $\frac{1}{k}$. If $R < A$, we check if

$$R + \frac{q}{k^n} \geq A; \quad (57)$$

if it is, then $r_i = \frac{1}{k}$. If $R + \frac{q}{k^n} < A$, then r_i is the solution to:

$$\frac{q}{k^n} + \int_{p_i=r_i}^{p_i=1} \frac{(1-q)}{\binom{n}{x_1, x_2, \dots, x_k}} \binom{n}{x_i} (p_i)^{x_i} (1-p_i)^{n-x_i} dp_i = A \quad (58)$$

which will be at most $\frac{1}{k}$. These are the values of ℓ_i and r_i provided by the program. The code may look strange for these credible intervals, as our algorithm is designed to minimize the risk of floating point errors.

External Links

Shortprint application: <https://shortprint.streamlit.app/>

Github code for the application: <https://github.com/flipflipshift/shortprint-calculator/blob/main/application.py>

Wikipedia article on Pearson's chi-square test: https://en.wikipedia.org/wiki/Pearson%27s_chi-squared_test

Wikipedia article on the Clopper–Pearson interval: https://en.wikipedia.org/wiki/Binomial_proportion_confidence_interval#Clopper%E2%80%9393Pearson_interval

Stackexchange discussion on calculating the volume of Δ : math.stackexchange.com/questions/769545/volume-of-t-n-x-i-ge0x-1-cdotsx-n-le1/

Wikipedia article on Dirichlet distributions: https://en.wikipedia.org/wiki/Dirichlet_distribution

Wikipedia article on the multivariate Beta distribution: https://en.wikipedia.org/wiki/Beta_function#Multivariate_beta_function.

Wikipedia article on the Dirac measure: https://en.wikipedia.org/wiki/Dirac_measure

Reference

[Goo65] Leo A. Goodman, *On simultaneous confidence intervals for multinomial proportions*, *Technometrics* **7** (1965), no. 2, 247–254.