

assignment EM2

Problem 1

Solution. Consider the boundary condition for graphene

$$\begin{aligned}n \times (E_1 - E_2) &= 0 \\n \times (H_1 - H_2) &= \sigma_s E_t \\E_{1t} &= E_{2t} = E_t \\H_{1t} - H_{2t} &= \sigma_s E_t\end{aligned}$$

For TE wave

$$\begin{aligned}E_y &= e_y(z)e^{-jk_x x} \\e_{1y}(z) &= e_0 \cdot e^{jk_{1z}z} + R \cdot e_0 \cdot e^{-jk_{1z}z} \\e_{2y}(z) &= T \cdot e_0 \cdot e^{jk_{2z}z} \\h_{1y}(z) &= -\frac{k_{1z}}{\omega\mu_1}(e_0 \cdot e^{jk_{1z}z} - R \cdot e_0 \cdot e^{-jk_{1z}z}) \\h_{2y}(z) &= -\frac{k_{2z}}{\omega\mu_2}(T \cdot e_0 \cdot e^{jk_{2z}z})\end{aligned}$$

At $z=0$

$$\begin{aligned}1 + R &= T \\-\frac{k_{1z}}{\omega\mu_1}(1 - R) &= -\frac{k_{2z}}{\omega\mu_2} \cdot T + \sigma_s \cdot T\end{aligned}$$

Solve the equation

$$\begin{aligned}R &= \frac{-k_{1z}\mu_2 + k_{2z}\mu_1 - \sigma_s \cdot \omega\mu_1\mu_2}{\sigma_s \cdot \omega\mu_1\mu_2 - k_{1z}\mu_2} \\T &= \frac{-2k_{1z}\mu_2 + k_{2z}\mu_1}{\sigma_s \cdot \omega\mu_1\mu_2 - k_{1z}\mu_2}\end{aligned}$$

For TM wave

$$\begin{aligned}H_y &= h_y(z)e^{-jk_x x} \\h_{1y}(z) &= h_0 \cdot e^{jk_{1z}z} + R \cdot h_0 \cdot e^{-jk_{1z}z} \\h_{2y}(z) &= T \cdot h_0 \cdot e^{jk_{2z}z} \\e_{1y}(z) &= -\frac{k_{1z}}{\omega\epsilon_1}(h_0 \cdot e^{jk_{1z}z} - R \cdot h_0 \cdot e^{-jk_{1z}z}) \\e_{2y}(z) &= -\frac{k_{2z}}{\omega\epsilon_2}(T \cdot h_0 \cdot e^{jk_{2z}z})\end{aligned}$$

assignment EM2

At $z=0$

$$\begin{aligned}1 + R &= T + \sigma_s \cdot \left(-\frac{k_{2z}}{\omega\epsilon_2} \cdot T\right) \\ -\frac{k_{1z}}{\omega\epsilon_1}(1 - R) &= -\frac{k_{2z}}{\omega\epsilon_2} \cdot T\end{aligned}$$

Solve the equation

■

Problem 2

Solution. $Die1 = 2.5^2 \cdot \text{ones}(\text{length}(y), 1)$; $Die2 = 1.25^2 \cdot \text{ones}(\text{length}(y), 1)$; $Die3 = 1.875^2 \cdot \text{ones}(\text{length}(y), 1)$;

It is shown that total reflection can not cover all incidence angles for TM incidence. And a pass band will be opened in the band gap when the defect is introduced. Possible reason is that the break of perfect periodic boundary condition induces destructive interference between incident and reflected wave within some certain frequency range.

■

assignment EM2




Figure 1: TE incidence




Figure 2: TM incidence




Figure 3: TE incidence with defect

assignment EM2




Figure 4: TM incidence with defect