# Enterovirus Surveillance in Thailand

Fanglin Kuo

2018/06

1

Country	Characteristics		
Vietnam	<ol> <li>Seroprevalence of EV-A71 infections: 55%</li> <li>1,500,000 newborns/year</li> <li>Ongoing vaccine development</li> <li>Member of ASEAN</li> </ol>		
Malaysia	<ol> <li>Seroprevalence of EV-A71 infections: 60%</li> <li>500,000 newborns/year</li> <li>Ongoing vaccine development</li> <li>Member of ASEAN and PIC/S</li> </ol>		
Cambodia	<ol> <li>Seroprevalence of EV-A71 infections: 88.8%</li> <li>400,000 newborns/year</li> <li>Member of ASEAN</li> </ol>		
Thailand	<ol> <li>Seroprevalence of infections: CV-A (31.2%), EV-A71 (9.1%)</li> <li>765,600 newborns/year</li> <li>Member of ASEAN and PIC/S</li> </ol>		

ASEAN: Association of Southeast Asian Nations

PIC/S: Pharmaceutical Inspection Convention and Pharmaceutical Inspection Co-operation Scheme


2018/06

#### Websites


- Hand, foot and mouth disease (HFMD)
  - Bureau of Epidemiology, Department of Disease Control
  - http://www.boe.moph.go.th/boedb/surdata/disease.php?ds=71 (Th)
- Data: 2003 2018 presence
- ICD-10: B08.4 Enteroviral vesicular stomatitis with exanthem

2018/06

3







5


2018/06

### 2018 HFMD

- 1 January 2018 15 May 2018, a total of 11672 cases were reported from 77 provinces. There were 0 deaths.
- The highest proportion of cases by age group: 1 years old (27.78%); 2 years old (22.53%); 3 years old (16.85%)

http://www.boe.moph.go.th/boedb/surdata/506wk/y61/en/d71 1361 en.pdf

2018/06



7

#### Case definitions

- Herpangina (HA): Having well-characterized vesicular enanthem, oral ulcers on the anterior tonsil pillars, soft palate, buccal mucosa, or uvula.
- HFMD: Oral ulcers mainly on the buccal mucosa and tongue, accompanied by typical vesicular rashes most commonly on the extensor surfaces of the hands, feet, knees, and/or buttock.
- A severe case was defined as having HFMD or HA and accompanied by the occurrence of at least one of the following complications: aseptic meningitis, myocarditis, encephalitis, pulmonary edema, hemorrhage, acute flaccid paralysis, and cardiopulmonary collapse

2018/06

## Virus characterization of HFMD (2008 – 2013)

Table 1 Demographic and clinical data associated with different enterovirus genotypes

	HFMD	НА	HFMD/HA	All
No of case	795 (67.3%)	197 (16.7%)	190 (16.1%)	1182
No of sample	833 (67.3%)	202 (16.4%)	202 (16.4%)	1237
Age range (median)	4d to 54y (2.4y)	3 m to 16y (2.6y)	1 m to 50y (3y)	4d to 54y (2.5y)
Mean age ± SD	$3.4y \pm 4.8y$	$3.4y \pm 3.0y$	5.1y ± 8.3y	$3.6y \pm 5.0y$
Sex ratio (F:M)	1:1.4	1:1.1	1:1.2	1:1.3
Virology: EV71	66 (8.3%)	2 (1%)	40 (21.1%)	108 (9.1%)
CV-A16	65 (8.2%)	2 (1%)	35 (18.4%)	102 (8.6%)
Other CV-A	217 (27.3%)	22 (11.2%)	28 (14.7%)	267 (22.6%)
Pan-EV	131 (16.5%)	88 (44.7%)	10 (5.3%)	229 (19.4%)
Negative	316 (39.7%)	83 (42.1%)	77 (40.5%)	476 (40.3%)

HFMD, Hand, foot, and mouth disease; HA, Herpangina; EV71, Human enterovirus 71; CV-A, Coxsackievirus species A; Pan-EV, Pan-enterovirus; y, year old; m, month old.

(Linsuwanon, Puenpa et al. 2014)

2018/06

Year	Sample	Sample size/disease	Age	%	Reference
2014 – 2016	Specimens	1049 Patients with HFMD		CV-A6: 32.9% CV-A16: 11.6%	(Puenpa, Suwannakarn et al. 2017)
2010 – 2016	Fecal samples	1310 HFMD patients with no diarrhea	1 – 66y	EV-A: 92.6% CV-A6: 54.8%	(Chansaenroj, Tuanthap et al. 2017)
2010 – 2014	Fecal samples	1266 at Chiang Mai Patients with acute gastroenteritis	< 12y	EV: 5.8% EV-C96: 12.3%	(Kumthip, Khamrin et al. 2017)
2011 – 2013	Specimens	11 at Lopburi Patients with HFMD	<15y	CV-A16: 44.5% EV-A71: 29.6%	(Owatanapanich, Wutthanarungsan et al. 2016)
2013	Specimens	203	16d - 50y	EV-A71: 30% CV-A8: 26% CV-A16: 21%	(Mauleekoonphairoj, Puenpa et al. 2015)
2008 – 2013	Specimens	1182 Patients with HFMD or HA	4d - 54y	EV-A71: 9.1% CV-A: 31.2%	(Linsuwanon, Puenpa et al. 2014)
2012	Specimens	704 Patients with HFMD or HA	1m - 54y	CV-A6: 33.5% CV-A16: 9.4% EV-A71: 8.8%	(Puenpa, Mauleekoonphairoj et al. 2014)
2010 – 2011	Specimens	6266 Children	6m - 10y	EV: 41.5%	(Taylor, Lopez et al. 2017)
2008 – 2011	Resp samples	121 at Bangkok		EV: 43% EV-A71	(Zhou, Fernandez et al. 2016)
2006 – 2011	Resp samples	1810 Patients with resp illnesses	< 15y	EV-D68: 1.4%	(Linsuwanon, Puenpa et al. 2012)
2008 – 2009	Specimens	48	4m - 26y	EV-A71: 47.9% CA-16: 8.3%	(Chatproedprai, Theanboonlers et al. 2010)

#### Summary

- HFMD occurred endemically during 2008-2017. CV-A6, CV-A16, and EV-A71 were the three most commonly identified EV associated with HFMD.
- EV-71 had a high prevalence in 2008-2009 and has been identified with a higher frequency since 2012. Co-circulation between different EV-71 genotypes was observed during the outbreaks.
- Surveillance of epidemiology and monitoring of disease severit should be continued

2018/06

11

#### References

- Chansaenroj, J., S. Tuanthap, T. Thanusuwannasak, A. Duang-in, S. Klinfueng, N. Thaneskongtong, V. Vutithanachot, S. Vongpunsawad and Y. Poovorawan (2017). "Human enteroviruses associated with and without diarrhea in Thailand between 2010 and 2016." PloS one 12(7): e0182078.
- Chatproedprai, S., A. Theanboonlers, S. Korkong, C. Thongmee, S. Wananukul and Y. Poovorawan (2010). "Clinical and molecular characterization of hand-foot-andmouth disease in Thailand, 2008-2009." Jpn J Infect Dis 63(4): 229-233.
- Kumthip, K., P. Khamrin, H. Ushijima and N. Maneekarn (2017). "Multiple enterovirus genotypes circulating in children hospitalized with acute gastroenteritis in Thailand." Infection, Genetics and Evolution 55: 324-331.
- Linsuwanon, P., J. Puenpa, S.-W. Huang, Y.-F. Wang, J. Mauleekoonphairoj, J.-R. Wang and Y. Poovorawan (2014). "Epidemiology and seroepidemiology of human enterovirus 71 among Thai populations." Journal of biomedical science 21(1): 16.
- Linsuwanon, P., J. Puenpa, K. Suwannakarn, V. Auksornkitti, P. Vichiwattana, S. Korkong, A. Theamboonlers and Y. Poovorawan (2012). "Molecular epidemiology and evolution of human enterovirus serotype 68 in Thailand, 2006–2011." PloS one 7(5): e35190.
- Mauleekoonphairoj, J., J. Puenpa, S. Korkong, S. Vongpunsawad and Y. Poovorawan (2015). "Prevalence of human enterovirus among patients with hand, foot, and mouth disease and herpangina in Thailand, 2013." Southeast Asian Journal of Tropical Medicine and Public Health 46(6): 1013.
- Owatanapanich, S., R. Wutthanarungsan, W. Jaksupa and U. Thisyakorn (2016). "Risk Factors for Severe Enteroviral Infections in Children." <u>Journal of the Medical Association of Thailand = Chotmainet thangphaet</u> 99(3): 322-330.
- Puenpa, J., J. Mauleekoonphairoj, P. Linsuwanon, K. Suwannakarn, T. Chieochansin, S. Korkong, A. Theamboonlers and Y. Poovorawan (2014). "Prevalence and characterization of enterovirus infections among pediatric patients with hand foot mouth disease, herpangina and influenza like illness in Thailand, 2012." PloS one 9(6): e98888.
- Puenpa, J., K. Suwannakarn, J. Chansaenroj, S. Vongpunsawad and Y. Poovorawan (2017). "Development of single-step multiplex real-time RT-PCR assays for rapid diagnosis of enterovirus 71, coxsackievirus A6, and A16 in patients with hand, foot, and mouth disease." <u>Journal of virological methods</u> 248: 92-99.
- Taylor, S., P. Lopez, L. Weckx, C. Borja-Tabora, R. Ulloa-Gutierrez, E. Lazcano-Ponce, A. Kerdpanich, M. A. R. Weber, A. M. de Los Santos and J.-C. Tinoco (2017). "Respiratory viruses and influenza-like illness: Epidemiology and outcomes in children aged 6 months to 10 years in a multi-country population sample." <u>Journal of Infection</u> 74(1): 29-41.
- Zhou, Y., S. Fernandez, I.-K. Yoon, S. Simasathien, V. Watanaveeradei, Y. Yang, O. A. Marte-Salcedo, D. J. Shuck-Lee, S. J. Thomas and J. Hang (2016). "Metagenomics study of viral pathogens in undiagnosed respiratory specimens and identification of human enteroviruses at a Thailand hospital." <u>The American journal of tropical medicine and hygiene</u> 95(3): 663-669.

2018/06