

Social evolution in structured populations

Florence Débarre



CNRS


Centre Interdisciplinaire de Recherche en Biologie, Paris


Phénomènes de propagation et d'organisation spatiale en biologie



Journée thématique interdisciplinaire maths-bio


Université Paris-Dauphine

4 décembre 2017











Assortment / Relatedness


Spatial structure, population viscosity and altruism




Spatial structure, population viscosity and altruism




Spatial structure, population viscosity and altruism



Spatial structure, population viscosity and altruism



Spatial structure, population viscosity and altruism



Evolutionary Ecology, 1992, 6, 352–356

Altruism in viscous populations – an inclusive fitness model

P.D. TAYLOR

Department of Mathematics and Statistics, Queen's University, Kingston Ont. K7L 3N6, Canada


Summary

A viscous population (Hamilton, 1964) is one in which the movement of organisms from their place of birth is relatively slow. This viscosity has two important effects: one is that local interactions tend to be among relatives, and the other is that competition for resources tends to be among relatives. The first effect tends to promote and the second to oppose the evolution of altruistic behaviour. In a simulation model of Wilson *et al.* (1992) these two factors appear to exactly balance one another, thus opposing the evolution of local altruistic behaviour. Here I show, with an inclusive fitness model, that the same result holds in a patch-structured population.

Keywords: altruism; inclusive fitness; competition; viscosity


Subdivided population – Island model

N_d demes




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)




Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)



Subdivided population – Island model

N_d demes of n individuals each (total population size $N = n N_d$)



The choice of life-cycle matters

Constant population size (N), so between two time steps, $\#\text{▀} = \#\text{👶}$.

The choice of life-cycle matters

Constant population size (N), so between two time steps, $\#\text{▀} = \#\text{▀}$.

Wright-Fisher



Moran Birth-Death



Moran Death-Birth



The choice of life-cycle matters

Constant population size (N), so between two time steps, $\#\text{▀} = \#\text{🚼}$.


Wright-Fisher



Moran Birth-Death





Moran Death-Birth





In homogeneously structured populations,
with effects of social interactions on **fecundity**.

A common feature of models



A common feature of models



A common feature of models



A common feature of models




What is the effect of population viscosity
on the evolution of altruism when parent-
offspring strategy transmission is **imperfect**?

Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission


- ▶ Mutation



Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission


- ▶ Mutation
- ▶ Partial heritability



Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission


- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)



Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission

- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)




In the model

Parent Offspring




Fidelity of parent-offspring transmission




Causes of imperfect strategy transmission

- ▶ Mutation
 - ▶ Partial heritability
 - ▶ Cultural transmission (vertical)

In the model




Fidelity of parent-offspring transmission



Causes of imperfect strategy transmission

- ▶ Mutation
 - ▶ Partial heritability
 - ▶ Cultural transmission (vertical)


In the model




Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission

- ▶ Mutation
 - ▶ Partial heritability
 - ▶ Cultural transmission (vertical)



In the model




$$\nu = \frac{\mu_B}{\mu_A + \mu_B}$$


Fidelity of parent-offspring transmission

Causes of imperfect strategy transmission

- ▶ Mutation
 - ▶ Partial heritability
 - ▶ Cultural transmission (vertical)



In the model




Notation

$$\mathbf{x}(t); \quad x_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{duck} \text{ at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{bird} \text{ at time } t (1 \leq i \leq N) \end{cases}$$

Notation

$$\mathbf{X}(t); \quad X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{duck} \text{ at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t (1 \leq i \leq N) \end{cases}$$




$$\mathbb{E}[Y_i] = (1 - \mu)X_i + \mu\nu \times 1 + \mu(1 - \nu) \times 0.$$

Expected trait of the
offspring of individual i

Notation

$$\mathbf{X}(t); \quad X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{duck} \text{ at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t (1 \leq i \leq N) \end{cases}$$



$$\mathbb{E}[Y_i] = (1 - \mu)X_i + \mu\nu$$

Expected trait of the offspring of individual i

Notation

$$\mathbf{x}(t); \quad x_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{duck} \text{ at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t (1 \leq i \leq N) \end{cases}$$

Proportion of altruists in the population:

$$\bar{x} = \frac{1}{N} \sum_{i=1}^N x_i.$$

We want to compute $\mathbb{E}[\bar{x}]$,
the expected proportion of altruists in the population.

Phenotype

$$\phi_i = \delta X_i,$$

and we assume that $\delta \ll 1$. (Selection is weak.)

Social interactions

Phenotype

$$\phi_i = \delta X_i,$$

and we assume that $\delta \ll 1$. (Selection is weak.)

Social interactions affect fecundity

At the first order in δ ,

$$f_i = 1 + \delta \left(b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

Social interactions

Phenotype

$$\phi_i = \delta X_i,$$

and we assume that $\delta \ll 1$. (Selection is weak.)

Social interactions affect fecundity

At the first order in δ ,

$$f_i = 1 + \delta \left(b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$


Social interactions

Phenotype

$$\phi_i = \delta X_i,$$

and we assume that $\delta \ll 1$. (Selection is weak.)

Social interactions affect fecundity

At the first order in δ ,

$$f_i = 1 + \delta \left(b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

Proportion of altruists
among the other deme-mates

Phenotype

$$\phi_i = \delta X_i,$$

and we assume that $\delta \ll 1$. (Selection is weak.)

Social interactions affect fecundity

At the first order in δ ,

$$f_i = 1 + \delta \left(b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

Proportion of altruists among the other deme-mates

The cost is only paid by altruists

Calculations

Notation

$B_i = B_i(\mathbf{X}, \delta)$: expected # of offspring of individual i ;

$D_i = D_i(\mathbf{X}, \delta)$: probability that i dies.

Calculations

Notation

$B_i = B_i(\mathbf{X}, \delta)$: expected # of offspring of individual i ;

$D_i = D_i(\mathbf{X}, \delta)$: probability that i dies.

- ▶ Expected proportion of altruists at $t + 1$ in the proportion of altruists, conditional on the state of the population at time t :

$$\mathbb{E}[\bar{X}(t+1) | \mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N [B_i(1 - \mu)X_i + (1 - D_i)X_i + B_i\mu\nu]$$

Calculations

Notation

$B_i = B_i(\mathbf{X}, \delta)$: expected # of offspring of individual i ;

$D_i = D_i(\mathbf{X}, \delta)$: probability that i dies.

- ▶ Expected proportion of altruists at $t + 1$ in the proportion of altruists, conditional on the state of the population at time t :

$$\mathbb{E}[\bar{X}(t+1) | \mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N [B_i(1 - \mu)X_i + (1 - D_i)X_i + B_i\mu\nu]$$

- ▶ Take expectation and let $t \rightarrow \infty$; consider stationary distribution ξ

$$0 = \frac{1}{N} \sum_{X \in \Omega} \left[\sum_{i=1}^N \underbrace{B_i(1 - \mu) - D_i}_{W_i} X_i + \sum_{i=1}^N B_i\mu\nu \right] \xi(\mathbf{X}, \delta, \mu)$$

Calculations

Notation

$B_i = B_i(\mathbf{X}, \delta)$: expected # of offspring of individual i ;

$D_i = D_i(\mathbf{X}, \delta)$: probability that i dies.

- ▶ Expected proportion of altruists at $t + 1$ in the proportion of altruists, conditional on the state of the population at time t :

$$\mathbb{E}[\bar{X}(t+1) | \mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N [B_i(1 - \mu)X_i + (1 - D_i)X_i + B_i\mu\nu]$$

- ▶ Take expectation and let $t \rightarrow \infty$; consider stationary distribution ξ

$$0 = \frac{1}{N} \sum_{X \in \Omega} \left[\sum_{i=1}^N \underbrace{B_i(1 - \mu) - D_i}_{W_i} X_i + \sum_{i=1}^N B_i\mu\nu \right] \xi(\mathbf{X}, \delta, \mu)$$

Calculations (2)

- Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

Calculations (2)

- Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} \chi_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* \chi_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[\frac{\partial W_i}{\partial \delta} \chi_i \right] + O(\delta^2).$$

Calculations (2)

- Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[\frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

Calculations (2)

- Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[\frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

Calculations (2)

- ▶ Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[\frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

- ▶ Using partial derivatives: phenotypes

$$\frac{\partial W_i}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \frac{\partial \phi_k}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} X_k.$$

Calculations (2)

- Selection is weak ($\delta \ll 1$) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[\sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[\frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

- Using partial derivatives: phenotypes

$$\frac{\partial W_i}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \frac{\partial \phi_k}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} X_k.$$

- We obtain

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \underbrace{\mathbb{E}_0 [X_i X_k]}_{P_{ik}} + O(\delta^2).$$

Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n - 1) \frac{\partial W_i}{\partial \phi_{in}} + (N - n) \frac{\partial W_i}{\partial \phi_{out}} = 0,$$

Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$

- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left(\underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$

Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$

- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left(\underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$

- ▶ Then further decompose with partial derivatives:

$$\frac{\partial W_i}{\partial \phi_k} = \sum_{\ell=1}^N \frac{\partial W_i}{\partial f_\ell} \frac{\partial f_\ell}{\partial \phi_k}$$

Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$


- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left(\underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$


- ▶ Then further decompose with partial derivatives:

$$\frac{\partial W_i}{\partial \phi_k} = \sum_{\ell=1}^N \frac{\partial W_i}{\partial f_\ell} \frac{\partial f_\ell}{\partial \phi_k} \quad \text{and} \quad \frac{\partial f_\ell}{\partial \phi_\ell} = -\text{c}; \quad \frac{\partial f_\ell}{\partial \phi_{\text{in}}} = \frac{\text{b}}{n-1}; \quad \frac{\partial f_\ell}{\partial \phi_{\text{out}}} = 0.$$


Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state




Genealogy, Identity by descent and Identity in state



Genealogy, Identity by descent and Identity in state



Genealogy, Identity by descent and Identity in state



Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

$$P_{ij}$$

Expected state
of the i,j pair

= Probability that the two
individuals are altruists

Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

$$P_{ij} = Q_{ij} \nu$$

Expected state
of the i, j pair
= Probability that the two
individuals are altruists

Probability that the individuals at
sites i and j are identical by descent
(no mutation since
their common ancestor)

Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

$$P_{ij} = Q_{ij} \nu$$

Expected state of the i, j pair
= Probability that the two individuals are altruists

Probability that a mutant is an altruist
= Probability that a given site is occupied by an altruist

Probability that the individuals at sites i and j are identical by descent
(no mutation since their common ancestor)

Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

$$P_{ij} = Q_{ij} \nu + (1 - Q_{ij}) \nu^2$$

Expected state of the i, j pair
= Probability that the two individuals are altruists

Probability that both sites are occupied by an altruist

Probability that the individuals at sites i and j are not identical by descent

```
graph TD; A["Pij = Qij ν + (1 - Qij) ν2"] --> B["Expected state of the i, j pair  
= Probability that the two individuals are altruists"]; A --> C["Probability that both sites are occupied by an altruist"]; A --> D["Probability that the individuals at sites i and j are not identical by descent"]
```

Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection, $\delta = 0$),

$$P_{ij} = Q_{ij} \nu + (1 - Q_{ij}) \nu^2$$

Expected state
of the i, j pair

= Probability that the two
individuals are altruists

Q_{in} , Q_{out}

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\begin{aligned}\mathbb{E}[\bar{X}] = & \nu + \delta \nu(1-\nu) \frac{1-\mu}{\mu} (1-Q_{\text{out}}) \times \\ & \left(-c - (b - c) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right. \\ & \left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n-1) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)\end{aligned}$$

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

Mutation-drift
equilibrium

$$\begin{aligned}\mathbb{E}[\bar{X}] = & \nu + \delta \nu(1-\nu) \frac{1-\mu}{\mu} (1-Q_{\text{out}}) \times \\ & \left(-c - (b - c) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right. \\ & \left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n-1) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)\end{aligned}$$

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

Mutation-drift equilibrium Selection strength

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \Big)$$

\mathcal{B}

Annotations:

- Mutation-drift equilibrium: Points to the term ν .
- Selection strength: Points to the term $\delta \nu(1 - \nu)$.
- Variance in the state of one site: Points to the term $\frac{1 - \mu}{\mu} (1 - Q_{\text{out}})$.

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right. \\ \left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

$R \qquad \mathcal{B}$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher


Mutation-drift equilibrium Selection strength Variance in the state of one site

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$


R \mathcal{B}

How does relatedness R change with the emigration probability m ?

Wright-Fisher (N deaths)




Moran (1 death)




$$(n = 4, N_d = 15)$$

How does relatedness R change with the emigration probability m ?

Wright-Fisher (N deaths)




Moran (1 death)




$$(n = 4, N_d = 15)$$

How does relatedness R change with the emigration probability m ?

Wright-Fisher (N deaths)



Moran (1 death)



$$(n = 4, N_d = 15)$$

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$

$R \searrow \mathcal{B}$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1-\nu) \frac{1-\mu}{\mu} (1-Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n-1) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$

$R \searrow \quad \mathcal{B}$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population

Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \nearrow \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n - 1) \left(\frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$

$R \searrow \quad \mathcal{B} \nearrow$

Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Expected frequency of altruists in the population


Equation for Moran Death-Birth and Wright-Fisher

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1-\nu) \frac{1-\mu}{\mu} (1-Q_{\text{out}}) \times$$
$$\left(-c - (b - c) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \nearrow \right. \\ \left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[b - (b - c)(n-1) \left(\frac{(1-m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

$R \searrow \quad \mathcal{B} \nearrow$


Mutation-drift equilibrium
Selection strength
Variance in the state of one site

Effect of the emigration probability m on the expected proportion of altruists




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists





$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists





$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists





$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists






$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists






$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists






$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists






$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists



$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Effect of the emigration probability m on the expected proportion of altruists




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

How to explain this result? (Moran Death-Birth)

$$-\mathcal{C} + \mathcal{B}\mathcal{R} > 0 \Leftrightarrow \mathcal{R} > \mathcal{C}/\mathcal{B}$$


How to explain this result? (Moran Death-Birth)

$$-\mathcal{C} + \mathcal{B}R > 0 \Leftrightarrow R > \mathcal{C}/\mathcal{B}$$




How to explain this result? (Moran Death-Birth)

$$-\mathcal{C} + \mathcal{B}R > 0 \Leftrightarrow R > \mathcal{C}/\mathcal{B}$$



Is the result robust?


Strong selection




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.005)$$

Strong selection


Wright-Fisher,
strong selection



Moran Death-Birth,
strong selection




Moran Death-Birth,
strong selection




$$(b = 15, c = 1, n = 4, N_d = 15, \delta = 0.1)$$

Heterogeneous deme sizes ($\bar{n} = 4$ as before, but $2 \leq n \leq 5$)


Wright-Fisher



Moran Death-Birth



Moran Death-Birth



$$(b = 15, c = 1, \bar{n} = 4, N_d = 15, \delta = 0.005)$$

Political implications



Take-Home Messages

- ▶ Under weak selection, it is possible to compute the expected frequency of social individuals, for any life-cycle, any regular population structure, any mutation probability. (D., 2017, JTB)

Take-Home Messages

- ▶ Under weak selection, it is possible to compute the expected frequency of social individuals, for any life-cycle, any regular population structure, any mutation probability. (D., 2017, JTB)
- ▶ $\mathbb{E}[\bar{\chi}] > \nu \Leftrightarrow \mathcal{B}R > \mathcal{C}$.

Take-Home Messages

- ▶ Under weak selection, it is possible to compute the expected frequency of social individuals, for any life-cycle, any regular population structure, any mutation probability. (D., 2017, JTB)
- ▶ $\mathbb{E}[\bar{X}] > \nu \Leftrightarrow \mathcal{B}R > \mathcal{C}$.
- ▶ In subdivided populations, $\mathbb{E}[\bar{X}]$ can increase with the emigration probability m when strategy transmission is imperfect ($\mu > 0$). (D., *in review.*)

Take-Home Messages

- ▶ Under weak selection, it is possible to compute the expected frequency of social individuals, for any life-cycle, any regular population structure, any mutation probability. (D., 2017, JTB)
- ▶ $\mathbb{E}[\bar{X}] > \nu \Leftrightarrow \mathcal{B}R > \mathcal{C}$.
- ▶ In subdivided populations, $\mathbb{E}[\bar{X}]$ can increase with the emigration probability m when strategy transmission is imperfect ($\mu > 0$). (D., *in review.*)
- ▶ This result seems to hold under stronger selection and in heterogeneous populations.

Take-Home Messages

- ▶ Under weak selection, it is possible to compute the expected frequency of social individuals, for any life-cycle, any regular population structure, any mutation probability. (D., 2017, JTB)
- ▶ $\mathbb{E}[\bar{X}] > \nu \Leftrightarrow \mathcal{B}R > \mathcal{C}$.
- ▶ In subdivided populations, $\mathbb{E}[\bar{X}]$ can increase with the emigration probability m when strategy transmission is imperfect ($\mu > 0$). (D., *in review.*)
- ▶ This result seems to hold under stronger selection and in heterogeneous populations.

Funding & Thanks



ANR-14-ACHN-0003-01

Amic Frouvelle & Emeric Bouin.
for organizing the workshop

and thank you for
your attention!