

Mon titre

1 Introduction

smaller groups, smaller emigration probabilities, both leading to increased within group relatedness are more conducive to the evolution of altruistic behavior. Living next to your kin however also means competing against them; the evolution of social traits hence depends on the balance between the positive effects of interactions with related individuals and the detrimental consequences of kin competition. With generations are synchronous (Wright-Fisher model), in infinite populations, Talor REF has shown that compensation + Gardner and Rodrigues.

Deriving analytical results often implies making simplifying assumptions. Include simple population structures (but see), weak selection approximations, and rare or absent mutation. Simple pop reduces the dimension / complexity of the system that one has to study; weak selection approximations allow a decomposition of time scales expliciter. Say what mutation means, fidelity of parent-offspring transmission. Here, we relax the assumption of rare or absent mutation and explore how imperfect strategy transmission from parents to their offspring affect the evolution of altruistic behavior in subdivided populations.

2 Model and methods

2.1 Assumptions

We consider a population of size N , subdivided into N_D demes, each hosting exactly n individuals (*i. e.*, containing n sites, each of which is occupied by exactly 1 individual; we have $nN_D = N$). Each site has a unique label i , $1 \leq i \leq N$. There are two types of individuals in the population, altruists and defectors. Reproduction is asexual. Parents transmit their strategy to their offspring with probability $1 - \mu$; this transmission can be genetic or cultural (vertical cultural transmission), but for simplicity, we refer to the parameter μ as a mutation probability. With probability μ , offspring do not inherit their strategy from their parent but instead get one randomly: with probability p , they become altruists, with probability $1 - p$ they become defectors. We call the parameter p the mutation bias.

Social interactions take place within each deme; each individual interacts with the $n - 1$ other deme members. We assume that social interactions affect individual fecundity, whose baseline is set to 1. Each interaction with an altruist increases an individual's fecundity by ωb , while altruists pay a fecundity cost ωc . The parameter ω scales the relative effect of social interactions on fecundity, and is assumed to be small ($\omega \ll 1$).

Denoting by e_{ij} the interaction probability between individuals living at sites i

37 and j , we have

$$e_{ij} = \begin{cases} 0 & \text{if } i = j; \\ \frac{1}{n-1} & \text{if } i \neq j \text{ and both sites are in the same deme;} \\ 0 & \text{if the two sites are in different demes.} \end{cases} \quad (1)$$

attention,
maybe
rather
{eq: defE}
1/(n-1)

38 Given our assumptions and with this notation, the fecundity of the individual
39 living at site k is given by

$$f_k(\mathbf{X}, \omega) = 1 + \omega \left(\sum_{\ell=1}^N e_{\ell k} b X_{\ell} - c X_k \right). \quad (2)$$

40 Although our assumptions may seem restrictive (unconditional benefits, addi-
41 tive effects), the same fecundities are obtained with a generic fecundity func-
42 tion, after linearization, under the assumption that altruists and defectors are
43 phenotypically close (see **APPENDIX** for details).

44 Offspring remain in the parental deme with probability $1 - m$; when they
45 do, they land on any site of the deme with equal probability (including the very
46 site of their parent). With probability m , offspring emigrate to a different deme,
47 chosen uniformly at random among the other demes. Denoting by d_{ij} the prob-
48 ability of moving from site i to site j , we have

$$d_{ij} = \begin{cases} \frac{1-m}{n} & \text{if both sites are in the same deme;} \\ \frac{m}{(N_D-1)n} & \text{if the two sites are in different demes.} \end{cases} \quad (3) \quad \{\text{eq: defD}\}$$

49 The way the population is updated from one time step to the next depends
50 on the chosen life-cycle (updating rule). We will specifically explore three dif-
51 ferent life-cycles. At the beginning of each step of each life-cycle, all individuals
52 produce offspring, that can be mutated; then these juveniles move, within the
53 parental deme or outside of it, and land on a site. The next events occurring
54 during the time step depend on the life-cycle:

55 **Moran Birth-Death** : One of the newly created juveniles is chosen at random; it
56 kills the adult who was living at the site, and replaces it; all other juveniles
57 die.

58 **Moran Death-Birth** : One of the adults is chosen to die (uniformly at random
59 among all adults). It is replaced by one of the juveniles who had landed in
60 its site. All other juveniles die.

61 **Wright-Fisher** : All the adults die. At each site of the entire population, one of
62 the juveniles that landed there is chosen and establishes at the site.

63 3 Results

64 3.1 Expected proportion of altruists

65 We want to compute the expected proportion of altruists in the population. Some
 66 steps can be done without specifying the life-cycle. We represent the state of the
 67 population at a given time t using indicator variables $X_i(t)$, $1 \leq i \leq N$, equal
 68 to 1 if the individual living at site i at time t is an altruist, and equal to 0 if it is
 69 a defector; these indicator variables are gathered in a N -long vector $\mathbf{X}(t)$. The
 70 set of all possible population states is $\Omega = \{0, 1\}^N$. The proportion of altruists in
 71 the population is written $\bar{X}(t) = \sum_{i=1}^N X_i(t)$. We denote by $B_{ji}(X(t), \omega)$, written
 72 B_{ji} for simplicity, the probability that the individual at site j at time $t+1$ is the
 73 newly established offspring of the individual living at site i at time t . We denote
 74 by $D_i(X(t), \omega)$ (D_i for simplicity) the probability that the individual living at site
 75 i at time t has been replaced (*i. e.*, died) at time $t+1$. Both quantities depend
 76 on the chosen life-cycle. Since a dead individual is immediately replaced by one
 77 new individual,

in a table?

$$D_i = \sum_{j=1}^N B_{ij} \quad (4a) \quad \{\text{eq:DBequiv}\}$$

78 holds for all sites i . The structure of the population is also such that in the ab-
 79 sence of selection ($\omega = 0$), all individuals have the same probability of dying and
 80 the same probability of having successful offspring (*i. e.*, offspring that become
 81 adults), so that

really
needed?

$$D_i^0 = \sum_{j=1}^N B_{ji}^0 = B^*, \quad (4b) \quad \{\text{eq:DBRV}\}$$

82 where the 0 subscript means that the quantities are evaluated for $\omega = 0$; this also
 83 implies that B_{ij}^0 and D_i^0 do not depend on the state \mathbf{X} of the population. For the
 84 Moran life-cycles, $B^* = 1/N$, while for the Wright-Fisher life-cycle, $B^* = 1$. (The
 85 difference with eq. (4a) is that we are now considering offspring produced by i
 86 landing on j).

87 Given that the population is in state $\mathbf{X}(t)$ at time t , the expected frequency of
 88 altruists at time $t+1$ is given by

$$\mathbb{E}[\bar{X}(t+1)|\mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N \left[\sum_{j=1}^N B_{ij} (X_j(1-\mu) + \mu p) + (1-D_i)X_i \right]. \quad (5a) \quad \{\text{eq:conditionalchange}\}$$

89 The first term within the brackets corresponds to births; the type of the indi-
 90 vidual living at i at time $t+1$ then depends on the type of its parent (living at
 91 site j), and on whether mutation occurred. The second term corresponds to the
 92 survival of the individual living at site i .

93 Given that there is no absorbing population state (a lost strategy can always
 94 be recreated by mutation), there is a stationary distribution of population states,
 95 and the expected frequency of altruists does not change anymore; we denote by
 96 $\xi(\mathbf{X}, \omega, \mu)$ the probability that the population is in state \mathbf{X} , given the strength of
 97 selection ω and the mutation probability μ . Taking the expectation of eq. (5a)
 98 ($\mathbb{E}[\bar{X}] = \sum_{\mathbf{X} \in \Omega} \bar{X} \xi(\mathbf{X}, \omega, \mu)$), we obtain, after reorganizing:

$$0 = \frac{1}{N} \sum_{\mathbf{X} \in \Omega} \sum_{i=1}^N \left[\sum_{j=1}^N B_{ij} (X_j(1-\mu) + \mu p) - D_i X_i \right] \xi(\mathbf{X}, \omega, \mu). \quad (6) \quad \{\text{eq:statdist}\}$$

99 Now, we use the assumption of weak selection ($\omega \ll 1$) and consider the first-
 100 order expansion of eq. (6) for ω close to 0. First, we note that in the absence
 101 of selection ($\omega = 0$), the population is at a mutation-drift balance, and the ex-
 102 pected state of every site i is then $\mathbb{E}_0[X_i] = \sum_{\mathbf{X} \in \Omega} X_i \xi(\mathbf{X}, 0, \mu) = p$, the mutation
 103 bias. Secondly, we further expand derivatives of B_{ji} and D_i using the chain rule,
 104 using the variables f_k ($1 \leq k \leq N$), corresponding to individual fecundities (also,
 105 recall that $f_k = 1$ when $\omega = 0$). Finally, we use the shorthand notation ∂_x to de-
 106 note $\frac{\partial}{\partial x} \Big|_{x=0}$. Thirdly, we note that for all the life-cycles that we consider, the
 107 number of deaths in the population during one time step does not depend on
 108 population composition (exactly 1 death for the Moran life-cycles, and exactly
 109 N for the Wright-Fisher life-cycle), so that $\partial_\omega \sum_{i,j=1}^N B_{ij}$ does not depend on ω .
 110 After simplification and reorganization, the first order expansion of eq. (6) yields
 111

$$\begin{aligned} 0 = \frac{1}{N} \sum_{i,k=1}^N \left[\frac{\partial \left(\sum_{j=1}^N (1-\mu) B_{ji} - D_i \right)}{\partial f_k} \Bigg|_{f_k=1} \right. \\ \left. \times \left(\sum_{\ell=1}^N e_{\ell k} \mathbf{b} \sum_{\mathbf{X} \in \Omega} X_\ell X_i \xi(\mathbf{X}, 0, \mu) - c \sum_{\mathbf{X} \in \Omega} X_k X_i \xi(\mathbf{X}, 0, \mu) \right) \right] \\ - B^* \mu \frac{\partial \mathbb{E}[\bar{X}]}{\partial \omega} \Bigg|_{\omega=0} + O(\omega^2). \end{aligned} \quad (7) \quad \{\text{eq:weaksel1}\}$$

112 The terms $\sum_{\mathbf{X} \in \Omega} X_i X_j \xi(\mathbf{X}, 0, \mu)$, that we will also denote by P_{ij} , correspond to
 113 the expected state of the pair of sites (i, j) , evaluated in the absence of selection
 114 ($\omega = 0$). We can also replace these terms by

$$Q_{ij} = \frac{P_{ij} - p^2}{p(1-p)}; \quad (8) \quad \{\text{eq:QP}\}$$

115 recursions on P_{ij} will reveal that Q_{ij} can be interpreted as a probability of iden-
 116 tity by descent, *i. e.*, the probability that the individuals at sites i and j have

117 a common ancestor and that no mutation has occurred on either lineage since
 118 the ancestor.

119 Finally, we obtain a first-order approximation of the expected frequency of
 120 altruists in the population with

$$\mathbb{E}[\bar{X}] = p + \omega \partial_{\omega} \mathbb{E}[\bar{X}] + O(\omega^2), \quad (9)$$

121 where $\partial_{\omega} \mathbb{E}[\bar{X}]$ is a shorthand notation for $\left. \frac{\partial \mathbb{E}[\bar{X}]}{\partial \omega} \right|_{\omega=0}$, which is given by eq. (7).

122 For each of the life-cycles that we consider, we can express $\partial_{\omega} \mathbb{E}[\bar{X}]$ as fol-
 123 lows:

$$\partial_{\omega} \mathbb{E}[\bar{X}] = b(\beta_D - \beta_I) - c(\gamma_D - \gamma_I), \quad (10)$$

124 where the subscript _D refers to “direct” effects, and the subscript _I to “indirect”
 125 effects. These indirect effects correspond to (kin) competition: by providing a
 126 benefit to a deme-mate and thereby increasing its fecundity, a focal altruist in-
 127 directly harms others by reducing their relative fecundity. Similarly, paying a
 128 fecundity cost indirectly helps others because it increases their relative fecundi-
 129 ties.

130 3.2 Identity by descent

131 We need to find equations for the expected state of pairs of sites (P_{ij}) and prob-
 132 abilities of identity by descent (Q_{ij}), quantities that are evaluated in the absence
 133 of selection (*i. e.*, for $\omega = 0$). To do so, we follow the same steps as in the pre-
 134 vious section: we first write expectations at the next time step given a current
 135 state, and we then take the expectation of this. Here we focus on identity by de-
 136 scent Q_{ij} , but expectations of the state of pairs of sites P_{ij} are simply recovered
 137 using eq. (8).


appendix

138 Because of the structure of the population, there are only three different val-
 139 ues of Q_{ij} :


$$Q_{ij} = \begin{cases} 1 & \text{when } i = j; \\ Q_{\text{in}} & \text{when } i \neq j \text{ and both sites are in the same deme;} \\ Q_{\text{out}} & \text{when sites } i \text{ and } j \text{ are in different demes.} \end{cases} \quad (11)$$

140 3.2.1 Moran model

(a) Moran



(b) Wright-Fisher



{fig:Q}
{fig:sub:QWF}

Figure 1: Probabilities of identity by descent, for two different individuals within the same deme (Q_{in} , full curves) and two individuals in different demes (Q_{out} , dashed curves), for different values of the mutation probability μ (10^{-4} , 0.005, 0.1), and for the two types of life-cycles: Moran (a) and Wright-Fisher (b). Other parameters: $n = 4$ individuals per deme, $N_D = 30$ demes.




Figure 2: Weak selection. Parameters: $\omega = 0.005$, $b = 15$, $c = 1$, **ndemes**, **size**, **nreps**. NOTE simulations running with 0.005 for μ and with 0.8 for m .




Figure 3: Strong selection

141 4 Figures




Figure 4: Weak selection, heterogeneous population




Figure 5: Strong selection, heterogeneous population

142 Adaptation of my equations to a subdivided population. Notation, for a
 143 quantity Y that depends on two sites ($Y = e, d, Q$):

$$Y_{\text{self}} := Y_{i,i} \quad (12a)$$

$$Y_{\text{in}} := Y_{i,j}, \quad i \text{ and } j \neq i \text{ in the same deme}; \quad (12b)$$

$$Y_{\text{out}} := Y_{i,j}, \quad i \text{ and } j \text{ in different demes.} \quad (12c)$$

144 For a site i , G_i denotes the deme the site belongs to, and notation $j \in G_i$ means
 145 that sites i and j are in the same deme.

146 The expected frequency of altruists in the population is given by

$$\mathbb{E}[\bar{X}] = p + \delta \frac{p(1-p)}{\mu} [b(\beta^D - \beta^I) - c(\gamma^D - \gamma^I)]. \quad (13)$$

Moran, Birth-Death

$$\begin{aligned} \beta_{\text{BD}}^D &= \sum_{k,\ell=1}^N \frac{1-\mu}{N} e_{k\ell} Q_{\ell k} \\ &= \sum_{k=1}^N \frac{1-\mu}{N} (e_{\text{self}} + (n-1)e_{\text{in}}Q_{\text{in}} + (N-n)e_{\text{out}}Q_{\text{out}}) \\ &= (1-\mu) (e_{\text{self}} + (n-1)e_{\text{in}}Q_{\text{in}} + (N-n)e_{\text{out}}Q_{\text{out}}). \end{aligned} \quad (14a)$$

$$\begin{aligned}
\beta_{\text{BD}}^I &= \sum_{j,k,l=1}^N \left(\frac{d_{lj}}{N} - \frac{\mu}{N^2} \right) e_{kl} Q_{jk} \\
&= \frac{1}{N} \sum_{j=1}^N \left[\left(\sum_{l=1}^N d_{lj} e_{jl} \right) + \sum_{\substack{k \in G_j \\ k \neq j}} \left(\sum_{l=1}^N d_{lj} e_{kl} Q_{\text{in}} Q_{\text{in}} \right) + \sum_{k \notin G_j} \sum_{l=1}^N d_{lj} (e_{kl} Q_{\text{out}} Q_{\text{out}}) \right] \\
&\quad + \frac{\mu}{N^2} \sum_{j=1}^N \left(\sum_{l=1}^N e_{kl} \right) \left(\sum_{k=1}^N Q_{jk} \right) \\
&= \frac{1}{N} \sum_{j=1}^N \left[d_{\text{self}} e_{\text{self}} + (n-1) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}} \right. \\
&\quad + \sum_{\substack{k \in G_j \\ k \neq j}} (d_{\text{in}} e_{\text{self}} + d_{\text{self}} e_{\text{in}} + (n-2) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}}) Q_{\text{in}} \\
&\quad + \sum_{k \notin G_j} (d_{\text{self}} e_{\text{out}} + (n-1) d_{\text{in}} e_{\text{out}} + d_{\text{out}} e_{\text{self}} + (n-1) d_{\text{out}} e_{\text{in}} + (N-2n) d_{\text{out}} e_{\text{out}}) Q_{\text{out}} \left. \right] \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) (e_{\text{self}} + (n-1) e_{\text{in}} + (N-n) e_{\text{out}}) \\
&= d_{\text{self}} e_{\text{self}} + (n-1) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}} \\
&\quad + (n-1) (d_{\text{in}} e_{\text{self}} + d_{\text{self}} e_{\text{in}} + (n-2) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}}) Q_{\text{in}} \\
&\quad + (N-n) (d_{\text{self}} e_{\text{out}} + (n-1) d_{\text{in}} e_{\text{out}} + d_{\text{out}} e_{\text{self}} + (n-1) d_{\text{out}} e_{\text{in}} + (N-2n) d_{\text{out}} e_{\text{out}}) Q_{\text{out}} \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) (e_{\text{self}} + (n-1) e_{\text{in}} + (N-n) e_{\text{out}}). \quad (14b)
\end{aligned}$$

$$\gamma_{\text{BD}}^D = 1 - \mu. \quad (14c)$$

$$\begin{aligned}
\gamma_{\text{BD}}^I &= \frac{1}{N} \sum_{j,k=1}^N \left(d_{kj} - \frac{\mu}{N} \right) Q_{jk} \\
&= \frac{1}{N} \sum_{j=1}^N \left[d_{\text{self}} - \frac{\mu}{N} + (n-1) \left(d_{\text{in}} - \frac{\mu}{N} \right) Q_{\text{in}} + (N-n) \left(d_{\text{out}} - \frac{\mu}{N} \right) Q_{\text{out}} \right] \\
&= d_{\text{self}} + (n-1) d_{\text{in}} Q_{\text{in}} + (N-n) d_{\text{out}} Q_{\text{out}} \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) \quad (14d)
\end{aligned}$$

Moran, Death-Birth

$$\begin{aligned}\beta_{\text{DB}}^D &= \frac{1-\mu}{N} \sum_{j,k=1}^N Q_{jk} e_{jk} = \beta_{\text{BD}}^D \\ &= (1-\mu) \left(e_{\text{self}} + (n-1) e_{\text{in}} Q_{\text{in}} + (N-n) e_{\text{out}} Q_{\text{out}} \right).\end{aligned}\quad (15a)$$

$$\beta_{\text{DB}}^I = \frac{1-\mu}{N} \sum_{i,j,k,l=1}^N d_{ji} d_{li} e_{kl} Q_{jk} \quad (15b)$$

147 Presented in the table in the appendix.

$$\gamma_{\text{DB}}^D = 1 - \mu = \gamma_{\text{BD}}^D. \quad (15c)$$

$$\begin{aligned}\gamma_{\text{DB}}^I &= (1-\mu) \sum_{i,j,k=1}^N \frac{d_{ji} d_{ki}}{N} Q_{jk} \\ &= \frac{1-\mu}{N} \sum_{j=1}^N \sum_{i=1}^N \left(d_{ji} d_{ji} + \sum_{\substack{k \neq j \\ k \in G_j}} d_{ji} d_{ki} Q_{\text{in}} + \sum_{k \notin G_j} d_{ji} d_{ki} Q_{\text{out}} \right) \\ &= \frac{1-\mu}{N} \sum_{j=1}^N \left[d_{\text{self}} d_{\text{self}} + (n-1) d_{\text{in}} d_{\text{in}} + (N-n) d_{\text{out}} d_{\text{out}} \right. \\ &\quad \left. + (n-1) \left(d_{\text{self}} d_{\text{in}} + d_{\text{in}} d_{\text{self}} + (n-2) d_{\text{in}} d_{\text{in}} + (N-n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{in}} \right. \\ &\quad \left. + (N-n) \left(d_{\text{self}} d_{\text{out}} + (n-1) d_{\text{in}} d_{\text{out}} + d_{\text{out}} d_{\text{self}} + (n-1) d_{\text{out}} d_{\text{in}} + (N-2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \right] \\ &\quad (15d)\end{aligned}$$

148 Probabilities of identity by descent

149 WF est faux. Il faut utiliser les formules Fourier...!

150 **Moran** For $i \neq j$,

$$Q_{ij} = \frac{1-\mu}{2} \sum_{k=1}^N (d_{kj} Q_{ki} + d_{ki} Q_{kj}). \quad (16a)$$

151 For $j \neq i, j \in G_i$,

$$\begin{aligned}
Q_{\text{in}} &= \frac{1-\mu}{2} \left((d_{\text{in}} + d_{\text{self}} Q_{\text{in}}) + (d_{\text{self}} Q_{\text{in}} + d_{\text{in}}) \right. \\
&\quad \left. + (n-2)(d_{\text{in}} Q_{\text{in}} + d_{\text{in}} Q_{\text{in}}) + (N-n)(d_{\text{out}} Q_{\text{out}} + d_{\text{out}} Q_{\text{out}}) \right) \\
&= (1-\mu) \left(d_{\text{in}} + d_{\text{self}} Q_{\text{in}} + (n-2)d_{\text{in}} Q_{\text{in}} + (N-n)d_{\text{out}} Q_{\text{out}} \right). \tag{16b}
\end{aligned}$$

152 And for $j \notin G_i$,

$$\begin{aligned}
Q_{\text{out}} &= \frac{1-\mu}{2} \left((d_{\text{out}} + d_{\text{self}} Q_{\text{out}}) + (n-1)(d_{\text{out}} Q_{\text{in}} + d_{\text{in}} Q_{\text{out}}) \right. \\
&\quad \left. + (d_{\text{self}} Q_{\text{out}} + d_{\text{out}}) + (n-1)(d_{\text{in}} Q_{\text{out}} + d_{\text{out}} Q_{\text{in}}) \right. \\
&\quad \left. + (N-2n)(d_{\text{out}} Q_{\text{out}} + d_{\text{out}} Q_{\text{out}}) \right) \\
&= (1-\mu) \left(d_{\text{out}} + d_{\text{self}} Q_{\text{out}} + (n-1)(d_{\text{out}} Q_{\text{in}} + d_{\text{in}} Q_{\text{out}}) + (N-2n)d_{\text{out}} Q_{\text{out}} \right) \tag{16c}
\end{aligned}$$

153 **Wright-Fisher** For $j \neq i$,

$$Q_{ij} = (1-\mu)^2 \sum_{k,l=1}^N d_{ki} d_{lj} Q_{kl}. \tag{17a}$$

154 When $j \neq i$, $j \in G_i$,

$$\begin{aligned}
Q_{\text{in}} &= (1 - \mu)^2 \left[\left(d_{\text{self}} d_{\text{in}} + d_{\text{in}} d_{\text{self}} + (n - 2) d_{\text{in}} d_{\text{in}} + (N - n) d_{\text{out}} d_{\text{out}} \right) \right. \\
&\quad + \left(d_{\text{self}} d_{\text{self}} + (n - 2) d_{\text{self}} d_{\text{in}} \right. \\
&\quad \quad + (n - 1) d_{\text{in}} d_{\text{in}} + (n - 2) d_{\text{in}} d_{\text{self}} \\
&\quad \quad \left. + (n - 2)(n - 2) d_{\text{in}} d_{\text{in}} + (N - n)(n - 1) d_{\text{out}} d_{\text{out}} \right) Q_{\text{in}} \\
&\quad + \left((N - n) d_{\text{self}} d_{\text{out}} + (N - n)(n - 1) d_{\text{in}} d_{\text{out}} \right. \\
&\quad \quad + (N - n) d_{\text{out}} d_{\text{self}} + (N - n)(n - 1) d_{\text{out}} d_{\text{in}} \\
&\quad \quad \left. + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big] \\
&= (1 - \mu)^2 \left[\left(2 d_{\text{in}} d_{\text{self}} + (n - 2) d_{\text{in}}^2 + (N - n) d_{\text{out}}^2 \right) \right. \\
&\quad + \left(d_{\text{self}}^2 + 2(n - 2) d_{\text{self}} d_{\text{in}} + (n^2 - 3n + 3) d_{\text{in}}^2 + (N - n)(n - 1) d_{\text{out}}^2 \right) Q_{\text{in}} \\
&\quad + \left(2(N - n) d_{\text{self}} d_{\text{out}} + 2(N - n)(n - 1) d_{\text{in}} d_{\text{out}} \right. \\
&\quad \quad \left. + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big] \tag{17b}
\end{aligned}$$

155 And when $j \notin G_i$, we have

$$\begin{aligned}
Q_{\text{out}} &= (1 - \mu)^2 \left[\left(2 d_{\text{self}} d_{\text{out}} + 2(n - 1) d_{\text{in}} d_{\text{out}} + (N - 2n) d_{\text{out}}^2 \right) \right. \\
&\quad + \left(2(n - 1) d_{\text{self}} d_{\text{out}} + 2(n - 1)^2 d_{\text{in}} d_{\text{out}} + (N - 2n)(n - 1) d_{\text{out}}^2 \right) Q_{\text{in}} \\
&\quad + \left(d_{\text{self}} d_{\text{self}} + (n - 1) d_{\text{self}} d_{\text{in}} + (N - 2n) d_{\text{self}} d_{\text{out}} \right. \\
&\quad \quad + (n - 1) d_{\text{in}} d_{\text{self}} + (n - 1)^2 d_{\text{in}}^2 + (n - 1)(N - 2n) d_{\text{in}} d_{\text{out}} \\
&\quad \quad \left. + (N - n) d_{\text{out}} d_{\text{self}} + (N - n)(n - 1) d_{\text{out}} d_{\text{in}} + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big]. \tag{17c}
\end{aligned}$$

156 **PAS FINI**

157 **Appendix**

158 All combinations for i, j, k, l . Notation: (i, j) means that i and j are in the same
159 deme, but are different; G_i refers to the deme containing site i .

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
1	$j = i$	$k = i$	$l = i$	$(i = j = k = l)$	1	d_{self}	d_{self}	e_{self}	1
2	$j = i$	$k = i$	$l \neq i; l \in G_i$	$(i = j = k, l)$	$n - 1$	d_{self}	d_{in}	e_{in}	1
3	$j = i$	$k = i$	$l \notin G_i$	$(i = j = k), (l)$	$N - n$	d_{self}	d_{out}	e_{out}	1
4	$j = i$	$k \neq i; k \in G_i$	$l = i$	$(i = j = l, k)$	$n - 1$	d_{self}	d_{self}	e_{in}	Q_{in}
5	$j = i$	$k \neq i; k \in G_i$	$l = k$	$(i = j, k = l)$	$n - 1$	d_{self}	d_{in}	e_{self}	Q_{in}
6	$j = i$	$k \neq i; k \in G_i$	$l \neq i, k; l \in G_i$	$(i = j, k, l)$	$(n - 1)(n - 2)$	d_{self}	d_{in}	e_{in}	Q_{in}
7	$j = i$	$k \neq i; k \in G_i$	$l \notin G_i$	$(i = j, k), (l)$	$(n - 1)(N - n)$	d_{self}	d_{out}	e_{out}	Q_{in}
8	$j = i$	$k \notin G_i$	$l = i = j$	$(i = j = l), (k)$	$(N - n)$	d_{self}	d_{self}	e_{out}	Q_{out}
9	$j = i$	$k \notin G_i$	$l \neq i, l \in G_i$	$(i = j, l), (k)$	$(N - n)(n - 1)$	d_{self}	d_{in}	e_{out}	Q_{out}
10	$j = i$	$k \notin G_i$	$l = k$	$(i = j), (k = l)$	$(N - n)$	d_{self}	d_{out}	e_{self}	Q_{out}
11	$j = i$	$k \notin G_i$	$l \neq k; l \in G_k$	$(i = j), (k, l)$	$(N - n)(n - 1)$	d_{self}	d_{out}	e_{in}	Q_{out}
12	$j = i$	$k \notin G_i$	$l \notin G_i, G_k$	$(i = j), (k), (l)$	$(N - n)(N - 2n)$	d_{self}	d_{out}	e_{out}	Q_{out}
13	$j \neq i, j \in G_i$	$k = i$	$l = i$	$(i = k = l, j)$	$(n - 1)$	d_{in}	d_{self}	e_{self}	Q_{in}
14	$j \neq i, j \in G_i$	$k = i$	$l = j$	$(i = k, j = l)$	$(n - 1)$	d_{in}	d_{in}	e_{in}	Q_{in}
15	$j \neq i, j \in G_i$	$k = i$	$l \neq i, j; l \in G_i$	$(i = k, j, l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	Q_{in}
16	$j \neq i, j \in G_i$	$k = i$	$l \notin G_i$	$(i = k, j), (l)$	$(n - 1)(N - n)$	d_{in}	d_{out}	e_{out}	Q_{in}
17	$j \neq i, j \in G_i$	$k = j$	$l = i$	$(i = l, j = k)$	$(n - 1)$	d_{in}	d_{self}	e_{in}	1
18	$j \neq i, j \in G_i$	$k = j$	$l = j$	$(i, j = k = l)$	$(n - 1)$	d_{in}	d_{in}	e_{self}	1
19	$j \neq i, j \in G_i$	$k = j$	$l \neq i, j; l \in G_i$	$(i, j = k, l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	1
20	$j \neq i, j \in G_i$	$k = j$	$l \notin G_i$	$(i, j = k), (l)$	$(n - 1)(N - n)$	d_{in}	d_{out}	e_{out}	1
21	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = i$	$(i = l, j, k)$	$(n - 1)(n - 2)$	d_{in}	d_{self}	e_{in}	Q_{in}
22	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = j$	$(i, j = l, k)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	Q_{in}
23	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = k$	$(i, j, k = l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{self}	Q_{in}
24	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l \neq i, j, k; l \in G_i$	(i, j, k, l)	$(n - 1)(n - 2)(n - 3)$	d_{in}	d_{in}	e_{in}	Q_{in}
25	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l \notin G_i$	$(i, j, k), (l)$	$(n - 1)(n - 2)(N - n)$	d_{in}	d_{out}	e_{out}	Q_{in}

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
26	$j \neq i; j \in G_i$	$k \notin G_i$	$l = i$	$(i = l, j), (k)$	$(n-1)(N-n)$	d_{in}	d_{self}	e_{out}	Q_{out}
27	$j \neq i; j \in G_i$	$k \notin G_i$	$l = j$	$(i, j = l), (k)$	$(n-1)(N-n)$	d_{in}	d_{in}	e_{out}	Q_{out}
28	$j \neq i; j \in G_i$	$k \notin G_i$	$l \neq i, j; l \in G_i$	$(i, j, l), (k)$	$(n-1)(N-n)(n-2)$	d_{in}	d_{in}	e_{out}	Q_{out}
29	$j \neq i; j \in G_i$	$k \notin G_i$	$l = k$	$(i, j), (k = l)$	$(n-1)(N-n)$	d_{in}	d_{out}	e_{self}	Q_{out}
30	$j \neq i; j \in G_i$	$k \notin G_i$	$l \neq k; l \in G_k$	$(i, j), (k, l)$	$(n-1)(N-n)(n-1)$	d_{in}	d_{out}	e_{in}	Q_{out}
31	$j \neq i; j \in G_i$	$k \notin G_i$	$l \notin G_i, G_k$	$(i, j), (k), (l)$	$(n-1)(N-n)(N-2n)$	d_{in}	d_{out}	e_{out}	Q_{out}
32	$j \notin G_i$	$k = i$	$l = i$	$(i = k = l), (j)$	$(N-n)$	d_{out}	d_{self}	e_{self}	Q_{out}
33	$j \notin G_i$	$k = i$	$l \neq i; l \in G_i$	$(i = k, l), (j)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{in}	Q_{out}
34	$j \notin G_i$	$k = i$	$l = j$	$(i = k), (j = l)$	$(N-n)$	d_{out}	d_{out}	e_{out}	Q_{out}
35	$j \notin G_i$	$k = i$	$l \neq j; l \in G_j$	$(i = k), (j, l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
36	$j \notin G_i$	$k = i$	$l \notin G_i, G_j$	$(i = k), (j), (l)$	$(N-n)(N-2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
37	$j \notin G_i$	$k \neq i; k \in G_i$	$l = i$	$(i = l, k), (j)$	$(N-n)(n-1)$	d_{out}	d_{self}	e_{in}	Q_{out}
38	$j \notin G_i$	$k \neq i; k \in G_i$	$l = k$	$(i, k = l), (j)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{self}	Q_{out}
39	$j \notin G_i$	$k \neq i; k \in G_i$	$l \neq i, k; l \in G_i$	$(i, k, l), (j)$	$(N-n)(n-1)(n-2)$	d_{out}	d_{in}	e_{in}	Q_{out}
40	$j \notin G_i$	$k \neq i; k \in G_i$	$l = j$	$(i, k), (j = l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
41	$j \notin G_i$	$k \neq i; k \in G_i$	$l \neq j; l \in G_j$	$(i, k), (j, l)$	$(N-n)(n-1)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
42	$j \notin G_i$	$k \neq i; k \in G_i$	$l \notin G_i, G_j$	$(i, k), (j), (l)$	$(N-n)(n-1)(N-2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
43	$j \notin G_i$	$k = j$	$l = i$	$(i = l), (j = k)$	$(N-n)$	d_{out}	d_{self}	e_{out}	1
44	$j \notin G_i$	$k = j$	$l \neq i; l \in G_i$	$(i, l), (j = k)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{out}	1
45	$j \notin G_i$	$k = j$	$l = j$	$(i), (j = k = l)$	$(N-n)$	d_{out}	d_{out}	e_{self}	1
46	$j \notin G_i$	$k = j$	$l \neq j; l \in G_j$	$(i), (j = k, l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{in}	1
47	$j \notin G_i$	$k = j$	$l \notin G_i, G_j$	$(i), (j = k), (l)$	$(N-n)(N-2n)$	d_{out}	d_{out}	e_{out}	1

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
48	$j \notin G_i$	$k \neq j; k \in G_j$	$l = i$	$(i = l), (j, k)$	$(N - n)(n - 1)$	d_{out}	d_{self}	e_{out}	Q_{in}
49	$j \notin G_i$	$k \neq j; k \in G_j$	$l \neq i; l \in G_i$	$(i, l), (j, k)$	$(N - n)(n - 1)(n - 1)$	d_{out}	d_{in}	e_{out}	Q_{in}
50	$j \notin G_i$	$k \neq j; k \in G_j$	$l = j$	$(i), (j = l, k)$	$(N - n)(n - 1)$	d_{out}	d_{out}	e_{in}	Q_{in}
51	$j \notin G_i$	$k \neq j; k \in G_j$	$l = k$	$(i), (j, k = l)$	$(N - n)(n - 1)$	d_{out}	d_{out}	e_{self}	Q_{in}
52	$j \notin G_i$	$k \neq j; k \in G_j$	$l \neq j, k; l \in G_j$	$(i), (j, k, l)$	$(N - n)(n - 1)(n - 2)$	d_{out}	d_{out}	e_{in}	Q_{in}
53	$j \notin G_i$	$k \neq j; k \in G_j$	$l \notin G_i, G_j$	$(i), (j, k), (l)$	$(N - n)(n - 1)(N - 2n)$	d_{out}	d_{out}	e_{out}	Q_{in}
54	$j \notin G_i$	$k \notin G_i, G_j$	$l = i$	$(i = l), (j), (k)$	$(N - n)(N - 2n)$	d_{out}	d_{self}	e_{out}	Q_{out}
55	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq i; l \in G_i$	$(i, l), (j), (k)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{in}	e_{out}	Q_{out}
56	$j \notin G_i$	$k \notin G_i, G_j$	$l = j$	$(i), (j = l), (k)$	$(N - n)(N - 2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
57	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq j; l \in G_j$	$(i), (j, l), (k)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{out}	e_{out}	Q_{out}
58	$j \notin G_i$	$k \notin G_i, G_j$	$l = k$	$(i), (j), (k = l)$	$(N - n)(N - 2n)$	d_{out}	d_{out}	e_{self}	Q_{out}
59	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq k; l \in G_k$	$(i), (j), (k, l)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{out}	e_{in}	Q_{out}
60	$j \notin G_i$	$k \notin G_i, G_j$	$l \notin G_i, G_j, G_k$	$(i), (j), (k), (l)$	$(N - n)(N - 2n)(N - 3n)$	d_{out}	d_{out}	e_{out}	Q_{out}

160 **A Island model**

161 With self replacement

$$d_{\text{self}} = d_{\text{in}} = \frac{1-m}{n}, \quad (18a)$$

$$d_{\text{out}} = \frac{m}{N-n}. \quad (18b)$$

162 Without self-replacement

$$d_{\text{self}} = 0, \quad (19a)$$

$$d_{\text{in}} = \frac{1-m}{n-1}, \quad (19b)$$

$$d_{\text{out}} = \frac{m}{N-n}. \quad (19c)$$

163 **B IDB**

164 **B.1 Moran**

165 Using the formulas for a 2D graph in REF Debarre 2017,

$$\tilde{\mathcal{D}}_{q_1} = \sum_{q_2} \sum_{l_1=0}^{N_1-1} \sum_{l_2=0}^{N_2-1} \tilde{d}_{l_1} \exp\left(-i \frac{2\pi q_1 l_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) \quad (20a)$$

$$\tilde{\mathcal{Q}}_{r_1} = \frac{1}{N} \sum_{q_1=0}^{N_1-1} \sum_{q_2=0}^{N_2-1} \frac{\mu \lambda'_M}{1 - (1-\mu) \tilde{\mathcal{D}}_{q_1}} \exp\left(i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(i \frac{2\pi q_2 r_2}{N_2}\right) \quad (20b)$$

166 We have

$$\begin{aligned} \tilde{\mathcal{D}}_{q_1} &= d_{\text{self}} + \sum_{l_2=1}^{N_2-1} d_{\text{in}} \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) + \sum_{l_1=1}^{N_1-1} \sum_{l_2=0}^{N_2-1} d_{\text{out}} \exp\left(-i \frac{2\pi q_1 l_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) \\ &= d_{\text{self}} + (\delta_{q_2}(N_2-1) + (1-\delta_{q_2})(-1)) d_{\text{in}} + (\delta_{q_1}(N_1-1) + (1-\delta_{q_1})(-1)) (\delta_{q_2} N_2) d_{\text{out}} \\ &= d_{\text{self}} + (\delta_{q_2} N_2 - 1) d_{\text{in}} + (\delta_{q_1} N_1 - 1) \delta_{q_2} N_2 d_{\text{out}}. \end{aligned} \quad (21a)$$

167 Whether there is self-replacement or not, we have $N_1 = D$ and $N_2 = n$, and

$$\tilde{\mathcal{D}}_0 = 1, \quad (22a)$$

$$\tilde{\mathcal{D}}_{q_1} = 1 - m - \frac{m}{d-1} \quad (q_1 \not\equiv 0 \pmod{N_1}), \quad (22b)$$

$$\tilde{\mathcal{D}}_{q_1} = d_{\text{self}} - d_{\text{in}} \quad (q_2 \not\equiv 0 \pmod{N_2}). \quad (22c)$$

168 So for \tilde{Q} ,

$$\begin{aligned}
\tilde{Q}_{r_1, r_2} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{1 - (1 - \mu) \tilde{D}_0} + \sum_{q_2=1}^{N_2-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_2}} \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) + \sum_{q_1=1}^{N_1-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_1}} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \right. \\
&\quad \left. + \sum_{q_1=1}^{N_1-1} \sum_{q_2=1}^{N_2-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_1, q_2}} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) \right] \\
&= \frac{\mu \lambda'_M}{N} \left[\frac{1}{1 - (1 - \mu)} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (\delta_{r_2} N_2 - 1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (\delta_{r_1} N_1 - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (\delta_{r_1} N_1 - 1)(\delta_{r_2} N_2 - 1) \right]. \tag{23a}
\end{aligned}$$

169 In particular,

$$\begin{aligned}
\tilde{Q}_0 &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (n - 1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (D - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (D - 1)(n - 1) \right] \\
&= 1. \tag{23b}
\end{aligned}$$

170 We find λ'_M using the above equation. When $r_1 = 0$, the two individuals are in
171 the same deme. They are different when $r_2 \neq 0$:

$$\begin{aligned}
Q_{\text{in}} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (-1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (D - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (D - 1)(-1) \right]. \tag{23c}
\end{aligned}$$

172 And when $r_1 \neq 0$, the two individuals are in different demes:

$$\begin{aligned}
Q_{\text{out}} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (-1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (-1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} \right]. \tag{23d}
\end{aligned}$$

B.2 Wright-Fisher

$$\begin{aligned}
\tilde{Q}_{r_1, r_2} &= \frac{1}{N} \sum_{q_1=0}^{N_1-1} \sum_{q_2=0}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_{q_1})^2} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) \\
&= \frac{1}{N} \left[\frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0)^2} + \sum_{q_2=1}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0)^2} \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) \right. \\
&\quad + \sum_{q_1=1}^{N_1-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0)^2} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \\
&\quad \left. + \sum_{q_1=1}^{N_1-1} \sum_{q_2=1}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_{q_1})^2} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) \right] \quad (24)
\end{aligned}$$

$$\begin{aligned}
&= \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_2} N_2 - 1) \right. \\
&\quad + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (\delta_{q_1} N_1 - 1) \\
&\quad \left. + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_1} N_1 - 1) (\delta_{q_2} N_2 - 1) \right] \\
&= \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_2} N_2 - 1) \delta_{q_1} N_1 \right. \\
&\quad \left. + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (\delta_{q_1} N_1 - 1) \right]. \quad (25)
\end{aligned}$$

174 To find λ'_{WF} , we solve

$$1 = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (N_2 - 1) N_1 + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (N_1 - 1) \right]. \quad (26a)$$

175 Then,

$$Q_{\text{in}} = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} - \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} N_1 + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (N_1 - 1) \right]. \quad (26b)$$

176 and

$$Q_{\text{out}} = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} - \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} \right]. \quad (26c)$$