

Adaptation of my equations to a subdivided population. Notation, for a quantity Y that depends on two sites ($Y = e, d, Q$):

$$Y_{\text{self}} = Y_{i,i} \quad (1a)$$

$$Y_{\text{in}} = Y_{i,j}, \quad i \text{ and } j \neq i \text{ in the same deme}; \quad (1b)$$

$$Y_{\text{out}} = Y_{i,j}, \quad i \text{ and } j \text{ in different demes}. \quad (1c)$$

For a site i , G_i denotes the deme it is in, and notation $j \in G_i$ means that sites i and j are in the same deme.

The expected frequency of altruists in the population is given by

$$\mathbb{E}[\bar{X}] = p + \delta \frac{p(1-p)}{\mu} [b(\beta^D - \beta^I) - c(\gamma^D - \gamma^I)]. \quad (2)$$

Moran, Birth-Death

$$\begin{aligned} \beta_{\text{BD}}^D &= \sum_{k,\ell=1}^N \frac{1-\mu}{N} e_{k\ell} Q_{\ell k} \\ &= \sum_{k=1}^N \frac{1-\mu}{N} (e_{\text{self}} + (n-1)e_{\text{in}}Q_{\text{in}} + (N-n)e_{\text{out}}Q_{\text{out}}) \\ &= (1-\mu) (e_{\text{self}} + (n-1)e_{\text{in}}Q_{\text{in}} + (N-n)e_{\text{out}}Q_{\text{out}}). \end{aligned} \quad (3a)$$

$$\begin{aligned}
\beta_{\text{BD}}^I &= \sum_{j,k,l=1}^N \left(\frac{d_{lj}}{N} - \frac{\mu}{N^2} \right) e_{kl} Q_{jk} \\
&= \frac{1}{N} \sum_{j=1}^N \left[\left(\sum_{l=1}^N d_{lj} e_{jl} \right) + \sum_{\substack{k \in G_j \\ k \neq j}} \left(\sum_{l=1}^N d_{lj} e_{kl} Q_{\text{in}} Q_{\text{in}} \right) + \sum_{k \notin G_j} \sum_{l=1}^N d_{lj} (e_{kl} Q_{\text{out}} Q_{\text{out}}) \right] \\
&\quad + \frac{\mu}{N^2} \sum_{j=1}^N \left(\sum_{l=1}^N e_{kl} \right) \left(\sum_{k=1}^N Q_{jk} \right) \\
&= \frac{1}{N} \sum_{j=1}^N \left[d_{\text{self}} e_{\text{self}} + (n-1) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}} \right. \\
&\quad + \sum_{\substack{k \in G_j \\ k \neq j}} (d_{\text{in}} e_{\text{self}} + d_{\text{self}} e_{\text{in}} + (n-2) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}}) Q_{\text{in}} \\
&\quad + \sum_{k \notin G_j} (d_{\text{self}} e_{\text{out}} + (n-1) d_{\text{in}} e_{\text{out}} + d_{\text{out}} e_{\text{self}} + (n-1) d_{\text{out}} e_{\text{in}} + (N-2n) d_{\text{out}} e_{\text{out}}) Q_{\text{out}} \left. \right] \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) (e_{\text{self}} + (n-1) e_{\text{in}} + (N-n) e_{\text{out}}) \\
&= d_{\text{self}} e_{\text{self}} + (n-1) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}} \\
&\quad + (n-1) (d_{\text{in}} e_{\text{self}} + d_{\text{self}} e_{\text{in}} + (n-2) d_{\text{in}} e_{\text{in}} + (N-n) d_{\text{out}} e_{\text{out}}) Q_{\text{in}} \\
&\quad + (N-n) (d_{\text{self}} e_{\text{out}} + (n-1) d_{\text{in}} e_{\text{out}} + d_{\text{out}} e_{\text{self}} + (n-1) d_{\text{out}} e_{\text{in}} + (N-2n) d_{\text{out}} e_{\text{out}}) Q_{\text{out}} \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) (e_{\text{self}} + (n-1) e_{\text{in}} + (N-n) e_{\text{out}}). \quad (3b)
\end{aligned}$$

$$\gamma_{\text{BD}}^D = 1 - \mu. \quad (3c)$$

$$\begin{aligned}
\gamma_{\text{BD}}^I &= \frac{1}{N} \sum_{j,k=1}^N \left(d_{kj} - \frac{\mu}{N} \right) Q_{jk} \\
&= \frac{1}{N} \sum_{j=1}^N \left[d_{\text{self}} - \frac{\mu}{N} + (n-1) \left(d_{\text{in}} - \frac{\mu}{N} \right) Q_{\text{in}} + (N-n) \left(d_{\text{out}} - \frac{\mu}{N} \right) Q_{\text{out}} \right] \\
&= d_{\text{self}} + (n-1) d_{\text{in}} Q_{\text{in}} + (N-n) d_{\text{out}} Q_{\text{out}} \\
&\quad - \frac{\mu}{N} (1 + (n-1) Q_{\text{in}} + (N-n) Q_{\text{out}}) \quad (3d)
\end{aligned}$$

Moran, Death-Birth

$$\begin{aligned}\beta_{\text{DB}}^D &= \frac{1-\mu}{N} \sum_{j,k=1}^N Q_{jk} e_{jk} = \beta_{\text{BD}}^D \\ &= (1-\mu) \left(e_{\text{self}} + (n-1) e_{\text{in}} Q_{\text{in}} + (N-n) e_{\text{out}} Q_{\text{out}} \right).\end{aligned}\quad (4a)$$

$$\beta_{\text{DB}}^I = \frac{1-\mu}{N} \sum_{i,j,k,l=1}^N d_{ji} d_{li} e_{kl} Q_{jk} \quad (4b)$$

Presented in the table in the appendix.

$$\gamma_{\text{DB}}^D = 1 - \mu = \gamma_{\text{BD}}^D. \quad (4c)$$

$$\begin{aligned}\gamma_{\text{DB}}^I &= (1-\mu) \sum_{i,j,k=1}^N \frac{d_{ji} d_{ki}}{N} Q_{jk} \\ &= \frac{1-\mu}{N} \sum_{j=1}^N \sum_{i=1}^N \left(d_{ji} d_{ji} + \sum_{\substack{k \neq j \\ k \in G_j}} d_{ji} d_{ki} Q_{\text{in}} + \sum_{k \notin G_j} d_{ji} d_{ki} Q_{\text{out}} \right) \\ &= \frac{1-\mu}{N} \sum_{j=1}^N \left[d_{\text{self}} d_{\text{self}} + (n-1) d_{\text{in}} d_{\text{in}} + (N-n) d_{\text{out}} d_{\text{out}} \right. \\ &\quad \left. + \left(d_{\text{self}} d_{\text{in}} + d_{\text{in}} d_{\text{self}} + (n-2) d_{\text{in}} d_{\text{in}} + (N-n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{in}} \right. \\ &\quad \left. + \left(d_{\text{self}} d_{\text{out}} + (n-1) d_{\text{in}} d_{\text{out}} + d_{\text{out}} d_{\text{self}} + (n-1) d_{\text{out}} d_{\text{in}} + (N-2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \right] \end{aligned} \quad (4d)$$

Probabilities of identity by descent

WF est faux. Il faut utiliser les formules Fourier...!

Moran For $i \neq j$,

$$Q_{ij} = \frac{1-\mu}{2} \sum_{k=1}^N (d_{kj} Q_{ki} + d_{ki} Q_{kj}). \quad (5a)$$

For $j \neq i, j \in G_i$,

$$\begin{aligned}
Q_{\text{in}} &= \frac{1-\mu}{2} \left((d_{\text{in}} + d_{\text{self}} Q_{\text{in}}) + (d_{\text{self}} Q_{\text{in}} + d_{\text{in}}) \right. \\
&\quad \left. + (n-2) (d_{\text{in}} Q_{\text{in}} + d_{\text{in}} Q_{\text{in}}) + (N-n) (d_{\text{out}} Q_{\text{out}} + d_{\text{out}} Q_{\text{out}}) \right) \\
&= (1-\mu) \left(d_{\text{in}} + d_{\text{self}} Q_{\text{in}} + (n-2) d_{\text{in}} Q_{\text{in}} + (N-n) d_{\text{out}} Q_{\text{out}} \right). \tag{5b}
\end{aligned}$$

And for $j \notin G_i$,

$$\begin{aligned}
Q_{\text{out}} &= \frac{1-\mu}{2} \left((d_{\text{out}} + d_{\text{self}} Q_{\text{out}}) + (n-1) (d_{\text{out}} Q_{\text{in}} + d_{\text{in}} Q_{\text{out}}) \right. \\
&\quad \left. + (d_{\text{self}} Q_{\text{out}} + d_{\text{out}}) + (n-1) (d_{\text{in}} Q_{\text{out}} + d_{\text{out}} Q_{\text{in}}) \right. \\
&\quad \left. + (N-2n) (d_{\text{out}} Q_{\text{out}} + d_{\text{out}} Q_{\text{out}}) \right) \\
&= (1-\mu) \left(d_{\text{out}} + d_{\text{self}} Q_{\text{out}} + (n-1) (d_{\text{out}} Q_{\text{in}} + d_{\text{in}} Q_{\text{out}}) + (N-2n) d_{\text{out}} Q_{\text{out}} \right) \tag{5c}
\end{aligned}$$

Wright-Fisher For $j \neq i$,

$$Q_{ij} = (1-\mu)^2 \sum_{k,l=1}^N d_{ki} d_{lj} Q_{kl}. \tag{6a}$$

When $j \neq i$, $j \in G_i$,

$$\begin{aligned}
Q_{\text{in}} &= (1 - \mu)^2 \left[\left(d_{\text{self}} d_{\text{in}} + d_{\text{in}} d_{\text{self}} + (n - 2) d_{\text{in}} d_{\text{in}} + (N - n) d_{\text{out}} d_{\text{out}} \right) \right. \\
&\quad + \left(d_{\text{self}} d_{\text{self}} + (n - 2) d_{\text{self}} d_{\text{in}} \right. \\
&\quad \quad + (n - 1) d_{\text{in}} d_{\text{in}} + (n - 2) d_{\text{in}} d_{\text{self}} \\
&\quad \quad \left. + (n - 2)(n - 2) d_{\text{in}} d_{\text{in}} + (N - n)(n - 1) d_{\text{out}} d_{\text{out}} \right) Q_{\text{in}} \\
&\quad + \left((N - n) d_{\text{self}} d_{\text{out}} + (N - n)(n - 1) d_{\text{in}} d_{\text{out}} \right. \\
&\quad \quad + (N - n) d_{\text{out}} d_{\text{self}} + (N - n)(n - 1) d_{\text{out}} d_{\text{in}} \\
&\quad \quad \left. + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big] \\
&= (1 - \mu)^2 \left[\left(2 d_{\text{in}} d_{\text{self}} + (n - 2) d_{\text{in}}^2 + (N - n) d_{\text{out}}^2 \right) \right. \\
&\quad + \left(d_{\text{self}}^2 + 2(n - 2) d_{\text{self}} d_{\text{in}} + (n^2 - 3n + 3) d_{\text{in}}^2 + (N - n)(n - 1) d_{\text{out}}^2 \right) Q_{\text{in}} \\
&\quad + \left(2(N - n) d_{\text{self}} d_{\text{out}} + 2(N - n)(n - 1) d_{\text{in}} d_{\text{out}} \right. \\
&\quad \quad \left. + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big] \tag{6b}
\end{aligned}$$

And when $j \notin G_i$, we have

$$\begin{aligned}
Q_{\text{out}} &= (1 - \mu)^2 \left[\left(2 d_{\text{self}} d_{\text{out}} + 2(n - 1) d_{\text{in}} d_{\text{out}} + (N - 2n) d_{\text{out}}^2 \right) \right. \\
&\quad + \left(2(n - 1) d_{\text{self}} d_{\text{out}} + 2(n - 1)^2 d_{\text{in}} d_{\text{out}} + (N - 2n)(n - 1) d_{\text{out}}^2 \right) Q_{\text{in}} \\
&\quad + \left(d_{\text{self}} d_{\text{self}} + (n - 1) d_{\text{self}} d_{\text{in}} + (N - 2n) d_{\text{self}} d_{\text{out}} \right. \\
&\quad \quad + (n - 1) d_{\text{in}} d_{\text{self}} + (n - 1)^2 d_{\text{in}}^2 + (n - 1)(N - 2n) d_{\text{in}} d_{\text{out}} \\
&\quad \quad \left. + (N - n) d_{\text{out}} d_{\text{self}} + (N - n)(n - 1) d_{\text{out}} d_{\text{in}} + (N - n)(N - 2n) d_{\text{out}} d_{\text{out}} \right) Q_{\text{out}} \Big]. \tag{6c}
\end{aligned}$$

PAS FINI

Appendix

All combinations for i, j, k, l . Notation: (i, j) means that i and j are in the same deme, but are different; G_i refers to the deme containing site i .

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
1	$j = i$	$k = i$	$l = i$	$(i = j = k = l)$	1	d_{self}	d_{self}	e_{self}	1
2	$j = i$	$k = i$	$l \neq i; l \in G_i$	$(i = j = k, l)$	$n - 1$	d_{self}	d_{in}	e_{in}	1
3	$j = i$	$k = i$	$l \notin G_i$	$(i = j = k), (l)$	$N - n$	d_{self}	d_{out}	e_{out}	1
4	$j = i$	$k \neq i; k \in G_i$	$l = i$	$(i = j = l, k)$	$n - 1$	d_{self}	d_{self}	e_{in}	Q_{in}
5	$j = i$	$k \neq i; k \in G_i$	$l = k$	$(i = j, k = l)$	$n - 1$	d_{self}	d_{in}	e_{self}	Q_{in}
6	$j = i$	$k \neq i; k \in G_i$	$l \neq i, k; l \in G_i$	$(i = j, k, l)$	$(n - 1)(n - 2)$	d_{self}	d_{in}	e_{in}	Q_{in}
7	$j = i$	$k \neq i; k \in G_i$	$l \notin G_i$	$(i = j, k), (l)$	$(n - 1)(N - n)$	d_{self}	d_{out}	e_{out}	Q_{in}
8	$j = i$	$k \notin G_i$	$l = i = j$	$(i = j = l), (k)$	$(N - n)$	d_{self}	d_{self}	e_{out}	Q_{out}
9	$j = i$	$k \notin G_i$	$l \neq i, l \in G_i$	$(i = j, l), (k)$	$(N - n)(n - 1)$	d_{self}	d_{in}	e_{out}	Q_{out}
10	$j = i$	$k \notin G_i$	$l = k$	$(i = j), (k = l)$	$(N - n)$	d_{self}	d_{out}	e_{self}	Q_{out}
11	$j = i$	$k \notin G_i$	$l \neq k; l \in G_k$	$(i = j), (k, l)$	$(N - n)(n - 1)$	d_{self}	d_{out}	e_{in}	Q_{out}
12	$j = i$	$k \notin G_i$	$l \notin G_i, G_k$	$(i = j), (k), (l)$	$(N - n)(N - 2n)$	d_{self}	d_{out}	e_{out}	Q_{out}
13	$j \neq i, j \in G_i$	$k = i$	$l = i$	$(i = k = l, j)$	$(n - 1)$	d_{in}	d_{self}	e_{self}	Q_{in}
14	$j \neq i, j \in G_i$	$k = i$	$l = j$	$(i = k, j = l)$	$(n - 1)$	d_{in}	d_{in}	e_{in}	Q_{in}
15	$j \neq i, j \in G_i$	$k = i$	$l \neq i, j; l \in G_i$	$(i = k, j, l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	Q_{in}
16	$j \neq i, j \in G_i$	$k = i$	$l \notin G_i$	$(i = k, j), (l)$	$(n - 1)(N - n)$	d_{in}	d_{out}	e_{out}	Q_{in}
17	$j \neq i, j \in G_i$	$k = j$	$l = i$	$(i = l, j = k)$	$(n - 1)$	d_{in}	d_{self}	e_{in}	1
18	$j \neq i, j \in G_i$	$k = j$	$l = j$	$(i, j = k = l)$	$(n - 1)$	d_{in}	d_{in}	e_{self}	1
19	$j \neq i, j \in G_i$	$k = j$	$l \neq i, j; l \in G_i$	$(i, j = k, l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	1
20	$j \neq i, j \in G_i$	$k = j$	$l \notin G_i$	$(i, j = k), (l)$	$(n - 1)(N - n)$	d_{in}	d_{out}	e_{out}	1
21	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = i$	$(i = l, j, k)$	$(n - 1)(n - 2)$	d_{in}	d_{self}	e_{in}	Q_{in}
22	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = j$	$(i, j = l, k)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{in}	Q_{in}
23	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l = k$	$(i, j, k = l)$	$(n - 1)(n - 2)$	d_{in}	d_{in}	e_{self}	Q_{in}
24	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l \neq i, j, k; l \in G_i$	(i, j, k, l)	$(n - 1)(n - 2)(n - 3)$	d_{in}	d_{in}	e_{in}	Q_{in}
25	$j \neq i, j \in G_i$	$k \neq i, j; k \in G_i$	$l \notin G_i$	$(i, j, k), (l)$	$(n - 1)(n - 2)(N - n)$	d_{in}	d_{out}	e_{out}	Q_{in}

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
26	$j \neq i; j \in G_i$	$k \notin G_i$	$l = i$	$(i = l, j), (k)$	$(n-1)(N-n)$	d_{in}	d_{self}	e_{out}	Q_{out}
27	$j \neq i; j \in G_i$	$k \notin G_i$	$l = j$	$(i, j = l), (k)$	$(n-1)(N-n)$	d_{in}	d_{in}	e_{out}	Q_{out}
28	$j \neq i; j \in G_i$	$k \notin G_i$	$l \neq i, j; l \in G_i$	$(i, j, l), (k)$	$(n-1)(N-n)(n-2)$	d_{in}	d_{in}	e_{out}	Q_{out}
29	$j \neq i; j \in G_i$	$k \notin G_i$	$l = k$	$(i, j), (k = l)$	$(n-1)(N-n)$	d_{in}	d_{out}	e_{self}	Q_{out}
30	$j \neq i; j \in G_i$	$k \notin G_i$	$l \neq k; l \in G_k$	$(i, j), (k, l)$	$(n-1)(N-n)(n-1)$	d_{in}	d_{out}	e_{in}	Q_{out}
31	$j \neq i; j \in G_i$	$k \notin G_i$	$l \notin G_i, G_k$	$(i, j), (k), (l)$	$(n-1)(N-n)(N-2n)$	d_{in}	d_{out}	e_{out}	Q_{out}
32	$j \notin G_i$	$k = i$	$l = i$	$(i = k = l), (j)$	$(N-n)$	d_{out}	d_{self}	e_{self}	Q_{out}
33	$j \notin G_i$	$k = i$	$l \neq i; l \in G_i$	$(i = k, l), (j)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{in}	Q_{out}
34	$j \notin G_i$	$k = i$	$l = j$	$(i = k), (j = l)$	$(N-n)$	d_{out}	d_{out}	e_{out}	Q_{out}
35	$j \notin G_i$	$k = i$	$l \neq j; l \in G_j$	$(i = k), (j, l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
36	$j \notin G_i$	$k = i$	$l \notin G_i, G_j$	$(i = k), (j), (l)$	$(N-n)(N-2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
37	$j \notin G_i$	$k \neq i; k \in G_i$	$l = i$	$(i = l, k), (j)$	$(N-n)(n-1)$	d_{out}	d_{self}	e_{in}	Q_{out}
38	$j \notin G_i$	$k \neq i; k \in G_i$	$l = k$	$(i, k = l), (j)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{self}	Q_{out}
39	$j \notin G_i$	$k \neq i; k \in G_i$	$l \neq i, k; l \in G_i$	$(i, k, l), (j)$	$(N-n)(n-1)(n-2)$	d_{out}	d_{in}	e_{in}	Q_{out}
40	$j \notin G_i$	$k \neq i; k \in G_i$	$l = j$	$(i, k), (j = l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
41	$j \notin G_i$	$k \neq i; k \in G_i$	$l \neq j; l \in G_j$	$(i, k), (j, l)$	$(N-n)(n-1)(n-1)$	d_{out}	d_{out}	e_{out}	Q_{out}
42	$j \notin G_i$	$k \neq i; k \in G_i$	$l \notin G_i, G_j$	$(i, k), (j), (l)$	$(N-n)(n-1)(N-2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
43	$j \notin G_i$	$k = j$	$l = i$	$(i = l), (j = k)$	$(N-n)$	d_{out}	d_{self}	e_{out}	1
44	$j \notin G_i$	$k = j$	$l \neq i; l \in G_i$	$(i, l), (j = k)$	$(N-n)(n-1)$	d_{out}	d_{in}	e_{out}	1
45	$j \notin G_i$	$k = j$	$l = j$	$(i), (j = k = l)$	$(N-n)$	d_{out}	d_{out}	e_{self}	1
46	$j \notin G_i$	$k = j$	$l \neq j; l \in G_j$	$(i), (j = k, l)$	$(N-n)(n-1)$	d_{out}	d_{out}	e_{in}	1
47	$j \notin G_i$	$k = j$	$l \notin G_i, G_j$	$(i), (j = k), (l)$	$(N-n)(N-2n)$	d_{out}	d_{out}	e_{out}	1

	j	k	l	Notation	Count	d_{ji}	d_{li}	e_{kl}	Q_{jk}
48	$j \notin G_i$	$k \neq j; k \in G_j$	$l = i$	$(i = l), (j, k)$	$(N - n)(n - 1)$	d_{out}	d_{self}	e_{out}	Q_{in}
49	$j \notin G_i$	$k \neq j; k \in G_j$	$l \neq i; l \in G_i$	$(i, l), (j, k)$	$(N - n)(n - 1)(n - 1)$	d_{out}	d_{in}	e_{out}	Q_{in}
50	$j \notin G_i$	$k \neq j; k \in G_j$	$l = j$	$(i), (j = l, k)$	$(N - n)(n - 1)$	d_{out}	d_{out}	e_{in}	Q_{in}
51	$j \notin G_i$	$k \neq j; k \in G_j$	$l = k$	$(i), (j, k = l)$	$(N - n)(n - 1)$	d_{out}	d_{out}	e_{self}	Q_{in}
52	$j \notin G_i$	$k \neq j; k \in G_j$	$l \neq j, k; l \in G_j$	$(i), (j, k, l)$	$(N - n)(n - 1)(n - 2)$	d_{out}	d_{out}	e_{in}	Q_{in}
53	$j \notin G_i$	$k \neq j; k \in G_j$	$l \notin G_i, G_j$	$(i), (j, k), (l)$	$(N - n)(n - 1)(N - 2n)$	d_{out}	d_{out}	e_{out}	Q_{in}
54	$j \notin G_i$	$k \notin G_i, G_j$	$l = i$	$(i = l), (j), (k)$	$(N - n)(N - 2n)$	d_{out}	d_{self}	e_{out}	Q_{out}
55	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq i; l \in G_i$	$(i, l), (j), (k)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{in}	e_{out}	Q_{out}
56	$j \notin G_i$	$k \notin G_i, G_j$	$l = j$	$(i), (j = l), (k)$	$(N - n)(N - 2n)$	d_{out}	d_{out}	e_{out}	Q_{out}
57	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq j; l \in G_j$	$(i), (j, l), (k)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{out}	e_{out}	Q_{out}
58	$j \notin G_i$	$k \notin G_i, G_j$	$l = k$	$(i), (j), (k = l)$	$(N - n)(N - 2n)$	d_{out}	d_{out}	e_{self}	Q_{out}
59	$j \notin G_i$	$k \notin G_i, G_j$	$l \neq k; l \in G_k$	$(i), (j), (k, l)$	$(N - n)(N - 2n)(n - 1)$	d_{out}	d_{out}	e_{in}	Q_{out}
60	$j \notin G_i$	$k \notin G_i, G_j$	$l \notin G_i, G_j, G_k$	$(i), (j), (k), (l)$	$(N - n)(N - 2n)(N - 3n)$	d_{out}	d_{out}	e_{out}	Q_{out}

A Island model

With self replacement

$$d_{\text{self}} = d_{\text{in}} = \frac{1-m}{n}, \quad (7a)$$

$$d_{\text{out}} = \frac{m}{N-n}. \quad (7b)$$

Without self-replacement

$$d_{\text{self}} = 0, \quad (8a)$$

$$d_{\text{in}} = \frac{1-m}{n-1}, \quad (8b)$$

$$d_{\text{out}} = \frac{m}{N-n}. \quad (8c)$$

B IDB

B.1 Moran

Using the formulas for a 2D graph in REF Debarre 2017,

$$\tilde{\mathcal{D}}_{q_1} = \sum_{q_2=0}^{N_1-1} \sum_{l_2=0}^{N_2-1} \tilde{d}_{l_1} \exp\left(-i \frac{2\pi q_1 l_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) \quad (9a)$$

$$\tilde{\mathcal{Q}}_{r_1} = \frac{1}{N} \sum_{q_1=0}^{N_1-1} \sum_{q_2=0}^{N_2-1} \frac{\mu \lambda'_M}{1 - (1-\mu) \tilde{\mathcal{D}}_{q_1}} \exp\left(i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(i \frac{2\pi q_2 r_2}{N_2}\right) \quad (9b)$$

We have

$$\begin{aligned} \tilde{\mathcal{D}}_{q_1} &= d_{\text{self}} + \sum_{l_2=1}^{N_2-1} d_{\text{in}} \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) + \sum_{l_1=1}^{N_1-1} \sum_{l_2=0}^{N_2-1} d_{\text{out}} \exp\left(-i \frac{2\pi q_1 l_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 l_2}{N_2}\right) \\ &= d_{\text{self}} + (\delta_{q_2}(N_2-1) + (1-\delta_{q_2})(-1)) d_{\text{in}} + (\delta_{q_1}(N_1-1) + (1-\delta_{q_1})(-1)) (\delta_{q_2} N_2) d_{\text{out}} \\ &= d_{\text{self}} + (\delta_{q_2} N_2 - 1) d_{\text{in}} + (\delta_{q_1} N_1 - 1) \delta_{q_2} N_2 d_{\text{out}}. \end{aligned} \quad (10a)$$

Whether there is self-replacement or not, we have $N_1 = D$ and $N_2 = n$, and

$$\tilde{\mathcal{D}}_0 = 1, \quad (11a)$$

$$\tilde{\mathcal{D}}_{q_1} = 1 - m - \frac{m}{d-1} \quad (q_1 \not\equiv 0 \pmod{N_1}), \quad (11b)$$

$$\tilde{\mathcal{D}}_{q_1} = d_{\text{self}} - d_{\text{in}} \quad (q_2 \not\equiv 0 \pmod{N_2}). \quad (11c)$$

So for \tilde{Q} ,

$$\begin{aligned}
\tilde{Q}_{r_1, r_2} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{1 - (1 - \mu) \tilde{D}_0} + \sum_{q_2=1}^{N_2-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_2}} \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) + \sum_{q_1=1}^{N_1-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_1}} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \right. \\
&\quad \left. + \sum_{q_1=1}^{N_1-1} \sum_{q_2=1}^{N_2-1} \frac{1}{1 - (1 - \mu) \tilde{D}_{q_1, q_2}} \exp\left(-i \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-i \frac{2\pi q_2 r_2}{N_2}\right) \right] \\
&= \frac{\mu \lambda'_M}{N} \left[\frac{1}{1 - (1 - \mu)} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (\delta_{r_2} N_2 - 1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (\delta_{r_1} N_1 - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (\delta_{r_1} N_1 - 1)(\delta_{r_2} N_2 - 1) \right]. \tag{12a}
\end{aligned}$$

In particular,

$$\begin{aligned}
\tilde{Q}_0 &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (n - 1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (D - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (D - 1)(n - 1) \right] \\
&= 1. \tag{12b}
\end{aligned}$$

We find λ'_M using the above equation. When $r_1 = 0$, the two individuals are in the same deme. They are different when $r_2 \neq 0$:

$$\begin{aligned}
Q_{\text{in}} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (-1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (D - 1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (D - 1)(-1) \right]. \tag{12c}
\end{aligned}$$

And when $r_1 \neq 0$, the two individuals are in different demes:

$$\begin{aligned}
Q_{\text{out}} &= \frac{\mu \lambda'_M}{N} \left[\frac{1}{\mu} + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} (-1) + \frac{1}{1 - (1 - \mu)(1 - m - \frac{m}{d-1})} (-1) \right. \\
&\quad \left. + \frac{1}{1 - (1 - \mu)(d_{\text{self}} - d_{\text{in}})} \right]. \tag{12d}
\end{aligned}$$

B.2 Wright-Fisher

$$\begin{aligned}
\tilde{Q}_{r_1 r_2} &= \frac{1}{N} \sum_{q_1=0}^{N_1-1} \sum_{q_2=0}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_{q_1}^2)_{q_2}^2} \exp\left(-\iota \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-\iota \frac{2\pi q_2 r_2}{N_2}\right) \\
&= \frac{1}{N} \left[\frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0^2)} + \sum_{q_2=1}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0^2)_{q_2}^2} \exp\left(-\iota \frac{2\pi q_2 r_2}{N_2}\right) \right. \\
&\quad + \sum_{q_1=1}^{N_1-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_0^2)_{q_1}^2} \exp\left(-\iota \frac{2\pi q_1 r_1}{N_1}\right) \\
&\quad \left. + \sum_{q_1=1}^{N_1-1} \sum_{q_2=1}^{N_2-1} \frac{\mu \lambda'_{WF}}{1 - (1-\mu)^2 (\tilde{\mathcal{D}}_{q_1}^2)_{q_2}^2} \exp\left(-\iota \frac{2\pi q_1 r_1}{N_1}\right) \exp\left(-\iota \frac{2\pi q_2 r_2}{N_2}\right) \right] \quad (13)
\end{aligned}$$

$$\begin{aligned}
&= \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_2} N_2 - 1) \right. \\
&\quad + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (\delta_{q_1} N_1 - 1) \\
&\quad \left. + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_1} N_1 - 1) (\delta_{q_2} N_2 - 1) \right] \\
&= \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (\delta_{q_2} N_2 - 1) \delta_{q_1} N_1 \right. \\
&\quad \left. + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (\delta_{q_1} N_1 - 1) \right]. \quad (14)
\end{aligned}$$

To find λ'_{WF} , we solve

$$1 = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} + \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} (N_2 - 1) N_1 + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (N_1 - 1) \right]. \quad (15a)$$

Then,

$$Q_{\text{in}} = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} - \frac{1}{1 - (1-\mu)^2 (d_{\text{self}} - d_{\text{in}})^2} N_1 + \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} (N_1 - 1) \right]. \quad (15b)$$

and

$$Q_{\text{out}} = \frac{\mu \lambda'_{WF}}{N} \left[\frac{1}{1 - (1-\mu)^2} - \frac{1}{1 - (1-\mu)^2 (1 - m - \frac{m}{d-1})^2} \right]. \quad (15c)$$