

# 4ème journée

Social evolution in structured populations

Florence Débarre




@flodebarre


CNRS



Centre de Interdisciplinaire de Recherche en Biologie, Paris


## Le 26 octobre 2017

# «Biologie & Mathématiques sur la Montagne»











Assortment / Relatedness


## Spatial structure, population viscosity and altruism




## Spatial structure, population viscosity and altruism




## Spatial structure, population viscosity and altruism



## Spatial structure, population viscosity and altruism



# Spatial structure, population viscosity and altruism



*Evolutionary Ecology*, 1992, 6, 352–356

## Altruism in viscous populations – an inclusive fitness model

P.D. TAYLOR

*Department of Mathematics and Statistics, Queen's University, Kingston Ont. K7L 3N6, Canada*


### Summary

A viscous population (Hamilton, 1964) is one in which the movement of organisms from their place of birth is relatively slow. This viscosity has two important effects: one is that local interactions tend to be among relatives, and the other is that competition for resources tends to be among relatives. The first effect tends to promote and the second to oppose the evolution of altruistic behaviour. In a simulation model of Wilson *et al.* (1992) these two factors appear to exactly balance one another, thus opposing the evolution of local altruistic behaviour. Here I show, with an inclusive fitness model, that the same result holds in a patch-structured population.

**Keywords:** altruism; inclusive fitness; competition; viscosity


## Subdivided population – Island model

$N_d$  demes




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )




## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )



## Subdivided population – Island model

$N_d$  demes of  $n$  individuals each (total population size  $N = n N_d$ )



## The choice of life-cycle matters

Constant population size ( $N$ ), so between two time steps,  $\#\text{▀} = \#\text{👶}$ .

## The choice of life-cycle matters

Constant population size ( $N$ ), so between two time steps,  $\#\text{▀} = \#\text{▀}$ .

Wright-Fisher



Moran Birth-Death



Moran Death-Birth



## The choice of life-cycle matters

Constant population size ( $N$ ), so between two time steps,  $\#\text{▀} = \#\text{🚼}$ .


Wright-Fisher



Moran Birth-Death





Moran Death-Birth





In homogeneously structured populations,  
with effects of social interactions on **fecundity**.

## A common feature of models



## A common feature of models



## A common feature of models



## A common feature of models




What is the effect of population viscosity  
on the evolution of altruism when parent-  
offspring strategy transmission is **imperfect**?

## Fidelity of parent-offspring transmission

### Causes of imperfect strategy transmission


- ▶ Mutation



# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission


- ▶ Mutation
- ▶ Partial heritability



# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission


- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)



# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission

- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)



In the model

Parent




# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission

- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)




In the model




# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission

- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)




In the model




# Fidelity of parent-offspring transmission

## Causes of imperfect strategy transmission

- ▶ Mutation
- ▶ Partial heritability
- ▶ Cultural transmission (vertical)



In the model




## Notation

$$X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{🐍 at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{🐦 at time } t (1 \leq i \leq N) \end{cases}$$

## Notation

$$X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{duck} \text{ at time } t (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t (1 \leq i \leq N) \end{cases}$$



$$\mathbb{E}[Y_i] = (1 - \mu) X_i + \mu \nu.$$

Expected trait of the  
offspring of individual  $i$

## Notation

$$X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{green bird} \text{ at time } t \ (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t \ (1 \leq i \leq N) \end{cases}$$

Proportion of altruists in the population:

$$\bar{X} = \frac{1}{N} \sum_{i=1}^N X_i.$$

## Notation

$$X_i(t) = \begin{cases} 1 & \text{if site } i \text{ occupied by } \text{green bird} \text{ at time } t \ (1 \leq i \leq N) \\ 0 & \text{if site } i \text{ occupied by } \text{blue bird} \text{ at time } t \ (1 \leq i \leq N) \end{cases}$$

Proportion of altruists in the population:

$$\bar{X} = \frac{1}{N} \sum_{i=1}^N X_i.$$

We want to compute  $\mathbb{E}[\bar{X}]$ ,  
the expected proportion of altruists in the population.

### Phenotype

$$\phi_i = \delta X_i,$$

and we assume that  $\delta \ll 1$ . (Selection is weak.)

## Social interactions

### Phenotype

$$\phi_i = \delta X_i,$$

and we assume that  $\delta \ll 1$ . (Selection is weak.)

### Social interactions affect fecundity

At the first order in  $\delta$ ,

$$f_i = 1 + \delta \left( b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

## Social interactions

### Phenotype

$$\phi_i = \delta X_i,$$

and we assume that  $\delta \ll 1$ . (Selection is weak.)

### Social interactions affect fecundity

At the first order in  $\delta$ ,

$$f_i = 1 + \delta \left( b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$


## Social interactions

### Phenotype

$$\phi_i = \delta X_i,$$

and we assume that  $\delta \ll 1$ . (Selection is weak.)

### Social interactions affect fecundity

At the first order in  $\delta$ ,

$$f_i = 1 + \delta \left( b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

Proportion of altruists  
among the other deme-mates

## Phenotype

$$\phi_i = \delta X_i,$$

and we assume that  $\delta \ll 1$ . (Selection is weak.)

## Social interactions affect fecundity

At the first order in  $\delta$ ,

$$f_i = 1 + \delta \left( b \sum_{j \in \mathcal{D}_i \setminus i} \frac{x_j}{n-1} - c X_i \right).$$

Proportion of altruists among the other deme-mates

The cost is only paid by altruists

## Calculations

### Notation

$B_i = B_i(\mathbf{X}, \delta)$ : expected # of offspring of individual  $i$ ;

$D_i = D_i(\mathbf{X}, \delta)$ : probability that  $i$  dies.

## Calculations

### Notation

$B_i = B_i(\mathbf{X}, \delta)$ : expected # of offspring of individual  $i$ ;

$D_i = D_i(\mathbf{X}, \delta)$ : probability that  $i$  dies.

- ▶ Expected proportion of altruists at  $t + 1$  in the proportion of altruists, conditional on the state of the population at time  $t$ :

$$\mathbb{E}[\bar{X}(t+1)|\mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N [B_i(1 - \mu)X_i + (1 - D_i)X_i + B_i\mu\nu]$$

## Calculations

### Notation

$B_i = B_i(\mathbf{X}, \delta)$ : expected # of offspring of individual  $i$ ;

$D_i = D_i(\mathbf{X}, \delta)$ : probability that  $i$  dies.

- ▶ Expected proportion of altruists at  $t + 1$  in the proportion of altruists, conditional on the state of the population at time  $t$ :

$$\mathbb{E}[\bar{X}(t+1)|\mathbf{X}(t)] = \frac{1}{N} \sum_{i=1}^N [B_i(1 - \mu)X_i + (1 - D_i)X_i + B_i\mu\nu]$$

- ▶ Take expectation and let  $t \rightarrow \infty$ ; consider stationary distribution  $\xi$

$$0 = \frac{1}{N} \sum_{x \in \Omega} \left[ \sum_{i=1}^N \underbrace{B_i(1 - \mu) - D_i}_{W_i} X_i + \sum_{i=1}^N B_i\mu\nu \right] \xi(\mathbf{X}, \delta, \mu)$$

## Calculations (2)

- Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

## Calculations (2)

- Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[ \frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

## Calculations (2)

- Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[ \frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

## Calculations (2)

- Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[ \frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

## Calculations (2)

- ▶ Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[ \frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

- ▶ Using partial derivatives: phenotypes

$$\frac{\partial W_i}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \frac{\partial \phi_k}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} X_k.$$

## Calculations (2)

- Selection is weak ( $\delta \ll 1$ ) and reproductive values are all equal:

$$0 = \frac{\delta}{N} \sum_{i=1}^N \left[ \sum_{X \in \Omega} \frac{\partial W_i}{\partial \delta} X_i \xi(\mathbf{X}, 0, \mu) - \sum_{X \in \Omega} \mu B^* X_i \frac{\partial \xi}{\partial \delta} \right] + O(\delta^2),$$

which we rewrite as

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \mathbb{E}_0 \left[ \frac{\partial W_i}{\partial \delta} X_i \right] + O(\delta^2).$$

- Using partial derivatives: phenotypes

$$\frac{\partial W_i}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \frac{\partial \phi_k}{\partial \delta} = \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} X_k.$$

- We obtain

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \sum_{k=1}^N \frac{\partial W_i}{\partial \phi_k} \underbrace{\mathbb{E}_0 [X_i X_k]}_{P_{ik}} + O(\delta^2).$$

## Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n - 1) \frac{\partial W_i}{\partial \phi_{in}} + (N - n) \frac{\partial W_i}{\partial \phi_{out}} = 0,$$

## Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$

- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left( \underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$

## Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$

- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left( \underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$

- ▶ Then further decompose with partial derivatives:

$$\frac{\partial W_i}{\partial \phi_k} = \sum_{\ell=1}^N \frac{\partial W_i}{\partial f_\ell} \frac{\partial f_\ell}{\partial \phi_k}$$

## Calculations (3)

- ▶ In a subdivided population,

$$\frac{\partial W_i}{\partial \phi_i} + (n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}} + (N-n) \frac{\partial W_i}{\partial \phi_{\text{out}}} = 0,$$


- ▶ So

$$\delta \mu B^* \frac{\partial \mathbb{E}[\bar{X}]}{\partial \delta} = \frac{\delta}{N} \sum_{i=1}^N \left( \underbrace{\frac{\partial W_i}{\partial \phi_i}}_{-c} + \underbrace{(n-1) \frac{\partial W_i}{\partial \phi_{\text{in}}}}_{\mathcal{B}} \underbrace{\frac{P_{\text{in}} - P_{\text{out}}}{P_{ii} - P_{\text{out}}}}_R \right) (P_{ii} - P_{\text{out}}) + O(\delta^2).$$


- ▶ Then further decompose with partial derivatives:

$$\frac{\partial W_i}{\partial \phi_k} = \sum_{\ell=1}^N \frac{\partial W_i}{\partial f_\ell} \frac{\partial f_\ell}{\partial \phi_k} \quad \text{and} \quad \frac{\partial f_\ell}{\partial \phi_\ell} = -\text{c}; \quad \frac{\partial f_\ell}{\partial \phi_{\text{in}}} = \frac{\text{b}}{n-1}; \quad \frac{\partial f_\ell}{\partial \phi_{\text{out}}} = 0.$$


## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state




## Genealogy, Identity by descent and Identity in state



## Genealogy, Identity by descent and Identity in state



## Genealogy, Identity by descent and Identity in state



## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

$$P_{ij}$$

Expected state  
of the  $i,j$  pair

= Probability that the two  
individuals are altruists

## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

$$P_{ij} = Q_{ij} \nu$$

Expected state  
of the  $i, j$  pair  
= Probability that the two  
individuals are altruists

Probability that the individuals at  
sites  $i$  and  $j$  are identical by descent  
(no mutation since  
their common ancestor)

## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

$$P_{ij} = Q_{ij} \nu$$

Expected state of the  $i, j$  pair  
= Probability that the two individuals are altruists

Probability that a mutant is an altruist  
= Probability that a given site is occupied by an altruist

Probability that the individuals at sites  $i$  and  $j$  are identical by descent  
(no mutation since their common ancestor)

## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

$$P_{ij} = Q_{ij} \nu + (1 - Q_{ij}) \nu^2$$

Expected state of the  $i, j$  pair  
= Probability that the two individuals are altruists

Probability that both sites are occupied by an altruist

Probability that the individuals at sites  $i$  and  $j$  are not identical by descent

```
graph TD; Pij["Pij = Qij ν + (1 - Qij) ν2"]; Pij --> Qij_nu["Qij ν"]; Qij_nu --> ExpectedState["Expected state of the i, j pair  
= Probability that the two individuals are altruists"]; Pij --> OneMinusQij_nu2["(1 - Qij) ν2"]; OneMinusQij_nu2 --> BothOccupied["Probability that both sites are occupied by an altruist"]; Pij --> BothNotIdentical["Probability that the individuals at sites i and j are not identical by descent"];
```

## Expected state of pairs of sites and identity by descent

At neutrality (i.e., in the absence of selection,  $\delta = 0$ ),

$$P_{ij} = Q_{ij} \nu + (1 - Q_{ij}) \nu^2$$

Expected state  
of the  $i, j$  pair

= Probability that the two  
individuals are altruists

$Q_{\text{in}}$ ,  $Q_{\text{out}}$

## Expected frequency of altruists in the population

$$\begin{aligned}\mathbb{E}[\bar{X}] = & \nu + \delta \nu(1-\nu) \frac{1-\mu}{\mu} (1-Q_{\text{out}}) \times \\ & \left( -c - (b-c) \left( \frac{(1-m)^2}{n} + \frac{m^2}{n(N_d-1)} \right) \right. \\ & \left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1-Q_{\text{out}}} \left[ b - (b-c)(n-1) \left( \frac{(1-m)^2}{n} + \frac{m^2}{n(N_d-1)} \right) \right] \right)\end{aligned}$$

## Expected frequency of altruists in the population

Mutation-drift  
equilibrium

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

## Expected frequency of altruists in the population

Mutation-drift  
equilibrium      Selection  
strength

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

## Expected frequency of altruists in the population

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$

$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right.$$

$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$

Mutation-drift equilibrium   Selection strength   Population variance  
 Variance in the state of one site

## Expected frequency of altruists in the population

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$\left. + \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \right)$$

Mutation-drift equilibrium   Selection strength   Population variance  
Variance in the state of one site

## Expected frequency of altruists in the population

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \Big)$$

$\mathcal{B}$

Mutation-drift equilibrium   Selection strength   Population variance  
Variance in the state of one site

## Expected frequency of altruists in the population

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \left. \right)$$

*R*      *B*

Mutation-drift equilibrium    Selection strength    Population variance  
Variance in the state of one site

## Expected frequency of altruists in the population

Mutation-drift equilibrium      Selection strength      Population variance  
Variance in the state of one site

$$\mathbb{E}[\bar{X}] = \nu + \delta \nu(1 - \nu) \frac{1 - \mu}{\mu} (1 - Q_{\text{out}}) \times$$
$$\left( -c - (b - c) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) - C \right.$$
$$+ \frac{Q_{\text{in}} - Q_{\text{out}}}{1 - Q_{\text{out}}} \left[ b - (b - c)(n - 1) \left( \frac{(1 - m)^2}{n} + \frac{m^2}{n(N_d - 1)} \right) \right] \Big)$$

*R*                            *B*