

Bertrand's Theorem: Closed, Perturbatively Stable Orbits

Daniel Topa dantopa@gmail.com

University of New Mexico
Department of Mathematics and Statistics

September 15, 2024





Overview

- Bertrand's Theorem and Proofs
- 2 Key Concepts
- Bertrand's Theorem
- Backup Slides



Proof Types

- Naïve expansion
- Oedicated Web Sites
- **3** Federation of American Scientists

The Theorem of Bertrand Rich Literature

Bertrand's Paper En français

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 4875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE.

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.



The Theorem of Bertrand Rich Literature

Bertrand's Paper in English

An English Translation of Bertrand's Theorem



F. C. Santos, V. Soares and A. C. Tort

Instituto de Física, Universidade Federal do Rio de Janeiro, Caixa Postal 68.528; CEP 21941-972 Rio de Janeiro, Brazil.

E-mail: tort@ufrj.br

(Received 26 November 2011 accepted 29 December 2011)

Abstract

A beautiful theorem due to J. L. F. Bertrand concerning the laws of attraction that admit bounded closed orbits for arbitrarily chosen initial conditions is translated from French into English.

Keywords: Classical Mechanics, Orbits.

Resumen

Un hermoso teorema debido a la J. L. F. Bertrand sobre las leyes de la atracción que admiten limitadas órbitas cerradas arbitrariamente escogidas para las condiciones iniciales es traducido del francés al inglés.

Palabras clave: Mecánica Clásica, Órbitas.

PACS: 45.20.D-. 01.40.G-. 01.40.Fk

ISSN 1870-9095

Santos, Soares, and Tort 2011



The Theorem of Bertrand Rich Literature

Bertrand's Paper in English

An English translation of Bertrand's theorem

F C Santos *
V Soares[†]
A C Tort [‡]
Instituto de Física

Universidade Federal do Rio de Janeiro Caixa Postal 68.528; CEP 21941-972 Rio de Janeiro, Brazil

October 30, 2018

Abstract

A beautiful theorem due to J. L. F. Bertrand concerning the laws of attraction that admit bounded closed orbits for arbitrarily chosen initial conditions is translated from French into English.

Santos, Soares, and Tort 2011





Proofs

- Théorème relatif au mouvement d'un point attiré vers un centre fixe (Bertrand 1873)
- An English translation of Bertrand's theorem (Bertrand 1873)
- A simplified proof of Bertrand's theorem (Tikochinsky 1988)
- A pedagogical relook at Bertrand's theorem (TU and Das 2019)
- Alternative proof of Bertrand's theorem using a phase space approach (Quilantán, Río-Correa, and Medina 1995)
- A proof of bertrand's theorem using the theory of isochronous potentials (Ortega and Rojas 2019)
- A partial proof of Bertrand's theorem (Musgrove 2020)

Lyapunov stability: Wikipedia PDE linearization Phase portraits Dimensional reduction

Proof Types

- Lyapunov stability
- PDE linearization
- Phase portraits
- Open Dimensional reduction





Lyapunov stability: Wikipedia PDE linearization Phase portraits Dimensional reduction

Lyapunov

Lyapunov's second method for stability [edit source]

Lyapunov, in his original 1892 work, proposed two methods for demonstrating stability. (1) The first method developed the solution in a series which was then proved convergent within limits. The second method, which is now referred to as the Lyapunov stability criterion or the Direct Method, makes use of a Lyapunov function Y(x) which has an analogy to the potential function of classical dynamics. It is introduced as follows for a system $\dot{x}=f(x)$ having a point of equilibrium at x=0. Consider a function $Y: \mathbb{R}^n \to \mathbb{R}$ such that

- ullet V(x)=0 if and only if x=0
- V(x)>0 if and only if $x\neq 0$
- $\bullet \ \dot{V}(x) = \frac{d}{dt}V(x) = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(x) = \nabla V \cdot f(x) \leq 0 \ \text{for all values of } x \neq 0 \ . \ \text{Note: for asymptotic stability, } \dot{V}(x) < 0 \ \text{for } x \neq 0 \ \text{is required.}$

Then V(y) is called a Lyapunov function and the system is stable in the sense of Lyapunov, (Note that V(0) = 0 is required; otherwise for example V(x) = 1/(1+|x|) would "prove" that $\dot{x}(t) = x$ is locally stable.) An additional condition called "properness" or "radial unboundedness" is required in order to conclude global stability. Global asymptotic stability (GAS) follows similarly.

It is easier to visualize this method of analysis by thinking of a physical system (e.g. vibrating spring and mass) and considering the energy of such a system. If the system loses energy over time and the energy is never restored then eventually the system must grind to a stop and reach some final resting state. This final state is called the attractor. However, finding a function that gives the precise energy of a physical system can be difficult, and for abstract mathematical systems, economic systems or biological systems, the concept of energy may not be applicable.

Lyapunov's realization was that stability can be proven without requiring knowledge of the true physical energy, provided a Lyapunov function can be found to satisfy the above constraints.

Lyapunov stability: Wikipedia PDE linearization Phase portraits Dimensional reduction

Papier de Bertrand en français

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 1875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE.

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.



Lyapunov stability: Wikipedia PDE linearization Phase portraits Dimensional reduction

Bertrand's Paper in French

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 1875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE,

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.



Lyapunov stability: Wikipedia PDE linearization Phase portraits Dimensional reduction

Bertrand's Paper in French

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 4875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE.

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.



Statements Survey of Proofs In the Literature Dimensional reduction

Bertrand's Theorem

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 4875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE.

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.





Statements Survey of Proofs In the Literature Dimensional reduction

Bertrand's Paper in French

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 1875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE,

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.





Statements
Survey of Proofs
In the Literature
Dimensional reduction

Lyapunov

Lyapunov's second method for stability [edit source]

Lyapunov, in his original 1892 work, proposed two methods for demonstrating stability. (I) The first method developed the solution in a series which was then proved convergent within limits. The second method, which is now referred to as the Lyapunov stability criterion or the Direct Method, makes use of a Lyapunov function V(x) which has an analogy to the potential function of classical dynamics. It is introduced as follows for a system $\dot{x} = f(x)$ having a point of equilibrium at x = 0. Consider a function $V: \mathbb{R}^n \to \mathbb{R}$ such that

- ullet V(x)=0 if and only if x=0
- V(x)>0 if and only if $x\neq 0$
- $\bullet \ \dot{V}(x) = \frac{d}{dt}V(x) = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(x) = \nabla V \cdot f(x) \leq 0 \ \text{for all values of } x \neq 0 \ . \ \text{Note: for asymptotic stability, } \dot{V}(x) < 0 \ \text{for } x \neq 0 \ \text{is required.}$

Then V(y) is called a Lyapunov function and the system is stable in the sense of Lyapunov, (Note that V(0) = 0 is required; otherwise for example V(x) = 1/(1+|x|) would "prove" that $\dot{x}(t) = x$ is locally stable.) An additional condition called "properness" or "radial unboundedness" is required in order to conclude global stability. Global asymptotic stability (GAS) follows similarly.

It is easier to visualize this method of analysis by thinking of a physical system (e.g. vibrating spring and mass) and considering the energy of such a system. If the system loses energy over time and the energy is never restored then eventually the system must grind to a stop and reach some final resting state. This final state is called the attractor. However, finding a function that gives the precise energy of a physical system can be difficult, and for abstract mathematical systems, economic systems or biological systems, the concept of energy may not be applicable.

Lyapunov's realization was that stability can be proven without requiring knowledge of the true physical energy, provided a Lyapunov function can be found to satisfy the above constraints.



Literature: Texts

- **1** Arnol'd 2013, §2.8D, p. 37
- Goldstein 1980, App. A, p. 601
- **3** José and Saletan 1998, §2.3.3, p. 88

Statements
Survey of Proofs
In the Literature
Dimensional reduction

Bertrand's Paper in French

COMPTES RENDUS

DES SÉANCES

DE L'ACADÉMIE DES SCIENCES.

SÉANCE DU LUNDI 20 OCTOBRE 4875.

PRÉSIDENCE DE M. DE QUATREFAGES.

MÉMOIRES ET COMMUNICATIONS

DES MEMBRES ET DES CORRESPONDANTS DE L'ACADÉMIE.

MÉCANIQUE ANALYTIQUE. — Théorème relatif au mouvement d'un point attiré vers un centre fixe; par M. J. Bertrand.



Conservative Force



Bibliography I

- [1] Vladimir Igorevich Arnol'd. Mathematical methods of classical mechanics. Vol. 60. Springer Science & Business Media, 2013.
- [2] Joseph Bertrand. "Théorème relatif au mouvement d'un point attiré vers un centre fixe". In: CR Acad. Sci 77.16 (July 1873), pp. 849–853. URL: https://gallica.bnf.fr/ark:/12148/bpt6k3034n.
- [3] Siu A Chin. "A truly elementary proof of Bertrand's theorem". In: American Journal of Physics 83.4 (2015), pp. 320–323.
- [4] Herbert Goldstein. Classical mechanics. 1980.





Bibliography II

- [5] Jorge V José and Eugene J Saletan. Classical dynamics: a contemporary approach. Cambridge University Press, 1998.
- [6] Vladimir Jovanović. "A note on the proof of Bertrand's theorem". In: Theoretical and Applied Mechanics 42.1 (2015), pp. 27–33.
- [7] John F Musgrove. "A Partial Proof of Bertrand's Theorem". In: (2020).
- [8] Rafael Ortega and David Rojas. "A proof of Bertrand's theorem using the theory of isochronous potentials". In: Journal of Dynamics and Differential Equations 31.4 (2019), pp. 2017–2028.





Bibliography III

- [9] José Luis Castro Quilantán, José Luis del Río-Correa, and Marco Antonio Rosales Medina. "Alternative proof of Bertrand's theorem using a phase space approach". In: Revista Mexicana de Física 42.5 (1995), pp. 867–877.
- [10] FC Santos, V Soares, and AC Tort. "An English translation of Bertrand's theorem". In: Lat. Am. J. Phys. Educ 5.4 (2011), p. 694.
- [11] Yoel Tikochinsky. "A simplified proof of Bertrand's theorem". In: American Journal of Physics 56.12 (1988), pp. 1073–1075.





Bibliography IV

[12] Jeevitha TU and Sanjit Das. "A Pedagogical Relook at Bertrand's Theorem". In: Resonance 24.11 (2019), pp. 1235–1251.



Bertrand's Theorem: Closed, Perturbatively Stable Orbits

Daniel Topa dantopa@gmail.com

University of New Mexico
Department of Mathematics and Statistics

September 15, 2024

