

Distribution of species, sequence types, plasmid backbones and carbapenemase genes

Among 70 isolates, 68 carried at least one carbapenemase gene. Sixty-three carbapenemase-positive isolates with an assigned plasmid backbone were included in the Sankey diagram. Four isolates carrying two carbapenemase-bearing plasmids were expanded into two backbone-level flows, yielding a total of 67 flows.




Figure 3. Distribution of bacterial species, multilocus sequence types (STs), plasmid backbone clusters, and carbapenemase genes among carbapenemase-producing isolates. The Sankey diagram summarizes plasmid backbone assignments based on the final curated reference backbone collection. Among the 70 isolates analyzed, 68 carried at least one carbapenemase gene. Sixty-three carbapenemase-positive isolates with an assigned plasmid backbone were included in the visualization. Four isolates carrying two carbapenemase-bearing plasmids (PAD003, ZCA001, ZCU001 and ZCZ001) were expanded into two independent backbone-level flows, resulting in 67 total flows. Clusters C01–C05 correspond to recurrent plasmid backbone groups, whereas S01 and S03 represent singleton plasmids. Carbapenemase-negative isolates and carbapenemase-positive isolates lacking an assigned plasmid backbone were excluded from the Sankey flows. Node counts correspond to backbone-level flows rather than unique isolates.