

Choose Your Own Project: Pancreatic ductal adenocarcinoma biomarkers

Daniel Flores Orozco

2021-12-16

1. Introduction

Pancreatic ductal adenocarcinoma (PDAC) is the most common pancreatic neoplasm and is considered a highly aggressive and lethal cancer (Sarantis et al., 2020). In fact, less than 20% of individuals diagnosed with PDAC survive the first year, and less than 10% survived more than five years (Debernardi et al., 2020; Sarantis et al., 2020). The poor prognosis is mainly attributed to late diagnosis and the lack of efficient treatments during an early stage. However, when the disease is detected in the early stages, the probability of surviving more than five years increases up to 70% (Debernardi et al., 2020). Therefore, the development of novel tests for the early detection of PDAC is essential to improve the patients' prognosis.

Debernardi et al. (2020) identified a group of proteins (creatinine, LYVE1, REG1A, REG1B, TFF1, and plasma CA19-9) in the urine that could potentially serve as biomarkers to detect PDAC. Debernardi et al. (2020) developed an algorithm to differentiate cancer samples from benign samples using LYVE1, REG1B, and TFF1 levels, as well as creatinine levels and age as predictors. Using logistic regression, they achieved a sensitivity and specificity of 0.963 and 0.967, respectively.

In this document, different Machine Learning algorithms, including random forest, kNN, SVM, logistic regression, and bayesian logistic regression, were used to predict if a given sample belonged to a benign or cancer group using the data provided by Debernardi et al. (2020). The ultimate objective of this project was to improve the performance reported by the original authors.

2. Methodology

2.1 Database

The pancreatic cancer biomarkers data was downloaded from the database repository Kaggle (<https://www.kaggle.com/johnjdavisiv/urinary-biomarkers-for-pancreatic-cancer>). The original data contain 14 columns: *sample_id*, *patient_cohort*, *sample_origin*, *age*, *sex*, *diagnosis*, *stage*, *benign_sample_diagnosis*, *plasma_CA19_9*, *creatinine*, *LYVE1*, *REG1B*, *TFF1*, *REG1A*. A description of each column information is presented in **Table 1**. The database contained 590 entries of the same number of patients. The patients' ages range from 26-89 years, with 299 males and 291 females. Around 33.7% of the entries came from patients diagnosed with cancer, and the rest from healthy or other benign pancreatic diagnostic.

Since the information *patient_cohort*, *sample_origin*, was not useful for the development of the different models, they were removed. Similarly, *stage*, and *benign_sample_diagnosis* were not included as the number of samples in each group was small.

Table 1: Columns details in original data set

Column name	Details
<i>sample_id</i>	Unique string identifying each subject
<i>patient_cohort</i>	Cohort 1, previously used samples; Cohort 2, newly added samples
<i>sample_origin</i>	BPTB: Barts Pancreas Tissue Bank, London, UK; ESP: Spanish National Cancer Research Centre, Madrid, Spain; LIV: Liverpool University, UK; UCL: University College London, UK
<i>age</i>	Age in years
<i>sex</i>	M = male, F = female
<i>diagnosis</i>	1 = control (no pancreatic disease), 2 = benign hepatobiliary disease (119 of which are chronic pancreatitis); 3 = Pancreatic ductal adenocarcinoma, i.e. pancreatic cancer
<i>stage</i>	For those with pancreatic cancer, what stage was it? One of IA, IB, IIA, IIIB, III, IV
<i>benign_sample_diagnosis</i>	For those with a benign, non-cancerous diagnosis, what was the diagnosis?
<i>plasma_CA19_9</i>	Blood plasma levels of CA 19â€“9 monoclonal antibody that is often elevated in patients with pancreatic cancer. Only assessed in 350 patients (one goal of the study was to compare various CA 19-9 cutpoints from a blood sample to the model developed using urinary samples).
<i>creatinine</i>	Urinary biomarker of kidney function
<i>LYVE1</i>	Urinary levels of Lymphatic vessel endothelial hyaluronan receptor 1, a protein that may play a role in tumor metastasis
<i>REG1B</i>	Urinary levels of a protein that may be associated with pancreas regeneration.
<i>TFF1</i>	Urinary levels of Trefoil Factor 1, which may be related to regeneration and repair of the urinary tract
<i>REG1A</i>	Urinary levels of a protein that may be associated with pancreas regeneration. Only assessed in 306 patients (one goal of the study was to assess REG1B vs REG1A)

2.2 Data exploration


2.2.1 Diagnosis by sex

The first step was to explore if PDAC diagnosis was similar in male and female patients. A contingency table combined with a Chi-squared test were used to evaluate this. The results indicated that cancer diagnosis was significantly (p-value = 0.0001) more common in males than females. For instance, 58.3% of the cancer group were males.

2.2.2 Diagnosis by age

The distribution of diagnosis by age was also evaluated (Fig.1). This analysis revealed that cancer was most common in older people. Samples from people between 65-85 were more likely to be from the cancer group. For instance, more than 93% of cancer cases came from people older than 50 years.


Figure 1. Diagnosis by age



2.2.3 Biomarkers by diagnosis group and sex

The evaluation of the concentration of the levels of the different biomarkers revealed that all of them, except creatinine, were overexpressed in the cancer groups compared to healthy and benign groups, as seen in **Fig. 2**. Moreover, **Fig.2** also shows that there is no noticeable difference in levels of the different biomarkers between males and females.


Figure 2. Biomarkers levels by diagnosis group and sex



2.2.4 Biomarkers by age and diagnosis group

Figure 3 shows the average concentration of each of the biomarkers in cancer, benign, and healthy groups at different ages. This figure shows that the creatine levels are similar and steady across all ages, which suggests that this protein may not be a good predictor. On the other hand, the levels of the other biomarkers seem to be constantly higher in cancer samples across all ages. These differences are substantially larger in older patients, which appears to result from a drop of these proteins in healthy patients rather than an increase in sick patients.

Figure 3. Average biomarker concentration by age and diagnosis group



2.2 Models development

3. Results

4. Conclusion

5. References