

Parity bias in fundamental units of real quadratic fields

Florian Breuer

School of Information and Physical Sciences
University of Newcastle, Australia
florian.breuer@newcastle.edu.au

Cameron Shaw-Carmody

School of Information and Physical Sciences
University of Newcastle, Australia
cameron.shaw-carmody@newcastle.edu.au

August 9, 2024

Abstract

We compute primes $p \equiv 5 \pmod{8}$ up to 10^{11} for which the Pellian equation $x^2 - py^2 = -4$ has no solutions in odd integers; these are the members of sequence A130229 in the OEIS. We find that the number of such primes $p \leq x$ is well approximated by $\frac{1}{12}\pi(x) - 0.037 \int_2^x \frac{dt}{t^{1/6} \log t}$, where $\pi(x)$ is the usual prime counting function.

Keywords. Pellian equation; fundamental units; primes; OEIS A130229

2020 MSC. 11D09; 11R27; 11Y65

1 Introduction

For a prime $p \equiv 5 \pmod{8}$, consider the real quadratic field $K = \mathbb{Q}(\sqrt{p})$, with ring of integers $\mathcal{O}_K = \mathbb{Z}[\frac{1+\sqrt{p}}{2}]$ and fundamental unit $\varepsilon_p = \frac{x_0+y_0\sqrt{p}}{2} > 1$. Then (x_0, y_0) is a fundamental solution to the Pellian equation

$$x^2 - py^2 = -4. \tag{1}$$

The prime 2 is inert in K/\mathbb{Q} , and $\varepsilon_p \equiv 1 \pmod{2\mathcal{O}_K}$ if and only if the equation (1) has no odd integer solutions. Primes $p \equiv 5 \pmod{8}$ satisfying the above equivalent conditions define sequence A130229 in [OEIS]. They also appear in [B19, B21, XYY16].

Since $\varepsilon_p \pmod{2\mathcal{O}_K}$ can take any of three non-zero values in $\mathcal{O}_K/2\mathcal{O}_K \cong \mathbb{F}_4$, it is reasonable to expect roughly one third of all primes $p \equiv 5 \pmod{8}$ to be members of this sequence.

Define

$$\chi(p) := \begin{cases} 1 & \text{if } p \equiv 5 \pmod{8} \text{ and } \varepsilon_p \equiv 1 \pmod{2\mathcal{O}_K} \\ -\frac{1}{2} & \text{if } p \equiv 5 \pmod{8} \text{ and } \varepsilon_p \not\equiv 1 \pmod{2\mathcal{O}_K} \\ 0 & \text{if } p \not\equiv 5 \pmod{8} \end{cases}$$

and the modified counting function

$$\theta_\chi(x) := \sum_{p \leq x} \chi(p) \log p.$$

Then the above heuristic leads us to expect $\theta_\chi(x) = o(x)$ as $x \rightarrow \infty$.

In this note, we report on computations of $\theta_\chi(x)$ for $x \leq 10^{11}$, which show a surprising bias away from the $\varepsilon_p \equiv 1 \pmod{2\mathcal{O}_K}$ case, hinted at in related computations reported in [B19, §4]. We thus pose

Conjecture 1.1 *There exists a constant $c \approx -0.066$ for which*

$$\theta_\chi(x) \sim cx^{\frac{5}{6}}$$

as $x \rightarrow \infty$.

2 Results

We computed ε_p using the continued fraction method in [JW09, §3.3], with the modification that B_i and G_i are only computed modulo 2, since we only need to know the parity of ε_p . This significantly reduces the memory requirements of the calculation.

We implemented the algorithm to run on a GPU using the Python Numba library [LP15]. The final computation for all $p < 10^{11}$ took about 17 hours on an entry-level gaming laptop with an Nvidia RTX 3050 GPU. The source code and data are available at <https://github.com/florianbreuer/A130229>.

Table 1 lists some values for the naive counting function

$$\pi_1(x) = \sum_{p \leq x, \chi(p)=1} 1.$$

It is advantageous to study the “smoothed” counting function $\theta_\chi(x) = \sum_{p \leq x} \chi(p) \log p$. Figure 1 plots $-\theta_\chi(x)$ for $x \leq 10^{11}$ on logarithmic axes. The plot resembles a straight line with slope $5/6$. The least squares best fit of the form $f(x) = cx^{5/6}$ is found to have $c \approx -0.06626$, computed using the `find_fit` method in SageMath v9.3 [Sage]. The error term $\theta_\chi(x) - cx^{5/6}$ is shown in Figure 2. This provides evidence for Conjecture 1.1. Moreover, it appears likely that the error is of the order $O(x^{\frac{1}{2}+\epsilon})$.

From this we may also deduce a good approximation for $\pi_1(x)$. Define

$$\pi_{-\frac{1}{2}}(x) = \sum_{p \leq x, \chi(p)=-\frac{1}{2}} 1 \quad \text{and} \quad \pi_\chi(x) = \sum_{p \leq x} \chi(p) = \pi_1(x) - \frac{1}{2}\pi_{-\frac{1}{2}}(x).$$

x	$\pi_1(x)$	approximation	x	$\pi_1(x)$	approximation
10^2	1	1	$2 \cdot 10^{10}$	72770931	72761719
10^3	15	11	$3 \cdot 10^{10}$	107298975	107293481
10^4	98	90	$4 \cdot 10^{10}$	141363308	141357259
10^5	741	735	$5 \cdot 10^{10}$	175085540	175080418
10^6	6200	6187	$6 \cdot 10^{10}$	208542967	208537579
10^7	53382	53348	$7 \cdot 10^{10}$	241775700	241776120
10^8	468223	468144	$8 \cdot 10^{10}$	274823028	274829667
10^9	4164936	4165422	$9 \cdot 10^{10}$	307723656	307723171
10^{10}	37490293	37483463	10^{11}	340472393	340476359

Table 1: Some values of the counting function $\pi_1(x)$ for sequence A130229.




Figure 1: log-log plot of $-\theta_\chi(x)$ for $x \leq 10^{11}$.

Then $\theta_\chi(x) \sim cx^{5/6} \approx c \cdot \frac{5}{6} \int_2^x t^{-1/6} dt$ suggests

$$\pi_\chi(x) \approx c \cdot \frac{5}{6} \int_2^x \frac{t^{-1/6}}{\log t} dt \sim c \frac{x^{5/6}}{\log x}. \quad (2)$$

Then from $\pi_1(x) + \pi_{-\frac{1}{2}}(x) \approx \frac{1}{4}\pi(x)$, where $\pi(x)$ is the usual prime counting function, we arrive at

$$\pi_1(x) \approx \frac{1}{12}\pi(x) + \frac{2}{3}\pi_\chi(x) \approx \frac{1}{12}\pi(x) + c \cdot \frac{5}{9} \int_2^x \frac{t^{-1/6}}{\log t} dt. \quad (3)$$

These approximations are compared to the computed values of $\pi_1(x)$ in Table 1.

Acknowledgements.

The first author would like to thank James Punch for useful discussions, Frank Calegari for suggesting that the correct exponent is likely to be 5/6 and GPT-4 for help with porting the algorithm to a GPU.




Figure 2: Plot of the error term $\theta_\chi(x) - cx^{5/6}$ for $c \approx -0.06626$

References

- [B19] Breuer, F., Periods of Ducci sequences and odd solutions to a Pellian equation, *Bull. Aust. Math. Soc.* **100** (2019), 201–205. [doi:10.1017/S0004972719000212](https://doi.org/10.1017/S0004972719000212) 1, 2
- [B21] Breuer, F., Multiplicative orders of Gauss periods and the arithmetic of real quadratic fields, *Finite Fields Appl.* **73** (2021), 101848. <https://doi.org/10.1016/j.ffa.2021.101848> 1
- [JW09] Jacobson, M. J. and Williams, H. C., Solving the Pell Equation, *CMS Books Math./Ouvrages Math. SMC*, Springer, New York, 2009. 2
- [LP15] Lam, S. T., Pitrou, A. and Seibert, S., Numba: a LLVM-based Python JIT compiler, In: *Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC* (2015), Association for Computing Machinery, New York, NY, USA. <https://doi.org/10.1145/2833157.2833162> 2
- [OEIS] On-line Encyclopedia of Integer Sequences, entry #A130229, <https://oeis.org/A130229>. 1
- [Sage] The Sage Developers, *SageMath, the Sage Mathematics Software System (Version 9.3)* <https://www.sagemath.org>, 2021. 2
- [XY16] Xue, J., Yang, T.-C. and Yu, C.-F., Numerical invariants of totally imaginary quadratic $\mathbb{Z}[\sqrt{p}]$ -orders. *Taiwanese J. Math.* **20** (2016), no.4, 723–741. <https://doi.org/10.11650/tjm.20.2016.6464> 1