

# TopoChronia: A QGIS plugin for the creation of fully quantified palaeogeographic maps

Florian Franziskakis<sup>1</sup>, Christian Vérard<sup>2</sup>, Sébastien Castelltort<sup>2</sup>, and Grégory Giuliani<sup>1</sup>

<sup>1</sup> enviroSPACE group, Institute for Environmental Sciences, University of Geneva <sup>2</sup> Earth Surface Dynamics group, Department of Earth Sciences, University of Geneva

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

## Summary

Reconstructing the palaeogeography and palaeotopography of the Earth has been a challenge since the advent of the plate tectonics theory in the 1960s. With the development of geographic information systems (GIS), many plate tectonics models have been created and allowed researchers to reconstruct the movements of plates back in time (up to 1 billion years for some models), based on geological evidence found in the present-day Earth.

We present TopoChronia, a QGIS plugin that converts input data from plate tectonic models into quantified and synthetic topography. This plugin is optimized to work with the PANALESES model, because it is the only one currently providing sufficient information in terms of geological features to reconstruct a fully quantified topography.

## Statement of need

Most of the plate tectonic models and reconstructions use the standalone GPlates software (Gurnis et al., 2012), which allows users to move plates in time steps and export geospatial data layers. These layers can later be used in GIS software, such as the QGIS plugin TerraAntiqua (Aminov et al., 2023), to reconstruct palaeotopography. Other models such as PANALESES (Vérard, 2019), are created and have processing functionalities that use commercial GIS software (ArcGIS). A preliminary version of the code to generate topography of the Earth based on PANALESES past was developed as an ArcGIS extension, written in Visual Basic .NET but never published. It is now fully updated as a QGIS plugin in Python.

Constraining the palaeotopography is critical in fields such as climate and mantle dynamics modelling, as the elevation of land and bathymetry of oceans are used to set the initial conditions of models (Bello et al., 2015; Ragon et al., 2023). Quantifying the Earth's topography and its evolution also allows to estimate the volume of rocks being eroded, for instance through sediment discharge (Lyster et al., 2020), as weathering of silicate rocks is a key controlling factor of CO<sub>2</sub> concentration in the atmosphere over geological time scales (Macdonald et al., 2019; Molnar & England, 1990).

The traditional method to create palaeotopographic maps (Scotese, 2021) is to use present-day geological evidence, rotate them back to their past location and derive semi-quantitative elevation typical of the environment they depict. Another method is to take the present-day Earth topography of an area of interest as it is, and rotate it back in time to its past location (Aminov et al., 2023). These methods have limitations, including that present-day features are the result of millions of years of plate movements and cannot be “copy-pasted” as such, and that one time step might not be coherent with the previous and next ones.

We provide here an open-source plugin to reconstruct palaeotopography and palaeogeography “from scratch” using the PANALESES model, which is based on present-day geological evidence

and uses a dual-control approach, meaning that one reconstruction is based on the state of the Earth in the previous time-step, and influences the next step. Synthetic values for elevations are generated in nodes (points) related to geological settings and based on their present-day counterparts (Vérard, 2017). The output maps of TopoChronia can be used for modelling purposes and to reconstruct sea-level curves, over the Phanerozoic and beyond (Vérard et al., 2015).

## Functionalities

TopoChronia is divided into three main parts:

### 1. Check Configuration

- Assess input data files (geometry, field names, values)
- Perform manual corrections if necessary, for wrongly named fields
- Define output folder path
- Extract available reconstruction ages

### 2. Create Node Grid

- Select input lines from plate model file
- Convert mid-oceanic ridge and isochron features and interpolate a preliminary raster for oceans
- Convert all other features (abandoned arcs, continents, cratons, lower subduction, upper subduction, passive margin wedges, continent sides, hot-spots, other margins, rifts, and collision zones)
- Merge all nodes and clean to avoid clashing between features

### 3. Interpolate to Raster

- Interpolate global raster
- Calculate oceanic volume under sea-level (elevation below 0m)
- Calculate required sea-level increase to match present-day oceanic volume
- Correct water load using Airy's model to adjust for sea-level increase
- Perform final raster interpolation with new sea-level

Each reconstruction will yield the following outputs:

- A palaeogeographic map in geotiff format with cylindrical equal-area projection (ESRI:54034): raster\_final\_filled\_{age}.tif
- A text file summarizing sea-level information **before** water load correction (initial volume and area, added water column, sea-level increase and subsidence): water\_load\_correction\_summary.txt
- A text file summarizing sea-level information **after** water load correction: water\_load\_correction\_summary\_f.txt
- All nodes both in EPSG:4326 and ESRI:54034 projections: all\_nodes\_{age}.geojson and reproj\_all\_nodes\_{age}.geojson
- All other processing products from line to points for each setting.

## Acknowledgements

The authors acknowledge financial support from the Swiss National Science Foundation (SNSF) under [Sinergia grant #213539](#): *Long-term evolution of the Earth from the base of the mantle to the top of the atmosphere: Understanding the mechanisms leading to 'greenhouse' and 'icehouse' regimes*.

The authors would like to thank Niklas Werner, Felipe Carlos and Bastien Deriaz for their help in testing the plugin.

## References

- Aminov, J., Dupont-Nivet, G., Ruiz, D., & Gailleton, B. (2023). Paleogeographic reconstructions using QGIS: Introducing Terra Antiqua plugin and its application to 30 and 50 Ma maps. *Earth-Science Reviews*, 240, 104401. <https://doi.org/10.1016/j.earscirev.2023.104401>
- Bello, L., Coltice, N., Tackley, P. J., Dietmar Müller, R., & Cannon, J. (2015). Assessing the role of slab rheology in coupled plate-mantle convection models. *Earth and Planetary Science Letters*, 430, 191–201. <https://doi.org/10.1016/j.epsl.2015.08.010>
- Gurnis, M., Turner, M., Zahirovic, S., DiCaprio, L., Spasojevic, S., Müller, R. D., Boyden, J., Seton, M., Manea, V. C., & Bower, D. J. (2012). Plate tectonic reconstructions with continuously closing plates. *Computers & Geosciences*, 38(1), 35–42. <https://doi.org/10.1016/j.cageo.2011.04.014>
- Lyster, S. J., Whittaker, A. C., Allison, P. A., Lunt, D. J., & Farnsworth, A. (2020). Predicting sediment discharges and erosion rates in deep time—examples from the late Cretaceous North American continent. *Basin Research*, 32(6), 1547–1573. <https://doi.org/10.1111/bre.12442>
- Macdonald, F. A., Swanson-Hysell, N. L., Park, Y., Lisiecki, L., & Jagoutz, O. (2019). Arc-continent collisions in the tropics set Earth's climate state. *Science*, 364(6436), 181–184. <https://doi.org/10.1126/science.aav5300>
- Molnar, P., & England, P. (1990). Late Cenozoic uplift of mountain ranges and global climate change: Chicken or egg? *Nature*, 346(6279), 29–34. <https://doi.org/10.1038/346029a0>
- Ragon, C., Vêrard, C., Kasparian, J., & Brunetti, M. (2023). Alternative climatic steady states for the Permian–Triassic paleogeography. *EGUsphere*, 1–31. <https://doi.org/10.5194/egusphere-2023-1808>
- Scotese, C. (2021). An Atlas of Phanerozoic Paleogeographic Maps: The Seas Come In and the Seas Go Out. *Annual Review of Earth and Planetary Sciences*, 49(Volume 49, 2021), 679–728. <https://doi.org/10.1146/annurev-earth-081320-064052>
- Vêrard, C. (2017). Statistics of the Earth's Topography. *OALib*, 04(06), 1–50. <https://doi.org/10.4236/oalib.1103398>
- Vêrard, C. (2019). Panalexis: Towards global synthetic palaeogeographies using integration and coupling of manifold models. *Geological Magazine*, 156(2), 320–330. <https://doi.org/10.1017/S0016756817001042>
- Vêrard, C., Hochard, C., Baumgartner, P. O., Stampfli, G. M., & Liu, M. (2015). 3D palaeogeographic reconstructions of the Phanerozoic versus sea-level and Sr-ratio variations. *Journal of Palaeogeography*, 4(1), 64–84. <https://doi.org/10.3724/SP.J.1261.2015.00068>