

# On the Erdős-Straus Conjecture: Algebraic Analysis and Modular Decomposition

Charles EDOU NZE

August 14, 2026

## Abstract

This document formally addresses the Erdős-Straus conjecture, which asserts that for all integers  $n \geq 2$ , the equation  $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  possesses positive integer solutions  $x, y, z$ . The structure encompasses strict axiomatic definitions, a review of pertinent literature focusing on modular sieves and divisor parametrization, a decomposition into congruence classes modulo 4 and 3, and detailed derivations covering these classes. A formal architecture in Lean 4 is subsequently outlined.

*Charles EDOU NZE, chercheur indépendant*

## Contents

|   |                                         |   |
|---|-----------------------------------------|---|
| 1 | Axiomatic Definitions                   | 2 |
| 2 | Contextual Literature                   | 2 |
| 3 | Lemma Isolation and Zero-Ellipse Proofs | 2 |
| 4 | Architecture for Autoformalization      | 4 |

# 1 Axiomatic Definitions

**Definition 1.1.** Let  $n \in \mathbb{N}$  with  $n \geq 2$ . A positive integer triplet  $(x, y, z) \in (\mathbb{N}_{>0})^3$  is defined as a solution to the Erdős-Straus Diophantine equation if it satisfies:

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

**Definition 1.2.** We define the predicate  $\mathcal{E}(n)$  to be true if and only if there exists a solution triplet  $(x, y, z) \in (\mathbb{N}_{>0})^3$  for a given integer  $n \geq 2$ .

# 2 Contextual Literature

The Erdős-Straus conjecture formulates a fundamental question concerning the representation of rational numbers as sums of unit fractions. Significant advances have relied on analyzing the equation via modular arithmetic. Approaches involving modular sieves have empirically verified the conjecture up to  $N = 10^{17}$  (Salez, 2014) and beyond. Theoretical studies extensively employ divisor parametrizations and polynomial identities. A prominent analytic result establishes that the number of solutions is bounded polylogarithmically on average. The methodology developed herein leverages these polynomial identities parameterized by congruence classes.

# 3 Lemma Isolation and Zero-Ellipse Proofs

The reduction of the conjecture to prime values  $n = p$  is standard, as a solution for a prime factor  $p$  can be extended to composite numbers via scalar multiplication. We establish specific lemmas for distinct congruence classes.

**Lemma 3.1.** For any integer  $k \geq 0$ , if  $n = 4k + 3$ , then  $\mathcal{E}(n)$  holds.

*Proof.* Let  $n = 4k + 3$  for  $k \in \mathbb{N}$ . We consider the rational function expansion of  $\frac{4}{4k+3}$ . We define the triplet of variables  $(x, y, z)$  as follows:

$$x = k + 1$$

$$y = 2(4k + 3)(k + 1)$$

$$z = 2(4k + 3)(k + 1)$$

We evaluate the sum of the reciprocal unit fractions corresponding to this assignment:

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{1}{k + 1} + \frac{1}{2(4k + 3)(k + 1)} + \frac{1}{2(4k + 3)(k + 1)}$$

Since the second and third fractions are identical, we sum them:

$$\frac{1}{2(4k + 3)(k + 1)} + \frac{1}{2(4k + 3)(k + 1)} = \frac{2}{2(4k + 3)(k + 1)} = \frac{1}{(4k + 3)(k + 1)}$$

Substituting this back into the original sum:

$$\frac{1}{k + 1} + \frac{1}{(4k + 3)(k + 1)}$$

To combine these fractions, we identify a common denominator, which is  $(4k+3)(k+1)$ . We multiply the numerator and denominator of the first fraction by  $(4k+3)$ :

$$\frac{4k+3}{(4k+3)(k+1)} + \frac{1}{(4k+3)(k+1)} = \frac{4k+4}{(4k+3)(k+1)}$$

We factor the numerator as  $4(k+1)$ :

$$\frac{4(k+1)}{(4k+3)(k+1)}$$

Since  $k \geq 0$ , the term  $(k+1)$  is non-zero, allowing us to cancel it from the numerator and denominator:

$$\frac{4}{4k+3} = \frac{4}{n}$$

This demonstrates that  $x = k+1$ ,  $y = 2(4k+3)(k+1)$ , and  $z = 2(4k+3)(k+1)$  constitute a valid integer triplet solution for  $n = 4k+3$ . Since  $k \geq 0$ , all denominators are strictly positive integers.  $\square$

**Lemma 3.2.** *For any integer  $k \geq 1$ , if  $n = 3k+2$ , then  $\mathcal{E}(n)$  holds.*

*Proof.* Let  $n = 3k+2$  for  $k \in \mathbb{N}_{>0}$ . We define  $x = k+1$ . The difference is evaluated as:

$$\frac{4}{3k+2} - \frac{1}{k+1} = \frac{4(k+1) - (3k+2)}{(3k+2)(k+1)} = \frac{4k+4-3k-2}{(3k+2)(k+1)} = \frac{k+2}{(3k+2)(k+1)}$$

We require  $\frac{k+2}{(3k+2)(k+1)}$  to be represented as  $\frac{1}{y} + \frac{1}{z}$ . We use the algebraic identity  $\frac{A+B}{C} = \frac{A}{C} + \frac{B}{C}$ . Let  $A = k+1$  and  $B = 1$ . Then  $A+B = k+2$ .

$$\frac{k+2}{(3k+2)(k+1)} = \frac{k+1}{(3k+2)(k+1)} + \frac{1}{(3k+2)(k+1)}$$

Simplifying the first fraction by canceling the common factor  $(k+1)$ :

$$\frac{k+1}{(3k+2)(k+1)} = \frac{1}{3k+2}$$

Thus, the difference decomposes exactly into two unit fractions:

$$\frac{1}{3k+2} + \frac{1}{(3k+2)(k+1)}$$

We assign  $y = 3k+2$  and  $z = (3k+2)(k+1)$ . Since  $k \geq 1$ ,  $x = k+1$ ,  $y = 3k+2$ , and  $z = (3k+2)(k+1)$  are all strictly positive integers. The full identity is:

$$\frac{4}{3k+2} = \frac{1}{k+1} + \frac{1}{3k+2} + \frac{1}{(3k+2)(k+1)}$$

This concludes the proof for this congruence class.  $\square$

## 4 Architecture for Autoformalization

The following definitions and types present a structure for translating the problem and established modular identities into Lean 4.

```
import Mathlib.Data.Nat.Basic
import Mathlib.Data.Rat.Basic

open Nat

-- Axiomatic definition of the Erdős-Straus property for an
-- integer n
def SatisfiesErdosStraus (n : Nat) : Prop :=
  Exists (fun x => Exists (fun y => Exists (fun z =>
    x > 0 /\ y > 0 /\ z > 0 /\
    (4 : Rat) / n = (1 : Rat) / x + (1 : Rat) / y + (1 : Rat) / z))
  )

-- Lemma for the congruence class  $n = 4k + 3$ 
lemma erdos_straus_mod_4_3 (k : Nat) : SatisfiesErdosStraus (4 * k
+ 3) := by
  admit

-- Lemma for the congruence class  $n = 3k + 2$ 
lemma erdos_straus_mod_3_2 (k : Nat) (hk : k >= 1) :
  SatisfiesErdosStraus (3 * k + 2) := by
  admit

-- General statement of the Erdős-Straus conjecture
theorem erdos_straus_conjecture (n : Nat) (hn : n >= 2) :
  SatisfiesErdosStraus n := by
  admit
```