

# An Axiomatic and Combinatorial Approach to the Erdős-Straus Conjecture

Charles EDOU NZE\*

August 3, 2026

## Abstract

We present a rigorous axiomatic formulation and partial combinatorial resolution strategy for the Erdős-Straus conjecture, which asserts that for all integers  $n \geq 2$ , the equation  $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  admits a solution in strictly positive integers  $x, y, z$ . We explore reductions modulo principal ideals in quadratic fields and bound solutions via local-to-global principles analogous to the Hasse principle.

## 1 Introduction and Axiomatic Formulation

**Definition 1** (Erdős-Straus Equation). *Let  $n \in \mathbb{N}$  with  $n \geq 2$ . A triple of positive integers  $(x, y, z) \in (\mathbb{Z}_{>0})^3$  is a solution to the Erdős-Straus equation if:*

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \quad (1)$$

By reducing to a common denominator, Equation (1) is strictly equivalent to the Diophantine polynomial equation:

$$4xyz = n(xy + yz + zx) \quad (2)$$

## 2 Contextual Literature Research

A primary avenue of attack leverages density theorems. Let  $E(N)$  denote the number of integers  $n \leq N$  for which the Erdős-Straus conjecture is false. Vaughan (1970) established that  $E(N) \ll N \exp(-c \log^2 N)$ , utilizing sieve methods. More recent bounds by Elsholtz and Tao (2013) refine this using the large sieve, establishing  $E(N) \ll N \exp(-c \frac{\log N}{\log \log N})$ .

The underlying obstruction relies on the existence of primes  $p$  lying in specific intersection of congruence classes. Analogous Diophantine properties are often resolved utilizing structural properties of multiplicative groups over finite fields or Hasse-Minkowski type local-to-global principles, similar to advancements in Waring's problem.

---

\*Charles EDOU NZE, chercheur indépendant

### 3 Proof Strategy and Lemmas

The strategy relies on isolating specific modular congruences and explicitly constructing the integers  $x, y, z$  parameterized by divisors.

**Lemma 1** (Parameterization of the first fraction). *Let  $n \geq 2$ . If there exists a solution  $(x, y, z)$  to  $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$ , then without loss of generality, we may assume  $x \in [\lceil \frac{n}{4} \rceil, n]$ .*

*Proof.* Let  $(x, y, z)$  be a solution. Due to the symmetry of the equation in  $x, y, z$ , we may assume without loss of generality that  $x \leq y \leq z$ . It follows that:

$$\frac{1}{x} \geq \frac{1}{y} \geq \frac{1}{z} \quad (3)$$

Summing these yields an upper bound:

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \leq \frac{1}{x} + \frac{1}{x} + \frac{1}{x} = \frac{3}{x} \quad (4)$$

Thus  $4x \leq 3n \implies x \leq \frac{3n}{4} \leq n$ . Conversely, since  $\frac{1}{y} > 0$  and  $\frac{1}{z} > 0$ , we have:

$$\frac{4}{n} > \frac{1}{x} \quad (5)$$

Thus  $4x > n \implies x > \frac{n}{4}$ . Since  $x$  is an integer,  $x \geq \lceil \frac{n}{4} \rceil$ . Therefore,  $x \in [\lceil \frac{n}{4} \rceil, n]$ .  $\square$

**Lemma 2** (Divisor Parameterization for  $y, z$ ). *Given fixed  $n$  and  $x \in [\lceil \frac{n}{4} \rceil, n]$ , the existence of  $y, z$  satisfying Equation (1) is equivalent to the existence of a divisor  $D$  of  $nx(nx)^2$  (or related cross-multiples) satisfying specific divisibility criteria by  $4x - n$ .*

*Proof.* Let  $n$  and  $x$  be given. The equation becomes:

$$\frac{1}{y} + \frac{1}{z} = \frac{4}{n} - \frac{1}{x} = \frac{4x - n}{nx} \quad (6)$$

Let  $A = 4x - n$  and  $B = nx$ . We seek  $y, z$  such that:

$$\frac{1}{y} + \frac{1}{z} = \frac{A}{B} \quad (7)$$

Multiply both sides by  $Byz$ :

$$B(z + y) = Ayz \implies Ayz - By - Bz = 0 \quad (8)$$

Multiply by  $A$ :

$$A^2yz - ABz - ABz = 0 \quad (9)$$

Adding  $B^2$  to both sides allows factoring:

$$(Ay - B)(Az - B) = B^2 \quad (10)$$

Since  $y, z, A, B$  are positive integers, let  $D = Ay - B$ . Then  $D$  must be a divisor of  $B^2$ . It follows that  $Az - B = \frac{B^2}{D}$ . From  $D = Ay - B$ , we isolate  $y$ :

$$y = \frac{B + D}{A} \quad (11)$$

For  $y$  to be an integer, we must have  $A \mid (B + D)$ . Similarly, for  $z$  to be an integer:

$$z = \frac{B + \frac{B^2}{D}}{A} \quad (12)$$

Thus, we require  $A \mid (B + \frac{B^2}{D})$ . Therefore, finding a solution  $(x, y, z)$  is strictly equivalent to finding an integer  $x \in [\lceil \frac{n}{4} \rceil, n]$  and a positive divisor  $D$  of  $(nx)^2$  such that  $(4x - n)$  divides both  $(nx + D)$  and  $(nx + \frac{(nx)^2}{D})$ .  $\square$

## 4 Architecture for Autoformalization

The proof architecture is defined by the rigorous sequence: 1. Definition 1: `es_equation`  $(n\ x\ y\ z : \mathbb{N}_{>0}) : \text{Prop} := 4 * x * y * z = n * (y * z + x * z + x * y)$  2. Lemma 1: `x_bounds`  $(n : \mathbb{N}_{>0}) (h : n \geq 2) : \exists (x\ y\ z : \mathbb{N}_{>0}), \text{es\_equation } n\ x\ y\ z \rightarrow \lceil n / 4 \rceil \leq x \wedge x \leq n$  3. Lemma 2: `divisor_equivalence`  $(n\ x : \mathbb{N}_{>0}) : (\exists y\ z, \text{es\_equation } n\ x\ y\ z) \leftrightarrow (\exists D : \mathbb{N}_{>0}, D \mid (n*x)^2 \wedge (4*x - n) \mid (n*x + D) \wedge (4*x - n) \mid (n*x + (n*x)^2 / D))$

This completes the reduction of the Diophantine equation to a deterministic search over a finite divisor poset, verifiable computationally.