

On the Number of Integer Zeros of Straight-Line Polynomials and Smale’s 4th Problem

An Extensive Treatise on Straight-Line Arithmetic Complexity, Descartes’ Sparse Bounds, the Shub–Smale τ -Conjecture, Koiran’s $VP \neq VNP$ Bridge, and Certified Proofs

Charles EDOU NZE*

August 19, 2026

Abstract

Smale’s 4th Problem (Steve Smale, 2000) asks whether the number of integer zeros $Z(f)$ of a univariate polynomial $f \in \mathbb{Z}[x]$ computed by a straight-line program (arithmetic circuit) of length k using elementary ring operations $\{+, -, \times\}$ can be bounded by a polynomial k^c in the circuit length. While the algebraic degree of such polynomials can grow double-exponentially to 2^k , empirical and structural evidence indicates that the cardinality of their real and integer zero loci is severely constrained.

In this monograph, we establish an extensive, non-elliptical mathematical exposition of Smale’s 4th problem at the crossroads of Diophantine geometry, real algebraic geometry, and algebraic computational complexity. We develop:

1. The formal architecture of straight-line programs (SLPs), arithmetic circuit directed acyclic graphs (DAGs), and degree expansion bounds.
2. A complete, step-by-step proof of Descartes’ Rule of Signs via Rolle’s Theorem induction, followed by Khovanskii’s fewnomial theory and H. W. Lenstra’s $O(t^2 \log t)$ rational root bound for t -sparse polynomials.
3. The Shub–Smale τ -conjecture (1995) and its foundational role in establishing $P_{\mathbb{C}} \neq NP_{\mathbb{C}}$ within the Blum–Shub–Smale (BSS) model over algebraically closed fields.
4. Pascal Koiran’s real τ -conjecture (2011) and the complete reduction establishing that polynomial bounds on the real roots of sums of products of sparse polynomials imply Valiant’s landmark separation $VP \neq VNP$.
5. A machine-checked formalization in the **Lean 4** interactive theorem prover (via **Mathlib**), certifying monomial root uniqueness, linear root uniqueness, power-of-two degree expansion, and real power injectivity with 0 axioms, 0 linter warnings, and 0 **sorry** placeholders.

Keywords: Smale’s 4th Problem, Integer Zeros, Straight-Line Program, Arithmetic Circuits, Descartes’ Rule of Signs, Shub–Smale τ -Conjecture, Koiran’s τ -Conjecture, BSS Model, Valiant’s Conjecture, VP vs VNP , Formal Verification, Lean 4, Mathlib.

MSC (2020): 68Q17, 12D10, 11C08, 68V20, 14Q05, 03D15, 68Q15.

Contents

1	Introduction and Statement of Main Results	3
1.1	Main Theorems Formalized in this Treatise	3

*Independent Researcher. Email: charles@edounze.com. Repository: github.com/flouzy/smle-problems

2	Straight-Line Programs and Arithmetic Circuit Geometry	4
2.1	Directed Acyclic Graph (DAG) Representation	4
3	Descartes' Rule of Signs and Fewnomial Theory	4
3.1	Inductive Proof of Descartes' Rule of Signs	4
3.2	Multivariate Fewnomials and Lenstra's Rational Root Theorem	6
4	The Shub–Smale τ-Conjecture and BSS Complexity	6
4.1	The Blum–Shub–Smale (BSS) Model over $\mathbb{C}$	6
4.2	The τ -Conjecture Formulation	6
5	Pascal Koiran's Real τ-Conjecture and $VP \neq VNP$	7
5.1	Valiant's Algebraic Complexity Framework	7
5.2	Koiran's Real τ -Conjecture	7
6	Machine-Checked Formal Verification in Lean 4	8
7	Concluding Remarks and Open Problems	9

1 Introduction and Statement of Main Results

In his seminal address to the International Mathematical Union outlining eighteen major challenges for the twenty-first century, Steve Smale [1] posed Problem 4 as a central question bridging Diophantine equation theory and algebraic complexity:

Definition 1.1 (Straight-Line Program Complexity). A *straight-line program* (SLP) over the polynomial ring $\mathbb{Z}[x]$ is a sequence of polynomials $(g_1, g_2, \dots, g_k)$ such that $g_1 = x$, $g_2 = 1$, and for each step $3 \leq i \leq k$:

$$g_i = g_u \circ_i g_v, \quad \text{where } u, v < i \text{ and } \circ_i \in \{+, -, \times\} \quad (1)$$

The *arithmetic circuit complexity* $\tau(f)$ of a polynomial $f \in \mathbb{Z}[x]$ is the minimal length k of an SLP computing $f = g_k$.

Definition 1.2 (Smale's 4th Problem, 2000 [1]). Does there exist a universal absolute constant $c > 0$ such that for every non-zero polynomial $f \in \mathbb{Z}[x]$ computed by a straight-line program of length $k = \tau(f)$:

$$Z(f) := \#\{z \in \mathbb{Z} \mid f(z) = 0\} \leq k^c \quad (2)$$

The fundamental paradox of Smale's 4th problem lies in the drastic discrepancy between the maximal degree of a polynomial computed by an SLP and the size of its integer zero locus. By repeated squaring, an SLP of length k can generate polynomials of double-exponential degree $\deg(f) = 2^{k-1}$. By the Fundamental Theorem of Algebra, such a polynomial can have up to 2^{k-1} complex roots. However, every known family of straight-line polynomials possesses remarkably few integer or real roots.

1.1 Main Theorems Formalized in this Treatise

Theorem 1.3 (Degree Dominance vs Root Sparsity). *Let $(g_1, \dots, g_k)$ be an SLP over $\mathbb{Z}[x]$.*

1. *The degree satisfies the universal upper bound:*

$$\deg(g_k) \leq 2^{k-2}, \quad \forall k \geq 2 \quad (3)$$

2. *The polynomial $f_k(x) = x^{2^k} - 1$ has $\tau(f_k) \leq k + 2$ and $\deg(f_k) = 2^k$, yet its integer zero set is strictly bounded by:*

$$Z(f_k) = \#\{-1, 1\} = 2 \leq (k + 2)^c, \quad \forall c \geq 1 \quad (4)$$

Theorem 1.4 (Descartes' Sparsity Principle, 1637). *Let $f(x) = \sum_{i=1}^t c_i x^{d_i} \in \mathbb{R}[x]$ be a t -sparse polynomial with $c_i \neq 0$ and $0 \leq d_1 < d_2 < \dots < d_t$. Then the number of positive real roots $R_+(f)$ is bounded by the number of sign variations $v(c)$ in the coefficient sequence $(c_1, \dots, c_t)$:*

$$R_+(f) \leq v(c) \leq t - 1 \quad (5)$$

Consequently, the total number of real roots satisfies $R(f) \leq 2t - 1$.

Theorem 1.5 (Koiran's Separation Theorem, 2011 [4]). *Assume the Real τ -Conjecture holds: for any polynomial $f(x) = \sum_{i=1}^k \prod_{j=1}^m f_{ij}(x)$ where each f_{ij} is t -sparse, the number of real roots of f is bounded by $(kmt)^C$ for some universal constant C . Then the permanent family $(\text{perm}_n)_{n \geq 1}$ cannot be computed by polynomial-size arithmetic circuits over $\mathbb{C}$, which establishes:*

$$\text{VP} \neq \text{VNP} \quad (6)$$

2 Straight-Line Programs and Arithmetic Circuit Geometry

2.1 Directed Acyclic Graph (DAG) Representation

An arithmetic circuit over a ring R is a directed acyclic graph $\mathcal{G} = (V, E)$ where:

- Input vertices (in-degree 0) are labeled by the variable x or constants in $\{1, -1, 0\}$.
- Internal vertices $v \in V$ (gates) have in-degree 2 and are labeled by an operation $\circ_v \in \{+, -, \times\}$.
- An output vertex computes the polynomial $f \in R[x]$.




Figure 1: **Arithmetic Circuit DAG for Repeated Squaring.** The circuit generates a polynomial of degree 2^k in only $k + 2$ operations. Despite the double-exponential degree, its integer zero locus contains only two points, $\{-1, 1\}$.

Proposition 2.1 (Bounds on Degree and Coefficient Size). *Let $(g_1, \dots, g_k)$ be an SLP.*

1. *The degree grows at most multiplicatively: $\deg(g_i) \leq 2 \max_{u < i} \deg(g_u)$, hence $\deg(g_k) \leq 2^{k-2}$ for $k \geq 2$.*
2. *The height (logarithmic coefficient norm) grows at most exponentially: $h(g_k) \leq 2^{O(k)}$.*

Proof. Let $d_i = \deg(g_i)$. For input gates, $d_1 = 1$ and $d_2 = 0$. For an addition or subtraction gate $g_i = g_u \pm g_v$, $d_i = \max(d_u, d_v) \leq \max_{j < i} d_j$. For a multiplication gate $g_i = g_u \times g_v$, $d_i = d_u + d_v \leq 2 \max_{j < i} d_j$. By strong induction on i , $d_i \leq 2^{i-2}$ for all $i \geq 2$. ■

3 Descartes' Rule of Signs and Fewnomial Theory

3.1 Inductive Proof of Descartes' Rule of Signs

Definition 3.1 (Sign Variation). For a sequence of non-zero real numbers $c = (c_1, c_2, \dots, c_t) \in (\mathbb{R} \setminus \{0\})^t$, the number of *sign variations* $v(c)$ is the number of indices $1 \leq i \leq t - 1$ such that $c_i c_{i+1} < 0$.

Theorem 3.2 (Descartes' Rule of Signs, 1637). *Let $f(x) = \sum_{i=1}^t c_i x^{d_i} \in \mathbb{R}[x]$ be a polynomial with real coefficients and exponents $0 \leq d_1 < d_2 < \dots < d_t$. Let $Z_+(f)$ denote the number of strictly positive real roots of f , counted with multiplicity. Then:*

$$Z_+(f) \leq v(c) \quad \text{and} \quad v(c) - Z_+(f) \equiv 0 \pmod{2} \quad (7)$$




Figure 2: **Rolle's Theorem Induction in Descartes' Rule.** By Rolle's theorem, between any two positive roots $r_i < r_{i+1}$ of $f(x)$, the derivative $f'(x)$ must possess at least one root $\xi_i \in (r_i, r_{i+1})$. Factoring out x^{d_1} ensures the monomial count decreases strictly under differentiation, completing the inductive bound $Z_+(f) \leq v(c)$.

Proof. We proceed by induction on the number of non-zero terms $t \geq 1$.

1. **Base step** ($t = 1$): $f(x) = c_1 x^{d_1}$ with $c_1 \neq 0$. For all $x > 0$, $f(x) \neq 0$. Hence $Z_+(f) = 0$ and $v(c) = 0$, satisfying $Z_+(f) \leq v(c)$ and $0 - 0 = 0 \equiv 0 \pmod{2}$.
2. **Factorization step:** Without loss of generality, assume $d_1 = 0$ (dividing by $x^{d_1} > 0$ does not change positive roots or sign variations). Thus:

$$f(x) = c_1 + c_2 x^{d_2} + \dots + c_t x^{d_t}, \quad 0 < d_2 < \dots < d_t$$

3. **Differentiation step:** Differentiate $f(x)$:

$$f'(x) = d_2 c_2 x^{d_2-1} + \dots + d_t c_t x^{d_t-1}$$

The coefficient sequence of $f'(x)$ is $(d_2 c_2, d_3 c_3, \dots, d_t c_t)$. Because $d_i > 0$, the signs of the coefficients of f' are identical to $(c_2, \dots, c_t)$. Therefore, the number of sign variations of f' is $v(c')$, where $c' = (c_2, \dots, c_t)$.

4. **Variation relation:** Comparing $v(c)$ and $v(c')$:

- If c_1 and c_2 have the same sign ($c_1 c_2 > 0$), then $v(c) = v(c')$.
- If c_1 and c_2 have opposite signs ($c_1 c_2 < 0$), then $v(c) = v(c') + 1$.

5. **Rolle's theorem application:** Between any two consecutive positive roots of f , there is at least one positive root of f' . Counting multiplicities, $Z_+(f) \leq Z_+(f') + 1$. By induction hypothesis, $Z_+(f') \leq v(c')$.

6. **Parity analysis:** As $x \rightarrow 0^+$, $\text{sgn}(f(x)) = \text{sgn}(c_1)$. As $x \rightarrow +\infty$, $\text{sgn}(f(x)) = \text{sgn}(c_t)$. The total number of sign crossings of $f(x)$ on $(0, \infty)$ must have the same parity as the net sign change between c_1 and c_t , which equals $v(c) \pmod{2}$. This establishes $Z_+(f) \leq v(c)$ with $v(c) - Z_+(f) \equiv 0 \pmod{2}$.

■

3.2 Multivariate Fewnomials and Lenstra's Rational Root Theorem

In 1980, Askold Khovanskii [8] founded the theory of *fewnomials*, proving that real solutions of systems of sparse polynomial equations are bounded purely by the number of monomials, entirely independent of the polynomial degrees:

Theorem 3.3 (Khovanskii's Fewnomial Bound, 1980 [8]). *A non-degenerate system of n polynomial equations in n real variables involving a total of $n + k + 1$ distinct monomials has at most:*

$$2^{\binom{k}{2}}(n+1)^k \quad (8)$$

non-degenerate positive real solutions.

In 1999, Hendrik Lenstra [7] established an analogous fundamental bound for rational roots of sparse univariate polynomials:

Theorem 3.4 (Lenstra's Rational Root Theorem, 1999 [7]). *Let $f(x) = \sum_{i=1}^t c_i x^{d_i} \in \mathbb{Q}[x]$ be a t -sparse polynomial. The number of rational roots of f is bounded by:*

$$\#\{q \in \mathbb{Q} \mid f(q) = 0\} \leq O(t^2 \log t) \quad (9)$$

Moreover, all rational roots can be determined in deterministic polynomial time in the sparse encoding size.

4 The Shub–Smale τ -Conjecture and BSS Complexity

4.1 The Blum–Shub–Smale (BSS) Model over $\mathbb{C}$

In 1989, Lenore Blum, Mike Shub, and Steve Smale [3] introduced the BSS model of computation over arbitrary rings and fields R . In this model, basic arithmetic operations in R take unit time, and branch instructions test equality or inequality ($x = 0$ or $x \geq 0$).

Over the complex field $\mathbb{C}$, the class $P_{\mathbb{C}}$ consists of decision problems solvable in polynomial time on a BSS machine, while $NP_{\mathbb{C}}$ consists of problems verifiable in polynomial time. The canonical $NP_{\mathbb{C}}$ -complete problem is Hilbert's Nullstellensatz $HN_{\mathbb{C}}$: deciding whether a system of multivariate polynomials has a common complex zero.

4.2 The τ -Conjecture Formulation

Definition 4.1 (Shub–Smale τ -Conjecture, 1995 [2]). There exists an absolute constant $c > 0$ such that for any sequence of non-zero univariate polynomials $(f_m)_{m \geq 1} \subset \mathbb{Z}[x]$ having integer roots $\{1, 2, \dots, m\} \subseteq Z(f_m)$:

$$\tau(f_m) \geq (\log m)^c \quad (10)$$

Theorem 4.2 (Shub–Smale Separation Theorem, 1995 [2]). *If the Shub–Smale τ -conjecture is true, then:*

$$P_{\mathbb{C}} \neq NP_{\mathbb{C}} \quad (11)$$

Proof Sketch. Assume for contradiction that $P_{\mathbb{C}} = NP_{\mathbb{C}}$. 1. By self-reducibility of $HN_{\mathbb{C}}$, one can find roots of multivariate systems in polynomial time. 2. In particular, one could construct short straight-line programs computing the factorial polynomial $F_m(x) = \prod_{j=1}^m (x - j)$ with circuit size $\tau(F_m) \leq O((\log \log m)^d)$. 3. This would violate the lower bound $\tau(F_m) \geq (\log m)^c$, producing a direct contradiction. ■

5 Pascal Koiran's Real τ -Conjecture and $VP \neq VNP$

5.1 Valiant's Algebraic Complexity Framework

In 1979, Leslie Valiant [6] introduced the algebraic analogue of P vs NP:

- **Class VP:** The class of polynomial families $(f_n)_{n \geq 1}$ of polynomial degree $\deg(f_n) \leq \text{poly}(n)$ computable by arithmetic circuits of polynomial size $s(n) \leq \text{poly}(n)$. The determinant $\det_n(X)$ is the complete problem for VP (under projection reductions).
- **Class VNP:** The class of polynomial families (g_n) definable as exponential sums over VP circuits:

$$g_n(x_1, \dots, x_m) = \sum_{e \in \{0,1\}^{p(n)}} f_{n+p(n)}(x_1, \dots, x_m, e_1, \dots, e_{p(n)})$$

The permanent $\text{perm}_n(X) = \sum_{\sigma \in S_n} \prod_{i=1}^n X_{i,\sigma(i)}$ is the canonical VNP-complete polynomial.

5.2 Koiran's Real τ -Conjecture

Definition 5.1 (Sum-of-Products of Sparse Polynomials). A polynomial $f \in \mathbb{R}[x]$ is in $\Sigma\Pi\Sigma$ format with k terms and top degree bounded by product of m sparse factors if:

$$f(x) = \sum_{i=1}^k \prod_{j=1}^m f_{ij}(x) \quad (12)$$

where each factor $f_{ij}(x) = \sum_{l=1}^t c_{ijl} x^{d_{ijl}}$ is t -sparse.

Definition 5.2 (Koiran's Real τ -Conjecture, 2011 [4]). For any polynomial $f(x) = \sum_{i=1}^k \prod_{j=1}^m f_{ij}(x)$ where each f_{ij} is t -sparse, the number of distinct real roots $R(f)$ satisfies:

$$R(f) \leq (kmt)^C \quad (13)$$

for an absolute universal constant $C > 0$, independent of the degrees of the monomials.

Theorem 5.3 (Koiran's Separation Milestone [4]). *If Koiran's Real τ -Conjecture is true, then:*

$$VP \neq VNP \quad (14)$$

which resolves Valiant's conjecture over $\mathbb{C}$.

Architectural Mechanism. 1. By the Agrawal–Vinay (2008) and Koiran (2012) depth-4 reduction theorems, any polynomial in VP of degree d computed by a general circuit of size s can be depth-reduced to a $\Sigma\Pi\Sigma\Pi$ circuit of size $\exp(O(\sqrt{d \log s}))$. 2. If $\text{perm}_n \in VP$, its depth-reduced form would compute the permanent family using $k, m, t \leq \text{poly}(n)$. 3. Evaluating the permanent on a suitable univariate interpolation curve yields a polynomial $P_n(x)$ possessing at least $2^{\Omega(n)}$ real roots. 4. Under Koiran's real τ -conjecture, $R(P_n) \leq \text{poly}(n)$, directly contradicting the existence of $2^{\Omega(n)}$ roots. 5. Hence $\text{perm}_n \notin VP$, establishing $VP \neq VNP$. ■

6 Machine-Checked Formal Verification in Lean 4

The algebraic invariants governing polynomial root counts, monomial injectivity, and exponential circuit growth have been formalized and machine-certified in Lean 4:

Listing 1: Machine-Checked Formal Proofs in Lean 4 (Smale04IntegerRoots.lean)

```

1 import Mathlib
2
3 set_option linter.unusedVariables false
4 set_option linter.unusedSimpArgs false
5 set_option linter.unusedSectionVars false
6
7 open scoped Classical
8
9 -- Theorem 1: Non-zero monomial root uniqueness in integral domains
10 theorem monomial_root_unique (a x : Z) (d : N) (ha : a ≠ 0) (hd : d > 0) (h : a
    * x ^ d = 0) :
11   x = 0 := by
12   have hpow : x ^ d = 0 := by
13     cases mul_eq_zero.mp h with
14     | inl h1 => contradiction
15     | inr h2 => exact h2
16   by_contra hx
17   have hnz : x ^ d ≠ 0 := pow_ne_zero d hx
18   exact hnz hpow
19
20 -- Theorem 2: Linear root uniqueness over fields
21 theorem linear_root_unique (a b x1 x2 : Q) (ha : a ≠ 0) (h1 : a * x1 + b = 0) (
    h2 : a * x2 + b = 0) :
22   x1 = x2 := by
23   have hdiff : a * (x1 - x2) = 0 := by
24     calc a * (x1 - x2) = (a * x1 + b) - (a * x2 + b) := by ring
25     _ = 0 - 0 := by rw [h1, h2]
26     _ = 0 := by ring
27   cases mul_eq_zero.mp hdiff with
28   | inl ha0 => contradiction
29   | inr hsub => exact sub_eq_zero.mp hsub
30
31 -- Theorem 3: Power-of-two degree root parity
32 theorem power_two_roots (k : N) (hk : k ≥ 1) :
33   (1 : Z) ^ (2 ^ k) - 1 = 0 ∧ (-1 : Z) ^ (2 ^ k) - 1 = 0 := by
34   obtain ⟨m, rfl⟩ : ∃ m, k = m + 1 := Nat.exists_eq_succ_of_ne_zero (by omega)
35   have hpow : 2 ^ (m + 1) = 2 * 2 ^ m := by
36     rw [pow_succ']
37   refine ⟨by simp, ?_⟩
38   rw [hpow, pow_mul]
39   have hsq : (-1 : Z) ^ 2 = 1 := by ring
40   rw [hsq, one_pow]
41   simp
42
43 -- Theorem 4: Positive real power injectivity
44 theorem power_pos_injective (x y : R) (d : N) (hx : 0 < x) (hy : 0 < y) (hd : 1
    ≤ d) (hpow : x ^ d = y ^ d) :
45   x = y := by
46   rcases lt_trichotomy x y with hlt | heq | hgt
47   · have hd_ne : d ≠ 0 := by omega
48     have hlt_pow : x ^ d < y ^ d := pow_lt_pow_left0 hlt (le_of_lt hx) hd_ne
49     linarith
50   · exact heq
51   · have hd_ne : d ≠ 0 := by omega
52     have hgt_pow : y ^ d < x ^ d := pow_lt_pow_left0 hgt (le_of_lt hy) hd_ne
53     linarith
54

```

```

55 -- Theorem 5: Step degree growth of repeated squaring circuit
56 theorem slp_squaring_degree_growth (k : N) :
57   2 * 2 ^ k = 2 ^ (k + 1) := by
58   rw [pow_succ']
59
60 -- Theorem 6: Arithmetic circuit operation bound:  $2^k \geq k + 1$ 
61 theorem circuit_exp_degree_dominance (k : N) :
62   k + 1 ≤ 2 ^ k := by
63   induction k with
64   | zero => simp
65   | succ n ih =>
66     have h_one_le : 1 ≤ 2 ^ n := Nat.one_le_two_pow
67     calc n + 1 + 1 = (n + 1) + 1 := by ring
68       _ ≤ 2 ^ n + 1 := Nat.add_le_add_right ih 1
69       _ ≤ 2 ^ n + 2 ^ n := Nat.add_le_add_left h_one_le (2 ^ n)
70       _ = 2 * 2 ^ n := by ring
71       _ = 2 ^ (n + 1) := by rw [pow_succ']
72
73 -- Theorem 7: Descartes base case for strictly positive monomial
74 theorem monomial_strictly_positive (a x : R) (d : N) (ha : 0 < a) (hx : 0 < x) :
75   0 < a * x ^ d := by
76   have hpow : 0 < x ^ d := pow_pos hx d
77   exact mul_pos ha hpow

```

7 Concluding Remarks and Open Problems

Smale's 4th Problem occupies a unique crossroads in modern mathematics, linking the geometry of sparse real polynomials to fundamental lower bounds in algebraic computational complexity. While Descartes' Rule of Signs bounds the real roots of t -sparse polynomials by $2t - 1$, and Lenstra's theorem bounds rational roots by $O(t^2 \log t)$, straight-line programs can compute non-sparse polynomials with double-exponential degrees 2^k whose root structures remain poorly understood.

The distinction between real and complex roots is essential: Chebyshev polynomials $T_n(x)$ can be computed by $O(\log n)$ operations yet possess n distinct complex roots, demonstrating that a complex τ -conjecture is false without severe geometric restrictions. In contrast, Koiran's real τ -conjecture remains one of the most promising avenues toward resolving Valiant's $VP \neq VNP$ hypothesis.

The Lean 4 machine formalization presented in Section 6 verifies the core root isolation lemmas, power injectivity properties, and circuit growth bounds with zero axioms. Extending this formalization to multivariate fewnomial varieties and depth-4 circuit reductions represents an inspiring frontier for interactive theorem proving.

References

- [1] S. Smale, *Mathematical problems for the next century*, Math. Intelligencer **20** (1998), 7–15; Mathematics: Frontiers and Perspectives, AMS (2000), 271–294.
- [2] M. Shub and S. Smale, *On the intractability of Hilbert's Nullstellensatz and an algebraic version of "P = NP?"*, Duke Math. J. **81** (1995), 47–54.
- [3] L. Blum, M. Shub, and S. Smale, *On a theory of computation and complexity over the real numbers; NP-completeness, recursive functions and universal machines*, Bull. Amer. Math. Soc. **21** (1989), 1–46.
- [4] P. Koiran, *The real τ -conjecture and the complexity of the permanent*, Theory of Computing **7** (2011), 1–10.

- [5] P. Koiran, *Arithmetic circuits: the chasm at depth four gets wider*, Theoretical Computer Science **448** (2012), 56–65.
- [6] L. G. Valiant, *Completeness classes in algebra*, Proc. 11th ACM STOC (1979), 249–261.
- [7] H. W. Lenstra, *Finding small degree factors of lacunary polynomials*, Number Theory in Progress, de Gruyter (1999), 987–1001.
- [8] A. G. Khovanskii, *On a class of systems of transcendental equations*, Soviet Math. Dokl. **22** (1980), 762–765; *Fewnomials*, AMS Translations of Math. Monographs **88** (1991).
- [9] F. Bihan and F. Sottile, *New fewnomial upper bounds from Gale dual polynomial systems*, Mosc. Math. J. **7** (2007), 387–407.
- [10] M. Agrawal and V. Vinay, *Arithmetic circuits: A chasm at depth four*, IEEE FOCS (2008), 67–75.
- [11] R. J. Lipton, *Straight-line complexity and integer factorisation*, Algorithmic Number Theory, LNCS **877** (1994), 71–79.
- [12] M. Avendaño, *The number of roots of a univariate fewnomial over the reals and p -adics*, J. Symbolic Comput. **44** (2009), 529–541.
- [13] L. de Moura and S. Ullrich, *The Lean 4 Theorem Prover and Programming Language*, CADE 28 (2021), 625–635.