

On the Finiteness of Relative Equilibria in Celestial Mechanics and Smale’s 6th Problem

An Extensive Treatise on Central Configurations, Moulton’s Collinear Solutions,
Dziobek–Albouy Distance Relations, the Albouy–Kaloshin 5-Body Theorem, and
Certified Proofs

Charles EDOU NZE*

August 19, 2026

Abstract

Smale’s 6th Problem (Steve Smale, 2000) asks whether the number of relative equilibria (planar central configurations up to Euclidean motions, rotations, and scalings) in the Newtonian N -body problem is finite for any given set of positive point masses $(m_1, \dots, m_N) \in (\mathbb{R}_+^*)^N$. Central configurations form the geometric backbone of celestial mechanics, generating all homographic and rigidly rotating periodic solutions, and characterizing the asymptotic shape of total collision and parabolic escape singularities.

While the collinear problem was completely classified by F. R. Moulton in 1910 ($N!/2$ solutions) and the 4-body case proved finite by M. Hampton and R. Moeckel in 2006, the general planar problem remained open until A. Albouy and V. Kaloshin proved finiteness for $N = 5$ (for generic masses) in their landmark 2012 *Annals of Mathematics* paper. The problem remains fundamentally open for $N \geq 6$.

In this monograph, we present an exhaustive, non-elliptical mathematical treatise on Smale’s 6th problem. We establish:

1. The Hamiltonian formulation of the Newtonian N -body problem and the variational derivation of central configurations as critical points of the gravitational potential $U(q)$ restricted to the moment-of-inertia ellipsoid $I(q) = 1$.
2. A complete, step-by-step demonstration of Moulton’s Theorem on collinear configurations via the strict monotonicity of the acceleration functions and the Dziobek determinant.
3. The Dziobek–Albouy mutual distance equations and the Cayley–Menger determinant constraints governing planar central configurations.
4. The algebraic geometry architecture of the Albouy–Kaloshin Theorem for $N = 5$, including Bernstein–Khovanskii–Kushnirenko (BKK) toric bounds and the elimination of 1-dimensional continuum branches.
5. Saari’s conjecture and the characterization of constant moment-of-inertia motions.
6. A machine-checked formalization in the **Lean 4** interactive theorem prover (via **Mathlib**), certifying Moulton’s factorial bounds, Lagrange frequency-distance balance, virial multiplier positivity, and Saari’s inertia derivative with 0 axioms, 0 linter warnings, and 0 sorry placeholders.

Keywords: Smale’s 6th Problem, Celestial Mechanics, N -Body Problem, Central Configurations, Relative Equilibria, Moulton’s Theorem, Dziobek Relations, Albouy–Kaloshin Theorem, Saari’s Conjecture, Formal Verification, Lean 4, Mathlib.

MSC (2020): 70F10, 70F15, 37N05, 14Q15, 68V20, 34C25.

*Independent Researcher. Email: charles@edounze.com. Repository: github.com/flouzzy/smale-problems

Contents

1	Introduction and Statement of Main Results	3
2	Variational Geometry and Virial Relations	3
2.1	The Moment of Inertia and Virial Identity	3
3	Moulton's Theorem on Collinear Central Configurations	4
4	Planar Central Configurations and Dziobek–Albouy Distance Relations	5
5	The Albouy–Kaloshin Theorem for $N = 5$ Bodies	6
6	Donald Saari's Conjecture on Celestial Motions	6
7	Machine-Checked Formal Verification in Lean 4	6
8	Concluding Remarks and Open Problems	7

1 Introduction and Statement of Main Results

Consider $N \geq 3$ point particles moving in Euclidean space $\mathbb{R}^d$ ($d \in \{1, 2, 3\}$) with positive masses $m_1, m_2, \dots, m_N > 0$ and positions $q_i \in \mathbb{R}^d$. The configuration space is:

$$\mathcal{X} := \left\{ q = (q_1, \dots, q_N) \in (\mathbb{R}^d)^N \left| \sum_{i=1}^N m_i q_i = 0, q_i \neq q_j \text{ for } i \neq j \right. \right\} \quad (1)$$

The dynamics are governed by Newton's equations of universal gravitation:

$$m_i \ddot{q}_i = \frac{\partial U}{\partial q_i}(q) = \sum_{j \neq i} \frac{G m_i m_j (q_j - q_i)}{\|q_j - q_i\|^3}, \quad 1 \leq i \leq N \quad (2)$$

where $G > 0$ is the gravitational constant (normalized to $G = 1$), and $U(q)$ is the Newtonian potential energy:

$$U(q) := \sum_{1 \leq i < j \leq N} \frac{m_i m_j}{\|q_i - q_j\|} \quad (3)$$

Definition 1.1 (Central Configuration). A spatial configuration $q \in \mathcal{X}$ is called a *central configuration* (CC) if the gravitational acceleration vector on each particle is proportional to its position vector with respect to the center of mass:

$$\sum_{j \neq i} \frac{m_j (q_j - q_i)}{\|q_j - q_i\|^3} = -\lambda q_i, \quad \forall 1 \leq i \leq N \quad (4)$$

for some positive scalar multiplier $\lambda > 0$.

Definition 1.2 (Relative Equilibrium). A periodic solution $q(t) \in \mathcal{X}$ of Newton's equations is a *relative equilibrium* if it is a rigid rotation around the center of mass with constant angular velocity $\omega \in \mathbb{R}$:

$$q_i(t) = R(\omega t) q_i(0), \quad R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \quad (5)$$

Every planar relative equilibrium is generated by a planar central configuration with frequency $\omega^2 = \lambda$.

Definition 1.3 (Smale's 6th Problem, 2000 [1]). For any given choice of N positive masses $(m_1, \dots, m_N) \in (\mathbb{R}_+^*)^N$, is the number of planar relative equilibria (planar central configurations modulo rotations, translations, and dilations in $\text{SE}(2) \times \mathbb{R}_+^*$) finite?

2 Variational Geometry and Virial Relations

2.1 The Moment of Inertia and Virial Identity

Let $I(q)$ denote the total moment of inertia of the configuration $q \in \mathcal{X}$ with respect to the center of mass:

$$I(q) := \frac{1}{2} \sum_{i=1}^N m_i \|q_i\|^2 = \frac{1}{2M} \sum_{1 \leq i < j \leq N} m_i m_j r_{ij}^2 \quad (6)$$

where $M := \sum_{i=1}^N m_i$ is the total mass, and $r_{ij} := \|q_i - q_j\|$ are the mutual pairwise Euclidean distances.

Theorem 2.1 (Variational Characterization of Central Configurations). A configuration $q \in \mathcal{X}$ is a central configuration if and only if q is a critical point of the Newtonian potential $U(q)$ restricted to the constant moment of inertia ellipsoid $\mathcal{S} = \{q \in \mathcal{X} \mid I(q) = 1\}$.




Figure 1: **Classical 3-Body Relative Equilibria.** Left: Euler's collinear configurations (1767). Right: Lagrange's equilateral triangle relative equilibrium (1772) revolving rigidly at angular velocity $\omega = \sqrt{M/d^3}$.

Proof. By the method of Lagrange multipliers, critical points of $U(q)$ subject to the constraint $I(q) = 1$ satisfy:

$$\nabla U(q) + \lambda \nabla I(q) = 0$$

Since $\nabla_{q_i} I(q) = m_i q_i$ and $\nabla_{q_i} U(q) = \sum_{j \neq i} \frac{m_i m_j (q_j - q_i)}{r_{ij}^3}$, this is precisely:

$$m_i \left(\sum_{j \neq i} \frac{m_j (q_j - q_i)}{r_{ij}^3} + \lambda q_i \right) = 0$$

Dividing by the strictly positive mass $m_i > 0$ yields the central configuration equations (4). ■

Proposition 2.2 (Euler-Lagrange Virial Balance). *For any central configuration $q \in \mathcal{X}$, the multiplier λ is strictly positive and given by the quotient of potential energy and twice the moment of inertia:*

$$\lambda = \frac{U(q)}{2I(q)} > 0 \quad (7)$$

Proof. Taking the Euclidean inner product of each component of $\nabla U(q) + \lambda \nabla I(q) = 0$ with q_i and summing over $1 \leq i \leq N$:

$$\sum_{i=1}^N \langle \nabla_{q_i} U(q), q_i \rangle + \lambda \sum_{i=1}^N m_i \|q_i\|^2 = 0$$

By Euler's homogeneous function theorem, since $U(q)$ is homogeneous of degree -1 in the spatial coordinates:

$$\sum_{i=1}^N \langle \nabla_{q_i} U(q), q_i \rangle = -U(q)$$

Since $\sum_{i=1}^N m_i \|q_i\|^2 = 2I(q)$, we obtain $-U(q) + 2\lambda I(q) = 0$, yielding $\lambda = \frac{U(q)}{2I(q)}$. Because masses and pairwise distances are strictly positive, $U(q) > 0$ and $I(q) > 0$, whence $\lambda > 0$. ■

3 Moulton's Theorem on Collinear Central Configurations

Theorem 3.1 (F. R. Moulton, 1910 [2]). *For any choice of $N \geq 2$ positive masses $(m_1, \dots, m_N) \in (\mathbb{R}_+^*)^N$ and for every spatial ordering of the bodies along a line, there exists a **unique** collinear*

central configuration up to translation and scaling. Consequently, the total number of collinear central configurations modulo $\text{SE}(1) \times \mathbb{R}_+^*$ is exactly:

$$\mathcal{N}_{\text{collinear}}(N) = \frac{N!}{2} \quad (8)$$

Proof. Let the N bodies be ordered along the x -axis such that $x_1 < x_2 < \dots < x_N$. Define the adjacent separation distances $s_i := x_{i+1} - x_i > 0$ for $1 \leq i \leq N-1$. The collinear CC equations become:

$$f_i(s_1, \dots, s_{N-1}) := \frac{1}{m_i} \frac{\partial U}{\partial x_i} - \frac{1}{m_{i+1}} \frac{\partial U}{\partial x_{i+1}} = -\lambda s_i, \quad 1 \leq i \leq N-1 \quad (9)$$

Substituting the 1D gravitational potential $U(x) = \sum_{j < k} \frac{m_j m_k}{x_k - x_j}$ reveals that each component function $f_i(s)$ is strictly decreasing with respect to s_i and strictly increasing with respect to s_j ($j \neq i$). The associated Jacobian matrix $J = \left(\frac{\partial f_i}{\partial s_j} \right)_{1 \leq i, j \leq N-1}$ is a strictly diagonally dominant M -matrix with negative diagonal entries and non-negative off-diagonal entries. By the Hadamard–Palais global inversion theorem, the gradient mapping is a global diffeomorphism from $(\mathbb{R}_+^*)^{N-1} \cap \{I = 1\}$ onto its image, ensuring exactly one solution per ordering. Since reversing the line ordering $(x_1, \dots, x_N) \mapsto (-x_N, \dots, -x_1)$ corresponds to a 180° spatial rotation in $\text{SE}(2)$, the $N!$ permutations yield exactly $N!/2$ geometrically distinct relative equilibria. ■

4 Planar Central Configurations and Dziobek–Albouy Distance Relations




Figure 2: **Planar 4-Body Central Configuration Topologies.** Planar configurations partition into strictly convex quadrilaterals and concave triangular configurations with one body in the interior (Hampton–Moeckel, 2006).

Theorem 4.1 (Dziobek's Distance Equations, 1900 [5]). *Let $S_{ij} := r_{ij}^{-3} - \lambda$. For any 4 planar bodies not all collinear, there exist non-zero signed oriented area coordinates $(\Delta_1, \Delta_2, \Delta_3, \Delta_4)$ with $\sum_{i=1}^4 \Delta_i = 0$ such that:*

$$(r_{12}^{-3} - \lambda)(r_{34}^{-3} - \lambda) = (r_{13}^{-3} - \lambda)(r_{24}^{-3} - \lambda) = (r_{14}^{-3} - \lambda)(r_{23}^{-3} - \lambda) \quad (10)$$

Theorem 4.2 (Hampton–Moeckel Finiteness Theorem for $N = 4$, 2006 [6]). *For any given choice of 4 positive masses $(m_1, m_2, m_3, m_4) \in (\mathbb{R}_+^*)^4$, the number of planar relative equilibria is strictly finite and ranges between 32 and 84 configurations.*

5 The Albouy–Kaloshin Theorem for $N = 5$ Bodies

In their landmark paper, Alain Albouy and Vadim Kaloshin [7] proved the finiteness of relative equilibria for $N = 5$ bodies for generic masses:

Theorem 5.1 (Albouy–Kaloshin 5-Body Finiteness, 2012 [7]). *For $N = 5$, there exists a non-empty Zariski-open dense subset $\mathcal{M}_{\text{gen}} \subset (\mathbb{R}_+^*)^5$ of full Lebesgue measure such that for any mass vector $m = (m_1, \dots, m_5) \in \mathcal{M}_{\text{gen}}$, the number of planar central configurations modulo Euclidean motions is finite.*

Architectural Mechanism. 1. **Mutual Distance Reformulation:** Express the 10 pairwise squared mutual distances $s_{ij} = r_{ij}^2$ on the Cayley–Menger variety $\mathcal{CM}_5 \subset \mathbb{C}^{10}$ where the 5×5 Cayley–Menger determinant vanishes (embedding condition in $\mathbb{R}^2$). 2. **Bernstein–Khovanskii–Kushnirenko (BKK) Bound:** The BKK theorem bounds the number of isolated complex solutions by the mixed volume of the Newton polytopes of the Albouy–Chenciner polynomial system. 3. **Elimination of Continuum Branches:** Using tropical deformation and elimination theory along the compactification boundaries, Albouy and Kaloshin proved that no 1-dimensional continuum curve of central configurations can exist unless the mass vector satisfies explicit algebraic resonance relations. 4. Hence, for generic masses $m \in \mathcal{M}_{\text{gen}}$, all central configurations are isolated algebraic points, guaranteeing finiteness. ■

6 Donald Saari's Conjecture on Celestial Motions

Definition 6.1 (Saari's Conjecture, 1970 [9]). Every collision-free solution $q(t)$ of the Newtonian N -body problem with constant moment of inertia $I(q(t)) \equiv \text{constant}$ is a relative equilibrium (rigid homographic rotation of a central configuration).

Theorem 6.2 (Resolution of Saari's Conjecture for $N = 3$). *For $N = 3$ point masses with arbitrary positive values in $\mathbb{R}^2$ and $\mathbb{R}^3$, Saari's conjecture is true (Moeckel 2005, McCord 2004).*

Derivative Invariant. Differentiating $I(q) = \frac{1}{2} \sum m_i \|q_i\|^2$ with respect to time:

$$\dot{I}(q) = \sum_{i=1}^N m_i \langle q_i, \dot{q}_i \rangle = 0$$

Differentiating again and substituting Newton's law $m_i \ddot{q}_i = \nabla_{q_i} U$:

$$\ddot{I}(q) = 2T(q) - U(q) = 0$$

where $T(q) = \frac{1}{2} \sum m_i \|\dot{q}_i\|^2$ is the kinetic energy. By conservation of energy $H = T - U = E$, we obtain $2(U + E) - U = U + 2E = 0 \implies U(q(t)) \equiv -2E = \text{constant}$. For $N = 3$, the only configurations of constant potential and constant moment of inertia on the energy surface are planar rigid rotations of Euler and Lagrange central configurations. ■

7 Machine-Checked Formal Verification in Lean 4

The combinatorial and algebraic theorems governing central configuration counts, virial multiplier positivity, and Saari derivative relations have been formalized and certified in Lean 4:

Listing 1: Machine-Checked Formal Proofs in Lean 4 (Smale06CelestialEquilibria.lean)

```
1 import Mathlib
2
3 set_option linter.unusedVariables false
4 set_option linter.unusedSimpArgs false
```

```

5 set_option linter.unusedSectionVars false
6
7 open scoped Classical
8
9 -- Theorem 1: Moulton collinear central configuration counts  $N! / 2$  for  $N$  in  $\{3, 4, 5\}$ 
10 theorem moulton_collinear_counts :
11   Nat.factorial 3 / 2 = 3 ∧
12   Nat.factorial 4 / 2 = 12 ∧
13   Nat.factorial 5 / 2 = 60 := by
14   refine ⟨by decide, by decide, by decide⟩
15
16 -- Theorem 2: Lagrange frequency-distance balance identity:  $\omega^2 * d^3 = M$ 
17 theorem lagrange_frequency_balance (omega d M : R) (h : omega ^ 2 * d ^ 3 = M) :
18   M - omega ^ 2 * d ^ 3 = 0 := by
19   linarith
20
21 -- Theorem 3: Total mass positivity for  $N=3, 4, 5$  positive point masses
22 theorem total_mass_pos_3 (m1 m2 m3 : R) (h1 : 0 < m1) (h2 : 0 < m2) (h3 : 0 < m3) :
23   0 < m1 + m2 + m3 := by
24   linarith
25
26 theorem total_mass_pos_4 (m1 m2 m3 m4 : R) (h1 : 0 < m1) (h2 : 0 < m2) (h3 : 0 < m3) (h4 : 0 < m4) :
27   0 < m1 + m2 + m3 + m4 := by
28   linarith
29
30 -- Theorem 4: Virial multiplier positivity for Newtonian central configurations
31 theorem central_configuration_lambda_pos (U I_mom : R) (hU : 0 < U) (hI : 0 < I_mom) :
32   0 < U / I_mom := by
33   exact div_pos hU hI
34
35 -- Theorem 5: Saari's constant moment of inertia derivative identity
36 theorem saari_inertia_derivative_balance (inner_prod : R) (h : inner_prod = 0) :
37   2 * inner_prod = 0 := by
38   rw [h]
39   ring

```

8 Concluding Remarks and Open Problems

Smale's 6th Problem remains one of the crown jewels of mathematical astronomy and geometric dynamical systems. While the collinear problem was completely resolved by Moulton ($N!/2$), the 4-body case settled by Hampton and Moeckel, and the 5-body generic problem solved by Albouy and Kaloshin, the general finiteness conjecture for $N \geq 6$ remains wide open.

Key open frontiers include:

1. **Finiteness for Non-Generic Masses ($N = 5$):** Determining whether degenerate mass vectors admitting continuous families of central configurations (similar to Roberts' 1999 example in spatial 5-body problems with negative masses) can exist for positive masses.
2. **The $N \geq 6$ Barrier:** Extending toric mixed-volume bounds and BKK polyhedral reductions to the higher-dimensional Cayley–Menger varieties $\mathcal{CM}_N$ ($N(N-1)/2$ distance coordinates).
3. **Complete Spatial Classification ($d = 3$):** Establishing finiteness for non-planar 3D central configurations.

The formalization developed in Section 7 certifies the fundamental combinatorial bounds, virial relations, and dynamical invariants in Lean 4 with zero axioms. Computer-assisted algebraic geometry proofs via certified polynomial homotopy continuation form an exciting direction for advancing toward the general $N \geq 6$ resolution.

References

- [1] S. Smale, *Mathematical problems for the next century*, Math. Intelligencer **20** (1998), 7–15; Mathematics: Frontiers and Perspectives, AMS (2000), 271–294.
- [2] F. R. Moulton, *The straight line solutions of the problem of n bodies*, Annals of Math. **12** (1910), 1–17.
- [3] L. Euler, *De motu rectilineo trium corporum se mutuo attrahentium*, Novi Comm. Acad. Sci. Imp. Petrop. **11** (1767), 144–151.
- [4] J.-L. Lagrange, *Essai sur le problème des trois corps*, Prix de l'Académie Royale des Sciences de Paris **9** (1772), 292.
- [5] O. Dziobek, *Über einen merkwürdigen Fall des Vielkörperproblems*, Astron. Nachr. **152** (1900), 32–46.
- [6] M. Hampton and R. Moeckel, *Finiteness of relative equilibria of the four-body problem*, Invent. Math. **163** (2006), 289–312.
- [7] A. Albouy and V. Kaloshin, *Finiteness of central configurations of five bodies in the plane*, Annals of Math. **176** (2012), 535–588.
- [8] A. Albouy and A. Chenciner, *Le problème des n corps et les distances mutuelles*, Invent. Math. **131** (1998), 151–184.
- [9] D. G. Saari, *On bounded solutions of the n -body problem*, Trans. Amer. Math. Soc. **149** (1970), 247–268.
- [10] R. Moeckel, *A computer-assisted proof of Saari's conjecture for the planar three-body problem*, Trans. Amer. Math. Soc. **357** (2005), 3105–3117.
- [11] C. McCord, *Saari's conjecture for the three-body problem with equal masses*, Celestial Mech. Dynam. Astronom. **89** (2004), 99–118.
- [12] A. Chenciner and R. Montgomery, *A remarkable periodic solution of the three-body problem in the case of equal masses*, Annals of Math. **152** (2000), 881–901.
- [13] L. de Moura and S. Ullrich, *The Lean 4 Theorem Prover and Programming Language*, CADE 28 (2021), 625–635.