

# Approach to Internal Medicine

A Resource Book for Clinical  
Practice

David Hui · Alexander A. Leung ·  
Christopher Ma *Editors*

*Fifth Edition*



Springer

*Editors*

David Hui  
The University of Texas MD  
Anderson Cancer Center  
Houston, TX  
USA

Alexander A. Leung  
University of Calgary  
Calgary, AB  
Canada

Christopher Ma  
University of Calgary  
Calgary, AB  
Canada

ISBN 978-3-030-72979-0      ISBN 978-3-030-72980-6 (eBook)  
<https://doi.org/10.1007/978-3-030-72980-6>

© The Editor(s) and The Author(s), under exclusive license to Springer Nature  
Switzerland AG 2022

This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

**Acute Dyspnea**     2015 ACCP/CTS Guideline Prevention of Acute Exacerbations of COPD

**DIFFERENTIAL DIAGNOSIS**

**RESPIRATORY**

- **AIRWAY**—COPD exacerbation, asthma exacerbation, acute bronchitis, infectious exacerbation of bronchiectasis, foreign body obstruction
- **PARENCHYMA**—pneumonia, cryptogenic organizing pneumonia (COP), ARDS, acute exacerbation of interstitial lung disease
- **VASCULAR**—pulmonary embolism, pulmonary hypertension
- **PLEURAL**—pneumothorax, pleural effusion

**CARDIAC**

- **MYOCARDIAL**—HF exacerbation, myocardial infarction
- **VALVULAR**—aortic stenosis, acute aortic regurgitation, mitral stenosis, endocarditis
- **PERICARDIAL**—pericardial effusion, tamponade

**SYSTEMIC**—sepsis, metabolic acidosis, anemia, cachexia

**OTHERS**—neuromuscular, psychogenic, anxiety

**PATHOPHYSIOLOGY**

**PRECIPITANTS OF COPD EXACERBATION**—infections, lifestyle/environmental (10% [cigarette smoke, dust, pollutants, cold air]), non-adherence to medications, pulmonary embolism, pulmonary edema, pneumothorax, progression of COPD

**CLINICAL FEATURES**

**RATIONAL CLINICAL EXAMINATION SERIES: DOES THE CLINICAL EXAMINATION PREDICT AIRFLOW LIMITATION?**

	LR+	LR–
<b>History</b>		
<i>Smoking &gt;40 pack-years</i>	12	0.63
<i>Smoking ever</i>	1.8	0.16
<i>Sputum &gt;1/4 cup</i>	4	0.84
<i>Chronic bronchitis symptoms</i>	3	0.78
<i>Wheezing</i>	3.8	0.66
<i>Any exertional dyspnea</i>	2.2	0.83
<i>Coughing</i>	1.8	0.69
<i>Any dyspnea</i>	1.2	0.55

**CLINICAL FEATURES (CONT'D)**

	LR+	LR–
<b>Physical</b>		
<i>Barrel chest</i>	10	0.90
<i>Decreased cardiac dullness</i>	10	0.88
<i>Match test</i>	7.1	0.43
<i>Rhonchi</i>	5.9	0.95
<i>Hyperresonance</i>	4.8	0.73
<i>FEV1 &gt;9 s</i>	4.8	–
<i>FEV1 6–9 s</i>	2.7	–
<i>FEV1 &lt;6 s</i>	0.45	–
<i>Subxiphoid cardiac apical impulse</i>	4.6	0.94
<i>Wheezing</i>	4.4	0.88
<i>Maximum laryngeal height ≤ 4 cm</i>	4.2	0.70
<i>Pulsus paradoxus (&gt;15 mmHg)</i>	3.7	0.62
<i>Decreased breath sounds</i>	2.6	0.66
<i>Accessory muscle use</i>	–	0.70
<b>Clinical Judgement</b>		
<i>Overall Clinical Prediction of Moderate-Severe Disease</i>	5.6	–
<i>Overall Clinical Prediction of Mild Disease</i>	2.3	–

**APPROACH**—“No single item or combination of items from the clinical examination rules out airflow limitation.” The best findings associated with increased likelihood of airflow limitation are objective wheezing, FEV1 >9 s, positive match test, barrel chest, hyperresonance, and subxiphoid cardiac impulse. “Three findings predict the likelihood of airflow limitation in men: years of cigarette smoking, subjective wheezing and either objective wheezing or peak expiratory flow rate.”

**Holleman et al. JAMA 1995;273(4)**

**UPDATE**—multivariate ‘Rule In’ Obstructive Disease Model (history of obstructive airways disease, smoking >40 pack-years, age ≥45, and laryngeal height ≤4 cm) has posterior odds of disease of 220. Multivariate ‘Rule Out’ Obstructive Disease Model (smoking <30 years, no wheezing symptoms, and no auscultated wheezing) has posterior odds of disease of 0.02

**Simel et al. The Rational Clinical Examination. McGraw-Hill, 2009**

## INVESTIGATIONS

### BASIC

- **LABS**—CBC, lytes, urea, Cr, troponin/Ck, Ca, Mg, PO<sub>4</sub>
- **MICROBIOLOGY**—sputum Gram stain/AFB/C&S/fungal, nasopharyngeal swab for viral studies
- **IMAGING**—CXR
- **ECG**—left atrial enlargement, atrial fibrillation, sinus tachycardia
- **SPIROMETRY/PFT**—FEV1/FVC <0.7, may be partially reversible. Severity based on FEV1
- **ABG**—if acute respiratory distress

### SPECIAL

- **BNP**—if suspect HF
- **D-DIMER, CT CHEST**—if suspect PE
- **ECHOCARDIOGRAM**

## DIAGNOSTIC & PROGNOSTIC ISSUES

**DIAGNOSIS OF COPD**—should be considered in patients at risk of developing disease; smoking history is most important risk factor; increased risk in patients with past history of asthma or severe childhood respiratory disease, exposed to passive smoke or biomass fuel; spirometry is essential for diagnosis (fixed post-bronchodilator FEV1/FVC ratio <0.70 or less than the lower limit of normal)

**GOLD CLASSIFICATION FOR COPD**—all have FEV1/FVC <0.7. Severity of airflow limitation based on post-bronchodilator FEV1

- **STAGE 1** (mild)—FEV1 ≥ 80% predicted
- **STAGE 2** (moderate)—FEV1 50–79% predicted
- **STAGE 3** (severe)—FEV1 30–49% predicted
- **STAGE 4** (very severe)—FEV1 <30% predicted

### MODIFIED MEDICAL RESEARCH COUNCIL (mMRC) DYSPNEA SCALE

- **0**—no breathlessness except on strenuous exercise
- **1**—short of breath when hurrying or walking up a slight hill
- **2**—walks slower than people of same age on level because of breathlessness or has to stop when walking at own pace
- **3**—stops for breath after walking 100m or after a few minutes
- **4**—too dyspneic to leave house; breathless when dressing

**COPD ASSESSMENT TEST**—a validated patient-reported outcome that consists of 8 items (cough, phlegm, chest tightness, shortness of breath on exertion, activity level, confidence to leave home, sleep, energy), each rated using a 6-point numeric rating scale from 0 to 5, with a higher total score indicating greater symptom burden

**GOLD ABCD GRADING**—assessment for initiation of COPD therapy

## DIAGNOSTIC & PROGNOSTIC ISSUES (CONT'D)

Exacerbations/ Hospitalizations	Assess symptoms	
	mMRC 0–1; CAT <10	mMRC ≥2; CAT ≥10
0–1 exacerbations without hospitalization	Gold A	Gold B
≥2 exacerbations or ≥1 hospitalization	Gold C	Gold D

**Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2020 Report**

**PROGNOSIS OF PATIENTS WITH ACUTE EXACERBATION OF COPD**—in-hospital mortality 5–10%

### BODE INDEX

- **BMI**—0 points = >21, 1 point = ≤21
- **OBSTRUCTION** (post-bronchodilator FEV1)—0 points = ≥65% predicted, 1 point = 50–64%, 2 points = 36–49%, 3 points = ≤35%
- **DISTANCE WALKED IN 6 MIN**—0 points = ≥350 m, 1 point = 250–349 m, 2 points = 150–249 m, 3 points = ≤149 m
- **EXERCISE MMRC DYSPNEA SCALE**—0 points = 0–1, 1 point = 2, 2 points = 3, 3 points = 4
- **SCORING**—total BODE score calculated as sum of all points. Relative risk for death (any cause) is increased by 34% per one-point increase in BODE score. Relative risk for death (from respiratory failure, pneumonia, or pulmonary embolism) is increased by 62% per one-point increase in BODE score

Celli et al. *NEJM* 2004;350:(10)

## ACUTE MANAGEMENT

**ABC**—O<sub>2</sub> to keep sat >90%, or 88–92% if CO<sub>2</sub> retainer, IV

**BRONCHODILATORS**—*salbutamol* 100 µg MDI 2 puffs q4h ATC + q1h PRN and *ipratropium* 20 µg MDI 2 puffs q4h ATC

**STEROIDS**—*prednisone* 40–60 mg PO daily × 5–14 days (tapering dose not always necessary) or *methylprednisolone* 60–125 mg IV q6–12 h (inpatient)

**ANTIBIOTICS**—give if any two of the following criteria are met: ↑ sputum purulence, ↑ dyspnea or ↑ sputum volume. Other considerations include the need for non-invasive mechanical ventilation and “at risk” for poor outcomes (substantial comorbidities, severe COPD, frequent exacerbations >3/year, recent antibiotics within 3 months); choices depend on clinical circumstance (*levofloxacin* 500 mg PO daily × 5–7 days [or 750 mg PO daily × 5

**ACUTE MANAGEMENT (CONT'D)**

days if no renal disease], *doxycycline* 100 mg PO BID  $\times$  5–7 days, *amoxicillin* 500 mg PO BID  $\times$  5–7 days, *cefuroxime* 250–500 mg PO BID  $\times$  5–7 days, *ceftriaxone* 1 g IV  $\times$  5–7 days, or *azithromycin* 500 mg PO  $\times$  1 day then 250 mg PO daily  $\times$  4 days)

**RESPIRATORY SUPPORT**—non-invasive ventilation, intubation and mechanical ventilation

**OTHERS**—DVT prophylaxis (*unfractionated heparin* 5000 U SC q8–12 h, *enoxaparin* 40 mg SC q24h, *dalteparin* 5000 U SC q24h, *tinzaparin* 75 IU/kg SC q24h), physiotherapy

**LONG-TERM MANAGEMENT**

**EDUCATION**—**smoking cessation** (see p. 490). Disease-specific self-management program. **Inhaler technique** education and review

**VACCINATIONS**—influenza vaccine annually and pneumococcal vaccine booster every 5 years

**REHABILITATION**—**education** and **exercise training** (increases quality of life and exercise tolerance); **pulmonary rehabilitation** associated with ↓ risk of recurrent exacerbation in patients with moderate to very severe COPD and recent AECOPD (<4 weeks)

**LONG-TERM MANAGEMENT (CONT'D)**

**LONG-TERM OXYGEN THERAPY**—if chronic hypoxemia

**INITIAL PHARMACOLOGIC THERAPY**—based on symptoms and risk of exacerbations

- **GOLD A (MINIMAL SYMPTOMS, LOW RISK OF EXACERBATION)**—short-acting bronchodilator with SABA (short-acting beta agonist) and/or SAMA (short-acting muscarinic antagonist)
- **GOLD B (MORE SYMPTOMS, LOW RISK OF EXACERBATION)**—regular treatment with long-acting bronchodilator (LAMA or LABA) plus SABA for symptom relief as needed
- **GOLD C (MINIMAL SYMPTOMS, HIGH RISK OF EXACERBATION)**—regular treatment with LAMA plus SABA for symptom relief as needed
- **GOLD D (MORE SYMPTOMS, HIGH RISK OF EXACERBATION)**—regular treatment with LAMA or combination LABA and LAMA if severe breathlessness (CAT >20); if features of asthma/COPD overlap syndrome, ICS/LABA combination may be preferred; plus SABA for symptom relief as needed

**SUBSEQUENT PHARMACOLOGIC THERAPY**

Current therapy	If persistent dyspnea or high COPD impact (i.e. mMRC $\geq$ 2 or CAT $\geq$ 10) with no exacerbations	If $\geq$ 1 exacerbations in past year $\pm$ persistent dyspnea or high COPD impact
SABA or SABA-SAMA PRN	Add LAMA or LABA	Add LAMA
LAMA or LABA monotherapy	Change to LAMA/LABA	LAMA/LABA if peripheral eosinophils normal LABA/ICS if LAMA contraindicated and 1 exacerbation in past year with peripheral eosinophils >300/ $\mu$ L or $\geq$ 2 exacerbations/1 hospitalization in past year with peripheral eosinophils $\geq$ 100/ $\mu$ L
LABA/ICS	LAMA/LABA/ICS or LAMA/LABA if no response to ICS or adverse effects from ICS	LAMA/LABA/ICS if prior indication for ICS LAMA/LABA if lack of response to ICS or adverse effects from ICS
LAMA/LABA	Trial of different LAMA/LABA or alternate delivery system Consider low dose theophylline, repeat pulmonary rehabilitation	LAMA/LABA/ICS or Continue LAMA/LABA and add phosphodiesterase-4 inhibitor (roflumilast 500 $\mu$ g PO daily) <sup>a</sup> or azithromycin <sup>b</sup>
LAMA/LABA/ICS	Continue LAMA/LABA/ICS Consider low dose theophylline (400 mg PO daily $\times$ 3 days, then 400–600 mg PO daily, therapeutic level 10–20 $\mu$ g/mL), repeat pulmonary rehabilitation Consider stopping ICS if lack of response or adverse effect to ICS	Add roflumilast <sup>a</sup> or azithromycin <sup>b</sup> Stop ICS if lack of response or adverse effect

<sup>a</sup>roflumilast for patients with FEV<sub>1</sub> <50% predicted and at least 1 hospitalization in past year

<sup>b</sup>azithromycin preventive therapy is more effective in patients who are not current smokers; consider development of resistant organisms such as non-Tuberculous mycobacterium

**LONG-TERM MANAGEMENT (CONT'D)**

**INVASIVE INTERVENTIONS**—if symptoms still persistent and/or decline in function, consider lung volume reduction procedures (surgery, endobronchial valves), lung transplantation

**TREATMENT ISSUES**

**COMMON INHALED MEDICATIONS** (DPI=dry powder inhaler; SMI=soft mist inhaler)

- **LAMA**—*tiotropium* DPI 18 mcg daily or SMI 2 inhalations of 2.5 mcg once daily, *glycopyrronium* DPI 50 mcg capsule once daily, *umeclidinium* DPI 62.5 mcg inhalation daily, *aclidinium* DPI 400 mcg BID
- **LABA**—*formoterol* DPI 12–24 mcg BID, *indacaterol* DPI 75 mcg daily, *salmeterol* DPI 50 mcg BID
- **LAMA/LABA COMBINATIONS**—*glycopyrrolate* 50 mcg/*indacaterol* 110 mcg 1 INH daily, *tiotropium* 2.5 mcg/*olodaterol* 2.5 mcg 2 INH daily, *umeclidinium* 62.5 mcg/*vilanterol* 25 mcg 1 INH daily
- **LAMA/LABA/ICS**—*fluticasone furoate* 100 mcg/*umeclidinium* 62.5 mcg/*vilanterol* 25 mcg 1 INH daily

**FACTORS FOR IMPENDING INTUBATION**—cardiac or respiratory failure, hemodynamic instability, markedly elevated respiratory rate ( $>35/\text{min}$ ), fatigue and labored respiration, use of accessory muscles, worsening hypercapnia, acidosis (especially lactic), stridor (impending upper airway obstruction), agonal breathing (impending respiratory arrest)

**LIFE-PROLONGING MEASURES FOR COPD**—smoking cessation, supplemental  $\text{O}_2$ , lung transplant

**INDICATIONS FOR SUPPLEMENTAL HOME  $\text{O}_2$** —ABG done at room air.  $\text{PaO}_2 <55$  mmHg alone or  $\text{PaO}_2 <60$  mmHg in the presence of bilateral ankle edema, cor pulmonale, or hematocrit  $>56\%$

**SPECIFIC ENTITIES** **$\alpha 1$ -ANTITRYPSIN DEFICIENCY**

- **PATHOPHYSIOLOGY**—production of an abnormal protease inhibitor (homozygous ZZ) with impaired transport out of the liver. Serum level is only 10–15% of normal  $\rightarrow$  increased protease activity leads to emphysema and cirrhosis (10%)
- **DIAGNOSIS**— $\alpha 1$ -antitrypsin levels; targeted testing should be considered in patients with COPD diagnosed before 65 years of age or with a smoking history of  $<20$  pack years
- **TREATMENTS**—similar to COPD,  $\alpha 1$ -antitrypsin replacement

**SPECIFIC ENTITIES (CONT'D)****ASTHMA AND COPD OVERLAP SYNDROME (ACOS)**

- **DIAGNOSIS**—patients with clinical features of both asthma and COPD. Airflow limitation not fully reversible, FEV1/FVC ratio  $<0.7$  or  $<\text{LLN}$  and bronchodilator increase in FEV1  $>12\%$  and 400 mL; history of atopy or allergies; exposure to risk factors (i.e.  $>10$  pack year smoking or equivalent, indoor/outdoor air pollutant exposure)
- **TREATMENTS**—similar to COPD and asthma

**BRONCHIOLITIS OBLITERANS**

- **PATHOPHYSIOLOGY**—severe inflammation of bronchioles  $\rightarrow$  airflow obstruction. Very different from bronchiolitis obliterans organizing pneumonia (BOOP)/cryptogenic organizing pneumonia (COP), a parenchymal lung disorder
- **CAUSES**—infection (viral, *Mycoplasma*), inflammatory (ulcerative colitis, rheumatoid arthritis), transplant (bone marrow, lung), toxic fumes, idiopathic
- **TREATMENTS**—bronchiolitis obliterans (with an organizing intraluminal exudate and proliferative granulation tissue polyp) is usually steroid responsive. Constrictive bronchiolitis (late, fibrotic, concentric) is not responsive to glucocorticoids

**BRONCHIECTASIS**

- **PATHOPHYSIOLOGY**—airway obstruction, destruction, altered immunity  $\rightarrow$   $\uparrow$  cellular and mediator inflammatory response  $\rightarrow$   $\uparrow$  elastase, sputum production  $\rightarrow$  recurrent infections  $\rightarrow$  vicious cycle  $\rightarrow$  permanent dilatation of bronchi. Major types of bronchiectasis include
  - **CYLINDRICAL OR TUBULAR BRONCHIECTASIS**—dilated airways alone, sometimes represents residual effect of pneumonia and may resolve
  - **VARICOSE BRONCHIECTASIS**—focal constrictive areas along the dilated airways
  - **SACULAR OR CYSTIC BRONCHIECTASIS**—most severe form. Progressive dilatation of the airways, resulting in large cysts or saccules
- **CAUSES**
  - **FOCAL**—broncholith, post-infectious, tumor, extrinsic lymph node compression, post-lobar resection, recurrent aspiration
  - **DIFFUSE**
    - **POST-INFECTIOUS**—bacterial (*Pseudomonas*, *Haemophilus*), mycobacterium, fungal, viral (adenovirus, measles, influenza, HIV)

**SPECIFIC ENTITIES (CONT'D)**

- **IMMUNODEFICIENCY**—cancer, chemotherapy, hypogammaglobulinemia, immunosuppression, sequelae of toxic inhalation or aspiration of foreign body
- **INTERSTITIAL LUNG DISEASE**—traction bronchiectasis
- **INFLAMMATORY**—RA, SLE, Sjögren syndrome, relapsing polychondritis, IBD
- **INHERITED**— $\alpha$ 1-antitrypsin deficiency, cystic fibrosis, primary ciliary dyskinesia (Kartagener syndrome, Young syndrome), tracheobronchomegaly (Mounier-Kuhn syndrome), cartilage deficiency (Williams-Campbell syndrome), Marfan syndrome

**SPECIFIC ENTITIES (CONT'D)**

- **DIAGNOSIS**—high-resolution CT chest (signet ring sign), PFT (obstruction  $\pm$  reversibility)
- **TREATMENTS**—exercises, chest physiotherapy, and bronchodilators similar to COPD; however, if reversible, inhaled corticosteroids should be given early. Ensure adequate systemic hydration. Effective treatment of exacerbations

**Related Topics**

Cryptogenic Organizing Pneumonia (p. 21)  
 Pulmonary Function Tests (p. 25)  
 Smoking (p. 490)