On The Gravitational Nature Of Time

Andrew Mueller[1*]

[1*]Independent Researcher, Milwaukee, 53219, WI, USA.

Corresponding author(s). E-mail(s): AndrewMueller414@gmail.com;

Abstract

We explore a modified set of coordinate transformations that describe time, cosmic inflation and gravitational acceleration as the same process as observed from different reference frames through an unwrapped Kaluza-Klein like geometry. These transformations satisfy existing experimental validations of special relativity while providing a set of 4 equations yielding our peculiar velocity, a resolution to the SNe/CMB velocity discrepancy, an orbital velocity equation that matches observation with only visible mass, and a direct physical mechanism to the equivalence principle.

Keywords: gravitation, time, relativity, inflation

1 Introduction

Albert Einstein was insistent on a static Universe during the period in which he wrote his specialized theory of relativity[1–4]. What is proposed here is largely a re-examination of this geometry under the assumption that the $\mathbb{R}^3$ space between two arbitrary points dilates as time progresses.

This dilation is then bound to γ, giving a velocity dependent function for $\vec{g}$ and a magnitude that extends into a 4th spatial coordinate, mathematically similar to an unwrapped Kaluza-Klein (KK) dimension[5, 6]. Through these coordinate transformations time, cosmic inflation and gravitational acceleration become the same process as observed from different reference frames in a space-time where the 4th KK dimension acts as a concurrent density axis.

This geometry gives an entirely different model of concurrency and synchronicity as a classical notion of simultaneity is maintained. While maintaining classical simultaneity resolves a long list of temporal paradoxes without the need for selective reference

frame preference, it produces a concurrent axis that describes the ultra-dense state of the early Universe being separated only by some Δ along this 4th axis.

1.1 The Lighthouse & The Clock-Tower

Let there be two overlapping coordinate systems, **S** and **U**. Let **S** and **U** represent static positions in time, with **U** representing a 'future' state from the perspective of an observer in **S** so that

$$v^{(S)} = (t, x_\mu) \qquad v^{(U)} = (t + \Delta t, x_\mu) \tag{1}$$

Let there also be two observers, A and B. Let A and B agree that B should travel at some relativistic velocity v' between two arbitrary points in space contained within **S** and **U**, p_1 and p_2. Let A remain at relative rest with respect to p_1 and at p_2 place a time keeping device large enough to be visible from p_1.

If A and B were to disagree upon the time at which B reaches p_2, this single, instantaneous event (B reaching p_2) would be required to happen more than once. Therefore, we posit that it is not possible for A and B to disagree upon the time at which B reaches p_2.

Instead, as the additional factor of γ has been demonstrated experimentally on a number of occasions[7–11], we should conclude that both A and B observe B reaching p_2 in $\frac{d}{v}\gamma$. This gives a set of coordinate transformations such that for each respective observer

$$\Delta t = \frac{\overbrace{d \cdot \gamma}^{B}}{v^{(U)}} = \underbrace{\frac{d}{v^{(S)}} = \frac{d}{v^{(U)}\left(\frac{1}{\gamma}\right)}}_{A} \tag{2}$$

This produces a model of relativistic dilation that loosely binds the additional γ factor in the $d = vt\gamma$ equivalence to cosmic inflation, where the additional velocity experienced by B results in a more rapid progression along this concurrent density axis leading to a further separation between any two points in the $\mathbb{R}^3$ set.

2 Theoretical Framework

Let us acknowledge the two primary properties of time:

1. A driving rate of change, as fundamental as the $\mathbb{R}^3$ dimensions.
2. A 4th displacement coordinate.

Throughout the following derivations, we presume that time has no physical significance, and acts only as a mathematical tool: $t = \frac{d}{v}$. In turn, we must attribute the 4th displacement coordinate to a position along the density axis that arises, now denoted y, and the driving rate of change to *motion*.

Likewise, we presume that in a co-moving frame of reference along the y axis, mechanical equations should limit at their Newtonian counterpart.

As we are removing time as a physically meaningful quantity, let us define

$$\tau = \frac{\dot{x}}{\widehat{v}} \tag{3}$$

where $\dot{x}$ indicates not a derivative with respect to time, but rather a fundamental driving rate of change. Let $\widehat{v} = v \cdot 1t$, or the velocity in units of distance.




Fig. 1 A plane-based geometric simplification of the concurrent density axis found by these coordinate transformations.

2.1 The Equivalence Principle

Here we examine these coordinate transformations in the context of the equivalence principle. While a well defined function remains elusive, a numerical intersection with the fine structure constant allows us to derive several important properties of an apparent electromagnetic function producing $\vec{g}$.

By solving γ for v, let us define $v_{(\gamma)}$ to be

$$v_{(\gamma)} = c\sqrt{1 - \frac{1}{\gamma^2}} \tag{4}$$

And δ to be

$$\delta_{(R)} = \frac{d}{dR}g = -2\frac{GM}{R^3} \tag{5}$$

If δ is induced by spatial dilation, we should presume that this dilation is radial (at least in a simplified single body system). We then find the magnitude of a single $\mathbb{R}^3$ axis by taking the inverse of the determinant of a diagonal 3×3 matrix:

$$\omega = \sqrt[3]{\frac{d}{dR}g} = \delta^{1/3} \tag{6}$$

If we investigate this quantity for Earth, we find:

$$\omega_{\oplus} = 0.014537\ldots \tag{7}$$

While this approximates 2α, it more closely gives

$$\omega_{\oplus} = \alpha\left(2-\alpha\right) \pm 0.0253\% \tag{8}$$

If we then expand equation 8 we find

$$\alpha\left(2-\alpha\right) = \sqrt[3]{\frac{d}{dR}g} \tag{9}$$

Let us then move R to the electromagnetic side of the equation algebraically. Integrating this function then gives

$$R^4\left(\frac{e^6\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{4h^3} + \frac{3\mu_0 e^{10}\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{64\epsilon_0 h^5} - \frac{3\mu_0^2 e^8}{16\epsilon_0^2 h^4} - \frac{\mu_0^3 e^{12}}{256\epsilon_0^3 h^6}\right) = 1.272\cdot 10^{21} \tag{10}$$

If we then take the radius independent magnitude, isolating the additional electromagnetic parameters we find

$$\frac{e^6\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{4h^3} + \frac{3\mu_0 e^{10}\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{64\epsilon_0 h^5} - \frac{3\mu_0^2 e^8}{16\epsilon_0^2 h^4} - \frac{\mu_0^3 e^{12}}{256\epsilon_0^3 h^6} \tag{11}$$

As γ represents 1 plus some small ds, let us add 1 to these additional parameters and substitute γ_{em} into equation 4. This gives

$$\gamma_{\text{em}} = 1 + \frac{e^6\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{4h^3} + \frac{3\mu_0 e^{10}\left(\frac{\mu_0}{\epsilon_0}\right)^{\frac{3}{2}}}{64\epsilon_0 h^5} - \frac{3\mu_0^2 e^8}{16\epsilon_0^2 h^4} - \frac{\mu_0^3 e^{12}}{256\epsilon_0^3 h^6} \tag{12}$$

$$v_{(\gamma_{\text{em}})} = 371,721\ \frac{m}{s} \tag{13}$$

3 Results

Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text.

4 This is an example for first level head—section head

Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text. Sample body text.




Fig. 2 The inverse of the determinant representing a change in volume, being equivalent to dy, plotted against mass density.

5 Equations

References

[1] Einstein, A.: On the electrodynamics of moving bodies. Annalen der Physik **17**(10), 891–921 (1905) https://doi.org/10.1002/andp.19053221004

[2] Einstein, A.: Kosmologische betrachtungen zur allgemeinen relativitätstheorie. Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 142–152 (1917)

[3] O'Raifeartaigh, C.: Albert einstein and the origins of modern cosmology. Physics Today (2017) https://doi.org/10.1063/PT.5.9085

[4] Nussbaumer, H.: Einstein's conversion from his static to an expanding universe. The European Physical Journal H **39**, 37–62 (2014) https://doi.org/10.1140/epjh/e2013-40037-6

[5] Kaluza, T.: Zum unitätsproblem der physik. Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften zu Berlin (Math. Phys.), 966–972 (1921)

[6] Klein, O.: Quantentheorie und fünfdimensionale relativitätstheorie. Zeitschrift für Physik **37**(12), 895–906 (1926) https://doi.org/10.1007/BF01397481




Fig. 3 A trigonometric derivation of a relativistic orbital velocity equation.

[7] Rossi, B., Hall, D.B.: Variation of the rate of decay of mesotrons with momentum. Physical Review **59**(3), 223–228 (1941) https://doi.org/10.1103/PhysRev.59.223

[8] Frisch, D.H., Smith, J.H.: Measurement of the relativistic time dilation using mu-mesons. American Journal of Physics **31**(5), 342–355 (1963) https://doi.org/10.1119/1.1969508

[9] Bailey, J., Borer, K., Combley, F., Drumm, H., Krienen, F., Lange, F., Picasso, E., Ruden, W., Farley, F.J.M., Field, J.H., Flegel, W., Hattersley, P.M.: Measurements of relativistic time dilatation for positive and negative muons in a circular orbit. Nature **268**, 301–305 (1977) https://doi.org/10.1038/268301a0

[10] Bailey, J., *et al.*: Final report on the cern muon storage ring including the anomalous magnetic moment and the electric dipole moment of the muon, and a direct test of relativistic time dilation. Nuclear Physics B **150**, 1–75 (1979) https://doi.org/10.1016/0550-3213(79)90292-X

[11] Botermann, B., Bing, D., Geppert, C., Gwinner, G., Hansch, T.W., Huber, G., Karpuk, S., Krieger, A., Kuhl, T., Nortershauser, W., Novotny, C., Reinhardt, S.,

Sanchez, R., Schwalm, D., Stohlker, T., Wolf, A., Saathoff, G.: Test of time dilation using stored li+ ions as clocks at relativistic speed. Physical Review Letters **113**(12), 120405 (2014) https://doi.org/10.1103/PhysRevLett.113.120405