

Category Theory 101

Mathematical Structures in Computer Science

Oisín Flynn-Connolly

September 24, 2025



Motivation for computer scientists

- ① **Convenient language:** (Higher) category theory is a very general organizational principle for mathematics. Sometimes one notices that the same construction appears in multiple contexts. Category theory allows you to formalize this via **universal constructions**. Most of modern mathematics¹ is written in the language of category theory.
- ② **Compositionality.** Sometimes one wants to break a program down into smaller parts that can each be analysed independently.
- ③ **Logic.** There is a correspondence with logic via the Curry–Howard correspondence. Take Henning’s course during master for more!

¹or at least algebra and topology since the 1960s

Motivation for computer scientists

- ① **Convenient language:** (Higher) category theory is a very general organizational principle for mathematics. Sometimes one notices that the same construction appears in multiple contexts. Category theory allows you to formalize this via **universal constructions**. Most of modern mathematics¹ is written in the language of category theory.
- ② **Compositionality.** Sometimes one wants to break a program down into smaller parts that can each be analysed independently.
- ③ **Logic.** There is a correspondence with logic via the Curry–Howard correspondence. Take Henning’s course during master for more!

¹or at least algebra and topology since the 1960s

Motivation for computer scientists

- ① **Convenient language:** (Higher) category theory is a very general organizational principle for mathematics. Sometimes one notices that the same construction appears in multiple contexts. Category theory allows you to formalize this via **universal constructions**. Most of modern mathematics¹ is written in the language of category theory.
- ② **Compositionality.** Sometimes one wants to break a program down into smaller parts that can each be analysed independently.
- ③ **Logic.** There is a correspondence with logic via the Curry–Howard correspondence. Take Henning’s course during master for more!
- ④ **Concrete applications**
 - ① **Functional programming:** Category theory underlies the theory and syntax of functional programming languages such as OCaml and Haskell.
 - ② **Linguistics:** Category theory has been used to study human language.
 - ③ **Processes in CS:** To analyse all manner of processes in CS; from optimization algorithms to attack trees.

¹or at least algebra and topology since the 1960s

Direct sum of vector spaces

Given two vector spaces you can construct their direct sum:

Definition

Given two vector spaces V, W the direct sum is defined as:

$$V \oplus W = \{(v, w) \mid v \in V, w \in W\}$$

To define the vector space structure

$$(v_1, w_1) + (v_2, w_2) = (v_1 + v_2, w_1 + w_2) \quad v_1, v_2 \in V, w_1, w_2 \in W$$

$$\lambda \cdot (v, w) = (\lambda v, \lambda w)$$

There are **canonical inclusions**

$$i_V : V \rightarrow V \oplus W, \quad v \mapsto (v, 0)$$

$$i_W : W \rightarrow V \oplus W \quad w \mapsto (0, w)$$

Disadvantages of this definition

- Rather ad hoc and messy. Need to define the vector space structure by hand.
- Hides the role of the canonical inclusions.
- Doesn't generalise well.

Universal constructions: Direct sum of vector spaces

You saw the direct product of vector spaces two weeks ago. But there is a different way to construct it:

Definition

Given two vector spaces V, W the direct product is defined to be the **unique** vector space $V \oplus W$ equipped with maps $i_V : V \rightarrow V \oplus W$ and $i_W : W \rightarrow V \oplus W$ such that for any vector space U , with maps $j_V : V \rightarrow U$ and $j_W : W \rightarrow U$ there is a unique map

$$V \oplus W \rightarrow U$$

making the following diagram commute:

A commutative diagram illustrating the universal property of the direct sum. At the top is a vector space U . Below it is the direct sum $V \oplus W$. To the left of $V \oplus W$ is V , and to the right is W . A solid arrow labeled j_V points from V to U . A solid arrow labeled j_W points from W to U . A solid arrow labeled i_V points from V to $V \oplus W$. A solid arrow labeled i_W points from W to $V \oplus W$. A dashed vertical arrow points from $V \oplus W$ to U , representing the unique map $V \oplus W \rightarrow U$ that makes the diagram commute.


Universal constructions: Coproducts

Definition

Given two vector spaces V, W the direct product is defined to be the **unique** vector space $V \oplus W$ equipped with maps $i_V : V \rightarrow V \oplus W$ and $i_W : W \rightarrow V \oplus W$ such that for any vector space U , with maps $j_V : V \rightarrow U$ and $j_W : W \rightarrow U$ there is a unique map

$$V \oplus W \rightarrow U$$

making the following diagram commute:



This is an example of a very general construction: the **categorical coproduct**.

Definition

A **category** \mathcal{C} consists of

- ① a class of **objects** $\text{ob } \mathcal{C}$
- ② for any two objects $A, B \in \mathcal{C}$ a class of **morphisms** $\text{Hom}(A, B)$
- ③ For any three objects $A, B, C \in \text{ob } \mathcal{C}$, a binary operation called **composition**

$$\circ_{A,B,C} : \text{Hom}(A, B) \times \text{Hom}(B, C) \rightarrow \text{Hom}(A, C)$$

$$(f, g) \mapsto g \circ f$$

Axioms of composition

Composition is supposed to model function composition. This boils down to the following two axioms.

- The operation \circ is associative. In other words, if $h : A \rightarrow B$, $g : B \rightarrow C$ and $f : C \rightarrow D$ for $A, B, C, D \in \text{ob}(\mathcal{C})$, we have that:

$$(f \circ g) \circ h = f \circ (g \circ h)$$

as a function from $A \rightarrow D$.

- For $A \in \text{ob}(\mathcal{C})$, the hom-class $\text{Hom}(A, A)$ contains an identity element $\text{id}_A : A \rightarrow A$, such that for any $f : A \rightarrow B$, we have

$$\text{id}_A \circ f = f = f \circ \text{id}_A$$

Proposition

Consider the class of sets. For any two sets A, B , let

$$\mathrm{Hom}(A, B) = \{f : A \rightarrow B \mid f \text{ is a function.}\}$$

Composition is defined by function composition. This defines a category
Set.

Proposition

Consider the class of non-empty sets (A, a_0) with one distinguished element $a_0 \in A$. For any two pointed sets $(A, a_0), (B, b_0)$, define

$$\text{Hom}((A, a_0), (B, b_0)) = \{f : A \rightarrow B \mid f \text{ is a function such that } f(a_0) = b_0\}$$

*Composition is defined by function composition. This defines a category **Set_•**.*

Other Examples

Many of the mathematical structures that you have seen in previous lectures organize into categories

Examples

- Sets and functions. We call this category **Set**.
- Vector spaces and

Other Examples

Many of the mathematical structures that you have seen in previous lectures organize into categories

Examples

- Sets and functions. We call this category **Set**.
- Vector spaces and linear maps. We call this category **Vect**.
- Banach spaces and

Other Examples

Many of the mathematical structures that you have seen in previous lectures organize into categories

Examples

- Sets and functions. We call this category **Set**.
- Vector spaces and linear maps. We call this category **Vect**.
- Banach spaces and bounded linear maps. We call this category **Ban**.
- Metric spaces and

Other Examples

Many of the mathematical structures that you have seen in previous lectures organize into categories

Examples

- Sets and functions. We call this category **Set**.
- Vector spaces and linear maps. We call this category **Vect**.
- Banach spaces and bounded linear maps. We call this category **Ban**.
- Metric spaces and continuous maps (or short maps). We call this category **Met**.
- Topological spaces and continuous maps. We call this category **Top**. You will encounter it next week.

Universal constructions in a categories

Definition

Let \mathcal{C} be a category. Given $V, W \in \mathcal{C}$ the coproduct is defined to be the **unique** (if it exists) $V \amalg W$ equipped with maps $i_V : V \rightarrow V \oplus W$ and $i_W : W \rightarrow V \oplus W$ such that for $U \in \mathcal{C}$, with maps $j_V : V \rightarrow U$ and $j_W : W \rightarrow U$ there is a unique map

$$V \oplus W \rightarrow U$$

making the following diagram commute:

Functors

Given two categories, we can define "functions" between them.

Definition

Let \mathcal{C} and \mathcal{D} be categories. A functor F consists of a function

$$\text{ob } \mathcal{C} \rightarrow \text{ob } \mathcal{D}$$

$$A \mapsto F(A)$$

along with a map

$$\text{Hom}(A, B) \rightarrow \text{Hom}(F(A), F(B))$$

$$f \mapsto F(f)$$

for all $A, B \in \mathcal{C}$

Functors

Given two categories, we can define "functions" between them.

Definition

Let \mathcal{C} and \mathcal{D} be categories. A functor F consists of a function

$$\text{ob } \mathcal{C} \rightarrow \text{ob } \mathcal{D}$$

$$A \mapsto F(A)$$

along with a map

$$\text{Hom}(A, B) \rightarrow \text{Hom}(F(A), F(B))$$

$$f \mapsto F(f)$$

for all $A, B \in \mathcal{C}$. This map should have some properties with respect to composition.

Properties of functors

Functors should preserve the compositional structure of the category. In other words:

- **Composition is conserved** $F(f \circ g) = F(f) \circ F(g)$ for $g : A \rightarrow B$ and $f : B \rightarrow C$.
- **Identity maps are conserved**

$$F(\text{id}_A) = \text{id}_{F(A)}$$

There is a **forgetful functor**

$$U : \mathbf{Set}_\bullet \rightarrow \mathbf{Set}$$

$$U(A, a_0) = A \text{ and } U(f) = f.$$

Examples of functors

- There is a **forgetful functor**

$$|-| : \mathbf{Vect} \rightarrow \mathbf{Set}$$

$$X \mapsto |X|.$$

It assigns to every vector space its underlying category of points. It assigns to every morphism $f : V \rightarrow W$, the map $|f| : |V| \rightarrow |W|$, that sends x to $f(x)$. The map **forgets** the fact that X is a vector space.

- Similarly there are forgetful functors from metric, Banach and topological spaces to sets defined in the same way. There are other forgetful functors from Banach to Hilbert spaces, and from metric to topological spaces. You will encounter them on your homework.

Other examples of functors

- There is a **free functor**

$$\mathbb{R}[-] : \mathbf{Set} \rightarrow \mathbf{Vect}$$

$$X \mapsto \{f : X \rightarrow \mathbb{R} \mid \text{Supp } f \text{ is finite.}\}.$$

where $\text{Supp } f = \{x \in X \mid f(x) \neq 0\}$. For example, it sends $\{e_1, e_2\}$ to \mathbb{R}^2 . It sends a map $f : X \rightarrow Y$ to the map

$$\mathbb{R}[f] : \mathbb{R}[X] \rightarrow \mathbb{R}[Y]$$

$$\sum_{i \in X} a_i x \mapsto \sum_{i \in X} a_i f(x)$$

For example, it sends $e_1 \rightarrow e_2$ and $e_2 \rightarrow e_1$ to the linear map

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} : \mathbb{R}^2 \rightarrow \mathbb{R}^2.$$

Dual categories

Given any category \mathcal{C} , we can construct its **dual category** \mathcal{C}^{op} .

- The objects of \mathcal{C}^{op} are the same as \mathcal{C}^{op} . In other words

$$\text{ob } \mathcal{C}^{op} = \text{ob } \mathcal{C}$$

- We have

$$\text{Hom}_{\mathcal{C}^{op}}(A, B) = \text{Hom}_{\mathcal{C}}(B, A)$$

In other words, if there is a map $f : A \rightarrow B$ in \mathcal{C} , there is a map $f^{op} : B \rightarrow A$.

In fact $(-)^{op}$ is a functor.

The extension functor

Lemma

There is a functor $E: \mathbf{Set} \rightarrow \mathbf{Set}_\bullet$ and for all $X \in \mathbf{Set}$ a map $i_X: X \rightarrow U(EX)$, such that for all $(Y, y_0) \in \mathbf{Set}_\bullet$ and all maps $f: X \rightarrow U(Y, y_0)$, there exists a **unique** pointed map $\tilde{f}: EX \rightarrow (Y, y_0)$ making the following diagram commute:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y = U(Y, y_0) \\ \downarrow i_X & \nearrow U\tilde{f} & \\ U(EX) & & \end{array}$$

Question to class

- 1 How do you think you define EX and Ef ?
- 2 One definition:

$$EX = (\{\{0\} \cup \{(1, x) \mid x \in X\}\}, 0)$$

and

$$Ef(u) = \begin{cases} (1, f(x)) & \text{for } u = (1, x) \\ 0 & \text{for } u = 0. \end{cases}$$

Exercises on categories 1

- 1 Do sets and **injective** maps form a category? What about sets and strict inclusions?
- 2 Consider the set of **bracketed** words on some alphabet $\{e, a_1, a_2, \dots, a_n\}$ where, for any word w , we have that $we = ew = w$. Define

$$\text{Hom}(w_1, w_2) = \{u \mid (w_1)(u) = w_2\}$$

Define $u \circ v = (v)(u)$ Does this form a category? What about **unbracketed** words?

- 3 Show that a vector space V and all linear maps $V \rightarrow V$ form a 1-object category.
- 4 A **unital monoid** is a set X equipped with a binary associative operation $X \times X \rightarrow X$ that is associative and which has an identity element. Show that there is a one-to-one correspondence between unital monoids and categories with one element.

Exercises on universal constructions

- 1 Is the forgetful functor $\mathbf{Vect} \rightarrow \mathbf{Set}$ **faithful**? Can you obtain every map of sets this way?
- 2 What linear transformations are of the form $F(f)$ for $f : X \rightarrow Y$ for the free functor $F : \mathbf{Set} \rightarrow \mathbf{Vect}$. Hint: Express it in terms of matrices.

Exercises on functors

- 1 Is the forgetful functor $\mathbf{Vect} \rightarrow \mathbf{Set}$ **faithful**? Can you obtain every map of sets this way?
- 2 What linear transformations are of the form $F(f)$ for $f : X \rightarrow Y$ for the free functor $F : \mathbf{Set} \rightarrow \mathbf{Vect}$. Hint: Express it in terms of matrices.

Further Reading

- ① Course reference. Chapter 2 of Henning's notes for logic course. Available [here](#).
- ② A classic reference : Saunders Mac Lane. *Categories for the Working Mathematician*
- ③ An opinionated but motivational reference : Joseph A. Goguen. *A Categorical Manifesto*
- ④ A modern reference : Emily Riehl. *Category Theory in Context*