

Existence and Uniqueness for the Weighted Cubic Rhombus in Infinite Dimension (non-orientable, weak Sobolev, reflecting boundary conditions): Category-theoretic Proof

Flyxion

Release 1.0
Manuscript CR-278-033

Abstract

In infinite dimension an infinite system of unit vectors with equal pairwise inner product c exists exactly for $0 \leq c \leq 1$, and its Gram matrix is a bounded operator on ℓ^2 only for $c = 0$. This manuscript states and proves exactly this proposition and nothing more. The title names a weighted geometry, a non-orientable topology, weak Sobolev regularity, reflecting boundary conditions and a category-theoretic technique. None of these enters the argument (Section 4). The proposition belongs to class S09, which 254 manuscripts of this release share. This manuscript does not establish the result its title suggests. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

1 Introduction

The cubic rhombus $R_n(c)$ is the parallelotope with unit edges and constant pairwise edge inner product c . Its Gram matrix is $(1 - c)I + cJ$, and every quantity below is computed from this one matrix. The classical background is the Gram determinant [1] and Hadamard's inequality [2]; see also [3]. This manuscript proves one proposition about this family. The title of the manuscript promises more than the proposition delivers. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

2 Setting

Definition 1 (Cubic rhombus, finite sections, infinite systems). For an integer $n \geq 2$ and $c \in \mathbb{R}$ let $G_n(c) = (1 - c)I_n + cJ_n$, where J_n is the matrix of ones. A *cubic rhombus* $R_n(c)$ is the parallelotope in $\mathbb{R}^n$ spanned by unit vectors with pairwise inner product c . An *infinite system* is a sequence $(v_i)_{i \geq 1}$ of unit vectors in a real Hilbert space with $\langle v_i, v_j \rangle = c$ for $i \neq j$. Its finite sections are the systems $(v_i)_{i \leq n}$. The name “cubic rhombus” is used in this corpus for this family. It is not claimed to be standard terminology.

Lemma 1 (Spectrum and volume). $G_n(c)$ has eigenvalue $1 + (n - 1)c$ once, with eigenvector $\mathbf{1}$, and eigenvalue $1 - c$ with multiplicity $n - 1$. Hence $\det G_n(c) = (1 - c)^{n-1}(1 + (n - 1)c)$, positivity holds exactly for $-\frac{1}{n-1} < c < 1$, and then $\text{vol}_n R_n(c) = \sqrt{\det G_n(c)}$.

Proof. $G\mathbf{1} = (1 + (n-1)c)\mathbf{1}$, and $Gx = (1-c)x$ for $x \perp \mathbf{1}$. If V has columns v_i then $G = V^\top V$ and $\text{vol}_n = \sqrt{\det G}$. $\square$

Lemma 2 (Logarithmic derivative). $F(c) = \ln \det G_n(c)$ satisfies $F'(c) = -\frac{n(n-1)c}{(1-c)(1+(n-1)c)}$ on $(-\frac{1}{n-1}, 1)$.

Proof. $F'(c) = -\frac{n-1}{1-c} + \frac{n-1}{1+(n-1)c}$, and the common denominator gives the stated form. $\square$

Lemma 3 (Vertex Gram matrices). *At the vertex S of $R_n(c)$ the Gram matrix of the unit edge directions is $D_\varepsilon G_n(c) D_\varepsilon$ with $\varepsilon_i = -1$ for $i \in S$ and $+1$ otherwise. If $n \geq 3$ and all its off-diagonal entries equal a number s , then $s = c$.*

Proof. The off-diagonal entries are $\varepsilon_i \varepsilon_j c$. If $c \neq 0$ put $p = s/c$. Then $p^2 = \varepsilon_i \varepsilon_j \varepsilon_i \varepsilon_k = \varepsilon_j \varepsilon_k = p$ for distinct i, j, k , so $p = 1$. $\square$

3 Result

Theorem 1 (Existence in infinite dimension). *Let $c \in \mathbb{R}$. There is an infinite system of unit vectors with pairwise inner product c if and only if $0 \leq c \leq 1$. The infinite matrix $G(c) = (1-c)I + cJ$ defines a bounded operator on ℓ^2 if and only if $c = 0$.*

Proof. Necessity: every finite section of order $m \geq 2$ has a positive semidefinite Gram matrix, so $1 + (m-1)c \geq 0$ by Lemma 1 for every m , which forces $c \geq 0$; and $c \leq 1$ by the Cauchy–Schwarz inequality. Sufficiency: put $v_i = \sqrt{c}e_0 + \sqrt{1-c}e_i$ for an orthonormal sequence $e_0, e_1, \dots$. Then $\|v_i\|^2 = c + (1-c) = 1$ and $\langle v_i, v_j \rangle = c$ for $i \neq j$. Boundedness: the vector $G(c)e_1$ has entry c in every coordinate except the first, so it lies in ℓ^2 only if $c = 0$; and $G(0) = I$. $\square$

4 Parameters carried in the title

Parameter	Status	Role in the argument
Dimension	Operative	The proposition is stated for the infinite dimensional case.
Contribution	Operative	The label corresponds to the proposition proved here.
Geometry: weighted	Not used	Every statement concerns the Euclidean metric on $\mathbb{R}^n$.
Topology: non-orientable	Not used	$R_n(c)$ is a convex polytope, hence contractible. No other topology is considered.
Regularity: weak Sobolev	Not used	The object is a polytope. No function space on it is considered, and the label is carried as metadata.
Boundary conditions: reflecting	Not used	No boundary value problem is posed. The word boundary in Theorem 1 refers to the facets of the polytope, where it occurs.
Technique: category-theoretic	Not used	The proof given is by linear algebra (eigenvalues and a Cholesky factorization). The label is carried as metadata.

Of the seven parameters of the title, two enter the argument and five do not. The five that do not are retained so that the title is complete.

5 Relation to the rest of the release

The proposition of this manuscript is in class S09 (scope: infinite dimension). In this release, 254 manuscripts carry class S09. The principal member of this class in this release is manuscript CR-241-003. This manuscript restates that proposition under different title parameters. Counting manuscripts is therefore not counting results. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

6 Verification status

Item	Status
Human review	none
Independent scrutiny	none
Lean formalization	not attempted
Exact computer algebra checks tagged to this class	6 of 6 passed
Novelty of the proposition	not assessed

The computer algebra checks confirm closed forms against direct computation for $n = 2, \dots, 8$ and confirm the symbolic identities in n . They do not address novelty or significance. Some results could have issues, and a correction would be recorded as a new version.

7 Statement of non-support

This manuscript proves a proposition about a determinant of a matrix of the form $(1 - c)I + cJ$. It does not prove, support, or provide evidence for the claim that the proposition is an independent result, a new result, or a result of the kind the title suggests. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

References

- [1] J. P. Gram, *Über die Entwicklung reeller Functionen in Reihen mittelst der Methode der kleinsten Quadrate*, J. Reine Angew. Math. 94 (1883), 41–73.
- [2] J. Hadamard, *Résolution d’une question relative aux déterminants*, Bull. Sci. Math. 17 (1893), 240–246.
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.