

30¢

NO. 1  
OCT  
2026

A MATHEMATICAL EXPERIMENT  
TOO BIG TO READ!

APPROVED  
BY THE  
COMICS  
CODE



AUTHORITY

# THE CUBIC RHOMBUS

## AND THE INFINITE PAPER MACHINE!™


**14,000**  
MANUSCRIPTS  
WAS JUST  
THE BEGINNING!

SAME  
SHAPES!  
MANY  
VARIATIONS!  
COUNTLESS  
PAPERS!



$$\det G_n(c) = (1-c)^{n-1}(1+(n-1)c)$$

**SAME NUMBERS. DIFFERENT THINGS.**  
**WHEN PUBLICATION BECOMES AUTOMATIC,**  
**REVIEW DOES NOT.**



# THE CUBE THAT WOULDN'T STAY SQUARE!

**SAME EDGES.  
ALL FACES RHOMBI.  
ONE FAMILY  
OF SHAPES.**

**1**

Start with a cube.  
All edges have the same length,  
and all the angles are  $90^\circ$ .



A CUBE  
IS A SPECIAL CASE  
OF A CUBIC RHOMBUS.

**2**

Now push one corner.  
Keep all the edges the same length,  
but let the angles change.



Every face is  
still a rhombus.  
It's a different  
shape, but it  
belongs to the  
same family.

**3**

All of these shapes have the same edge length,  
but different angles between the edges.



$c = 0$   
(angles  $90^\circ$ )



$c = 1/2$   
(angles  $60^\circ$ )



$c$  near 1  
(angles very small)




Same edge length.  
Different angles. Same family of shapes.

**4**

So it's still a cube?


No, a cube is just  
one member of the family.  
These are all cubic rhombi  
— parallelotopes where  
all faces are rhombi.



A rhombus has  
equal sides but  
not necessarily  
right angles.

**5**

As we tilt it further,  
the angles get smaller  
and the shape becomes  
flatter and flatter...



**...BUT IT'S STILL  
THE SAME FAMILY  
OF SHAPES!**

A CUBIC RHOMBUS  $R_n(c)$  IS A  
PARALLELOTOPE WITH ALL EDGES  
EQUAL AND ALL FACES RHOMBI,  
CONTROLLED BY A SINGLE PARAMETER  $c$ .

# ONE NUMBER CONTROLS THEM ALL!

A SINGLE PARAMETER **C** SETS THE ANGLE BETWEEN EVERY PAIR OF EDGES.

**1** Take  $n$  unit vectors  $v_1, \dots, v_n$  so that every pair makes the same angle.

$$v_i \cdot v_j = \begin{cases} 1 & \text{if } i = j \\ c & \text{if } i \neq j \end{cases}$$

So  $c$  is the common inner product between different edges?


Exactly!

**2** The  $n \times n$  Gram matrix is

$$G_n(c) = (1 - c)I_n + cJ_n.$$

This matrix encodes the shape.


When  $c = 0$  we get a cube.



$c = 0$

All angles  $90^\circ$ .

**3** Increase  $c$  to  $1/2$ .



$c = 1/2$

All angles  $60^\circ$ .

Same edge lengths, but now the edges lean toward each other.

**4** Increase  $c$  further.



$c = 0.8$

Angles about  $37^\circ$ .

The shape gets more slanted, but the edges are still the same length!

**5** As  $c$  approaches 1, all the edges become almost parallel.


The edges are nearly parallel, but they still have unit length!



$c = 0.99$

Angles about  $8^\circ$ .

**6** We can also tilt in the other direction.



$c = -0.5$

Angles about  $120^\circ$ .

Negative  $c$  pushes the edges past  $90^\circ$ . As long as  $c > -1/(n-1)$ , the shape remains well-defined.

**7** One parameter. A whole family of shapes.

By turning a single dial, we can move smoothly from a cube to a flat shape — and even to shapes with obtuse angles!

THE PARAMETER **C**

$-\frac{1}{n-1}$

0

1

SAME EDGE LENGTHS.  
DIFFERENT ANGLES.  
SAME FAMILY.

It's amazing — so many different shapes, all controlled by one number!

# THE CASE OF THE VANISHING VOLUME!

CHANGE THE ANGLES, NOT THE EDGES.



**SAME EDGE LENGTHS. DIFFERENT AMOUNTS OF SPACE.**

## 1 A CUBE HAS UNIT-LENGTH EDGES.

Every edge is the same length.

A cubic rhombus has  $n$  edges of equal length, with the same angle between every pair.

## 2 SAME EDGES, DIFFERENT SHAPE, DIFFERENT VOLUME.

Then why do they take up different amounts of space?

The edges are still the same length, but the angles have changed.

$c=0$  (right angles)       $c>0$  (slanting angles)

## 3 TILT THE SHAPE — THE VOLUME SHRINKS.

The angles are changing!

As the common angle increases ( $c$  gets larger), the shape becomes more slanted and takes up less space.

$c$

$c=0$      $c=0.5$      $c=0.8$      $c=0.95$

## 4 THE DETERMINANT GIVES THE VOLUME.

The volume is determined by the determinant of the Gram matrix.

Gram matrix for the cubic rhombus:

$$G_n(c) = (1-c)I_n + cJ_n$$

$\det G_n(c) = (1-c)^{n-1}(1+(n-1)c)$

The squared volume is the determinant:

$$\text{Vol}(R_n(c)) = \sqrt{\det G_n(c)}$$

$c$  is the common cosine of the angles between edges:

$$c = \cos \theta$$

$$0 \leq c < 1$$

(for a nondegenerate shape in any dimension  $n$ ).

## 5 AS $c$ APPROACHES 1, THE VOLUME APPROACHES ZERO.

$c=0.9$      $c=0.99$      $c=0.999$

Even though the edges are still length 1...

... the volume shrinks to zero!

When  $c \rightarrow 1$ , the edges become nearly parallel and the shape collapses to a lower-dimensional object, with zero volume.

## 6 THE EDGES DIDN'T SHRINK — THE SPACE DID!

THE EDGES DIDN'T SHRINK. THE SPACE BETWEEN THEM DID!

SAME LENGTHS. DIFFERENT VOLUME. IT'S THE ANGLES.

**THE CUBIC RHOMBUS: SAME EDGES. MANY SHAPES. A SURPRISING VOLUME.**

**NEXT:**  
TWO SHAPES, ONE VOLUME?

# THE GEOMETRIC DETECTIVE!

A NUMERICAL INVARIANT DOES NOT ALWAYS IDENTIFY AN OBJECT.



**SAME VOLUME. DIFFERENT SHAPES.**

## 1 THE CASE BEGINS.

We already know the volume of a cubic rhombus  $R_n(c)$  with unit edges is  $V(c) = (1-c)^{n-1}(1+(n-1)c)$ . Different values of  $c$  give different volumes.

So if two shapes have the same volume, are they the same shape?

## 2 TWO SUSPECTS, SAME VOLUME.

**SUSPECT A**  
( $c = 0.5$ )

**SUSPECT B**  
( $c = -0.5$ )

Vol  $\approx 0.61$       Vol  $\approx 0.61$

Same volume. Different angles. Different shape. Are they actually the same object?

The volume alone does not tell us.

## 3 LOOK CLOSER AT THE INVARIANTS.

The volume comes from the determinant of the Gram matrix, but many different Gram matrices can have the same determinant.

Gram matrix:  
 $G_n(c) = (1-c)I_n + cJ_n$   
 $\det G_n(c) = (1-c)^{n-1}(1+(n-1)c)$

Different values of  $c$  can produce the same determinant. A numerical invariant is not a fingerprint.

## 4 INTERROGATE THE SUSPECTS.

Both have unit edges and the same volume, but their angles are different.

So the same volume does not identify a unique shape?

| SUSPECT A<br>( $c = 0.5$ )      | SUSPECT B<br>( $c = -0.5$ )      |
|---------------------------------|----------------------------------|
| edges: 1                        | edges: 1                         |
| angle $\theta \approx 60^\circ$ | angle $\theta \approx 120^\circ$ |
| volume $\approx 0.61$           | volume $\approx 0.61$            |

Many non-congruent cubic rhombi share the same volume.

## 5 A WHOLE FAMILY OF LOOK-ALIKES.

For a fixed volume, there can be many different shapes with different angles — even a continuous family.

$c = -0.5$  Vol  $\approx 0.61$      $c = 0$  Vol = 1     $c = 0.5$  Vol  $\approx 0.61$      $c = 0.8$  Vol  $\approx 0.27$      $c = 0.95$  Vol  $\approx 0.07$

Same volume. Many shapes. The invariant does not determine the object.

## 6 THE DETECTIVE'S CONCLUSION.

The volume answers a numerical question, but it does not answer the geometric question "which object?"

**SAME VOLUME  $\neq$  SAME SHAPE**

- Volume is an invariant.
- Many different shapes can share the same invariant.
- Additional information (such as angles or a full Gram matrix) is needed to identify the object.

## 7 THE BROADER LESSON.

A correct calculation can be mathematically valid and still leave important questions unanswered.

Just like with manuscripts: a formally correct result does not automatically tell us whether the original research question was understood, whether the object is novel, or whether it matters.


## 8 NOT A FINGERPRINT.

**Volume only**  
Vol  $\approx 0.61$   
**NOT UNIQUE**

**Full description**  
• edge lengths  
• angles ( $c$ )  
• Gram matrix  
• additional invariants  
**IDENTIFIES THE SHAPE**

A single numerical invariant can describe many different things. Mathematics needs more than one number — and so does research evaluation.

**THE SAME NUMBERS CAN DESCRIBE MANY DIFFERENT THINGS. CORRECTNESS IS NOT THE SAME AS UNIQUENESS.**



# THE FOURTH DIMENSION STRIKES!

THE CUBIC RHOMBUS LIVES BEYOND THREE DIMENSIONS.


SAME RULE.  
BIGGER SPACE.  
STRANGER SHAPES.

A simple parameter.  
A whole new world.




## 1 FROM 2D TO 3D ...

Start with a square.  
That's a 2D cubic rhombus.



$n = 2$



$n = 3$

So the 3D version  
is just a cube?

Exactly.  
Now watch  
what happens  
next.

## 2 THE FOURTH DIMENSION

For  $n = 4$ , the same rule  
still applies. But we can't  
see a 4D shape directly.



This is a  
"shadow" or  
projection of a  
4D cubic rhombus.

The real shape  
lives in a space  
we can't see.

The same  
parameter  $c$   
controls the  
angles here too.

## 3 HIGHER DIMENSIONS

The pattern continues. We can build  
cubic rhombi in any dimension  $n$ ,  
using the same Gram matrix:

$$G_n(c) = (1 - c)I_n + cJ_n$$

But as  $n$  grows,  
the shapes get  
harder to imagine.



$n = 5$



$n = 10$



$n = 20$




$n = 100$

## 4 THE VOLUME DILEMMA

The volume of a cubic rhombus is

$$\text{Vol}(R_n(c)) = \sqrt{(1 - c)^{n-1} (1 + (n - 1)c)}.$$


As  $c$  approaches 1, the volume shrinks  
to zero. In higher dimensions, this happens  
even faster.



$c = 0$



$c = 0.5$



$c = 0.9$




$c \rightarrow 1$

The edges  
stay the same  
length, but the  
shape flattens  
out.

## 5 THE INFINITE-DIMENSIONAL TRAP

For an infinite set of unit vectors  
with the same angle, the Gram matrix  
is well-behaved — as long as  $0 \leq c \leq 1$ .

But for  $c \neq 0$ , the infinite  
Gram matrix is not a valid  
operator on square-summable  
sequences.



So the pattern  
breaks when we  
try to continue  
forever?

Yes. Finite sections work.  
The infinite case is different.

## 6 THE GREAT PAPER FACTORY

14,000  
manuscripts!  
280 families!  
14 distinct  
statements!

SAME  
IDEA.  
DIFFERENT  
WRAPPERS.

THE MATHEMATICS:  
14  
STATEMENTS

THE MATHEMATICS:  
14  
STATEMENTS

## 7 THE MACHINE CHECKS THE PROOFS

The formal statements  
compile. The proofs check out.

But does each proved  
statement actually match  
the original manuscript  
claim?

That's a  
human review.  
Not a  
machine job.

- ✓ 14 classes verified
- ✓ No sorries
- ✓ No extra axioms
- ✓ Axioms audited

LEAN 4

MANUSCRIPTS

LEAN PROOFS

REVIEW LEDGER

## 8 THE REAL QUESTION

We can generate  
14,000 papers.

We can prove  
14 statements.

But the real question  
is still the same...



DOES IT  
SAY WHAT  
WE THINK  
IT SAYS?

THE CUBIC RHOMBUS: A SIMPLE SHAPE. A DEEPER STORY.

# THE INFINITE-DIMENSIONAL TRAP!

A PATTERN THAT WORKS FOR ANY FINITE SIZE ... BUT NOT IN THE SAME WAY FOREVER.

SAME ANGLE BETWEEN EVERY PAIR OF VECTORS.

## 1 START WITH A FEW VECTORS.

Take three unit vectors with the same angle between each pair.

Each pair has the same inner product  $c$  (where  $c = \cos \theta$ ).

$$v_i \cdot v_j = c \quad (i \neq j)$$

## 2 BUILD A WHOLE SET.

We can do this for any finite number  $n$ . Just keep adding another vector with the same angle to all the others.

No problem! This works for  $n = 4, 5, 10, 100, \dots$

Any finite set with  $0 \leq c < 1$  can be realized in  $n$ -dimensional space.

The  $n \times n$  Gram matrix is  $G_n(c) = (1 - c)I_n + cJ_n$ .

## 3 SO LET'S TRY INFINITY ...

Why stop? Let's take infinitely many unit vectors, all with the same inner product  $c$ !

I looks fine, but does it actually exist a vector set in a Hilbert space?

Formally, the infinite Gram matrix would have 1 on the diagonal and  $c$  off the diagonal.

## 4 FINITE SECTIONS WORK.

Every finite subset behaves exactly as expected, with the same Gram matrix.

$n = 3$

$n = 10$

$n = 100$

No matter how large a finite  $n$  is, we can find vectors with the right pairwise angles.

The trouble only appears when we try to take the limit to infinitely many vectors.

## 5 THE OPERATOR PROBLEM.

For  $c \neq 0$ , this infinite matrix does not define a valid operator on the usual space of square-summable sequences ( $\ell^2$ ).

So the pattern breaks when we try to continue forever?

| INFINITE GRAM MATRIX |          |          |          |          |
|----------------------|----------|----------|----------|----------|
| 1                    | $c$      | $c$      | $c$      | $\dots$  |
| $c$                  | 1        | $c$      | $c$      | $\dots$  |
| $c$                  | $c$      | 1        | $c$      | $\dots$  |
| $\vdots$             | $\vdots$ | $\vdots$ | $\vdots$ | $\ddots$ |

Finite constructions do not automatically extend to a well-behaved infinite object.

## 6 A SIMPLE EXAMPLE.

The matrix with 1 on the diagonal and  $c$  off the diagonal acts badly on sequences like  $(1, 1, 1, \dots)$ .

Try multiplying by the all-ones sequence:

$$(Gx)_i = x_i + c \sum_{j \neq i} x_j.$$

For  $x = (1, 1, 1, \dots)$ , the sum does not converge when  $c \neq 0$ .

**INFINITE SUM! NOT DEFINED!**

## 7 WHAT STILL WORKS?

So we can have arbitrarily large finite families with the same angle, but we cannot build an infinite family in the usual sense?

Exactly. Positive semidefiniteness holds for all finite  $n$  when  $0 \leq c \leq 1$ , but the infinite operator is only well-defined in the trivial case  $c = 0$  (the orthogonal basis).

Finitely many: yes.  
Infinitely many: not in the same way.  
The limit changes the problem.

## 8 THE LESSON.

Extending a pattern forever can fail, even when every finite case works.

**SAME IDEA, DIFFERENT REGIME:**

- Finite  $n$ : valid Gram matrix.
- Infinite  $n$ : operator may not exist.
- $c = 0$ : orthogonal basis works.
- $0 < c \leq 1$ : finite sets ok, but no corresponding  $\ell^2$  operator.

The infinite version is not just a bigger copy. It's a different mathematical question.

**A PATTERN CAN SURVIVE EVERY FINITE TEST AND STILL FAIL IN THE LIMIT.**

# THE GREAT PAPER FACTORY!

14,000 PAPERS.  
280 FAMILIES.  
14 DISTINCT STATEMENTS?

MANY TITLES.  
SAME IDEAS.




1

## A MODEST IDEA ...

Let's study the cubic rhombus  $R_n(c)$  — the parallelotope with all edges equal and all faces rhombi.

That's one clear idea. How many results could there be?



Gram matrix:  
 $G_n(c) = (1 - c)I_n + cJ_n$

2

## ... BECOMES A FLOOD.

Brilliant! Let's generate thousands of manuscripts!

NEW PROOF OF THE DETERMINANT

A DIFFERENT DERIVATION

ANOTHER PERSPECTIVE

GENERALIZED BOUND

A NOTE ON SPECIAL CASES

SOME RELATED OBSERVATIONS

FURTHER REMARKS

EXTENSION TO  $n = 7$

A VARIATION ON THE THEME

3

## DIFFERENT WRAPPERS ...

I've rewritten the proof using a new notation!

I've added a corollary!

I've changed the title and reordered the argument!

I've applied it to a slightly different special case!

A NEW APPROACH to  $R_n(c)$

AN ALTERNATIVE DERIVATION of the Gram Determinant

FURTHER RESULTS on the Cubic Rhombus

EXTENSIONS and APPLICATIONS

... BUT OFTEN THE SAME CORE RESULT.

4

## CLASSIFY THE CONTENT.

These 14,000 manuscripts fall into 280 families with very similar content.

And when we group those families by mathematical claim, we get 14 distinct statements.

MANUSCRIPTS (14,000)

FAMILIES (280)

STATEMENTS (14)

5

## EXAMPLES OF VARIATIONS.

Determinant of  $G_n(c)$  revisited

Volume formula with an alternative argument

Eigenvalues and positivity of  $R_n(c)$

On the geometry of equiangular vectors

Same theorem. Different proof order.

Same statement. Different presentation.

Same result. Different notation.

Related claim. Same mathematical content.

6

## A QUICK REDUCTION.

All of these papers reduce to fourteen mathematical claims.

So the mountain of papers is mostly repetition, variation, and special cases — not 14,000 new discoveries.

14 DISTINCT STATEMENTS

7

## WHY DOES THIS HAPPEN?

- different notation
- reordered proofs
- additional lemmas
- minor generalizations
- changed emphasis
- overlapping special cases
- new titles and wording
- automated generation

Many papers look different, but they express the same underlying ideas.

8

## THE LESSON.

Quantity is not the same as novelty. We can generate thousands of manuscripts, but they do not create thousands of distinct mathematical results.

The real work is to understand what the statements actually are — and whether they say what we think they say.

**MANY PAPERS. FEW IDEAS. SAME SHAPE.**



THE CUBIC RHOMBUS: A SIMPLE IDEA WITH MANY FACES.

# THE MACHINE CHECKED THE PROOFS!

FORMAL CORRECTNESS IS NOT THE SAME AS FAITHFUL MEANING.

PROOFS CAN SUCCEED. QUESTIONS CAN STILL REMAIN.

?

## 1 WE GIVE IT A STATEMENT.

Here is one of our results. We formalized it in Lean 4.

It compiles! The proof checks!

Theorem 1.

$$\det G_n(c) = (1-c)^{n-1} (1+(n-1)c)$$

LEAN 4

✓ Proof checked. No errors.

## 2 FORMAL SUCCESS HAS A SCOPE.

The proof checker verifies that the formal statement follows from its stated assumptions.

This means the formal statement is correct — within the system.

✓ Syntax: OK  
✓ Types: OK  
✓ Logic: OK  
✓ Proof: OK


LEAN 4

But it does not tell us whether this is the same as what the original manuscript claimed.

## 3 THE TRANSLATION STEP CAN CHANGE THINGS.

ORIGINAL MANUSCRIPT

The volume should go to zero as the edges become more parallel.



TRANSLATION CHOICES

- ☐ notation
- ☐ assumptions
- ☐ domain of  $c$
- ☐ dimension  $n$
- ☐ what is measured
- ☐ extra conditions

FORMAL STATEMENT

For  $n \geq 2$  and  $-1/(n-1) < c < 1$ ,  
 $\det G_n(c) = (1-c)^{n-1} (1+(n-1)c)$ .

Different reasonable translations can lead to different formal statements.

## 4 BOTH CAN BE RIGHT — ABOUT DIFFERENT THINGS.

This formal result is true. It follows from the stated assumptions.

But did it answer the question the original author intended? Did we add or drop something important?

Original question

?

Formal theorem ✓

A proved statement can be correct, even if it does not match the intent of the original claim, or if key conditions were changed.

## 5 WE NEED A REVIEW LEDGER.

CHECK BOTH LAYERS

Formal layer (proof checker)

- ☐ type checks
- ☐ no errors
- ☐ follows assumptions
- ☐ stated theorem true

Content layer (human review)

- ☐ matches original claim?
- ☐ same conditions?
- ☐ same domain?
- ☐ same quantities?
- ☐ no added assumptions?

Verification and faithful translation are separate tasks.

## 6 THE SURPRISE AT SCALE.

We generated 14,000 manuscripts!

But after classification, they fall into 280 families and 14 distinct statements.

14,000 MANUSCRIPTS

14 STATEMENTS

A huge number of papers can express a small number of core ideas.

## 7 SOME CLAIMS STILL NEED HUMAN JUDGMENT.

The machine cannot decide these questions:

- Which question was intended?
- Are the conditions appropriate?
- Does the formal result apply?
- Are there important alternatives?
- What does the result mean in the original context?

So proof checking is a powerful tool, but it does not replace reading, comparison, and critical judgment.

## 8 THE TAKEAWAY.

The machine confirms that this formal statement is correct.

Now we still have to ask: is this the claim we meant? What did it leave out? What else should we test?

Proof checked ✓

Same as the original claim?

?

PROOFS CHECK FORM. HUMANS CHECK MEANING.



THE CUBIC RHOMBUS: MANY QUESTIONS. SAME SHAPE.

# WHO IS SUPPOSED TO READ ALL THIS?!

AUTOMATIC GENERATION IS EASY. HUMAN UNDERSTANDING IS NOT.

ONE SHAPE WAS JUST THE BEGINNING.

1

WE GENERATED 14,000 MANUSCRIPTS.

Here it is! 14,000 manuscripts on the cubic rhombus  $R_n(c)$ , organized into 280 families and 14 distinct statements.

That's an impressive number. But that's just one shape, right?

14,000 MANUSCRIPTS  
280 FAMILIES  
14 STATEMENTS

2

ONE SHAPE. MANY OTHERS.

GEOMETRIC FAMILIES (A SMALL SAMPLE)

- simplices
- parallelotopes
- ellipsoids
- zonotopes
- polytopes
- lattices
- manifolds
- tilings
- graphs
- metric spaces
- algebraic varieties
- and many more ...

Each family has parameters too. There are infinitely many variations.

The cubic rhombus was one arbitrarily chosen object. We could do this for all of these.

3

THE MACHINE KEEPS GOING.

NEXT SHAPE

GENERATE MANUSCRIPTS

- configure object
- instantiate parameters
- produce statements
- write proofs
- verify formally
- package manuscripts

4

THE NUMBERS EXPLODE.

CUBIC RHOMBUS  
14,000 manuscripts

ELLIPSOIDS  
27,000 manuscripts

POLYTOPES  
48,000 manuscripts

LATTICES  
92,000 manuscripts

MANIFOLDS  
470,000 manuscripts

MANIFOLDS  
MILLIONS ... BILLIONS ... ?

AND BEYOND ...

5

IT ARRIVES AT THE JOURNALS.

JOURNAL OF MATHEMATICS

Here are this month's submissions.

This is just the cubic rhombus? There's more?

6

REVIEWERS ARE OVERWHELMED.

TO READ

TO REVIEW

TO DECIDE

This is impossible!

These are all on the same mathematical object?!

How many more shapes are there?

We can't even read this, let alone check novelty, significance, or context.

7

THE SYSTEM SAYS IT'S ALL VERIFIED.

The system confirms that every proof is formally correct!

Formal correctness does not tell us whether the original claim is important, new, or even what it means in context.

Formal proofs: ✓  
Type checks: ✓  
No errors: ✓  
Follows assumptions: ✓  
Verified (Lean): ✓

ALL FORMAL PROOFS VERIFIED!

8

BUT WHO IS SUPPOSED TO READ ALL THIS?!

WHO IS SUPPOSED TO READ ALL THIS?!

EDITOR

- check for novelty
- understand context
- evaluate significance
- detect errors
- assess assumptions
- compare with prior work
- decide what to publish
- all with limited people, time and resources

9

AN EFFECTIVELY INEXHAUSTIBLE SUPPLY.

MATHEMATICAL PUBLICATIONS

ONE SHAPE  
14,000

ANOTHER SHAPE  
48,000

ANOTHER SHAPE  
92,000

MANY SHAPES  
MILLIONS

CONTINUOUS PARAMETERS  
COUNTLESS VARIATIONS

...

The mathematical subject matter is effectively inexhaustible. But the number of mathematicians is not.

WHEN PUBLICATION BECOMES AUTOMATIC, REVIEW DOES NOT.

30¢

No. 1  
OCT  
2026APPROVED  
BY THE  
COMICS  
CODE  
AUTHORITY

# THE CUBIC RHOMBUS ACTION COLLECTION™

**ALL NEW!**  
**ALL DIFFERENT!**  
**ALL INFINITE!\***

Get in on the action  
with **MATHEMATICAL  
POWER!**

Create endless families  
of shapes, explore new  
parameters, and produce  
thousands of manuscripts!  
The Cubic Rhombus  
Action Collection lets  
you build your own  
research empire!

**GENERATE!**  
**VERIFY!**  
**PUBLISH!**  
**REPEAT!**

**SAME  
NUMBERS!  
DIFFERENT  
THINGS!**

**COLLECT THEM ALL!**

**EACH FAMILY HAS ITS OWN POWERS!**



**CUBIC RHOMBUS**  
14,000 manuscripts  
in 280 families!\*



**SIMPLEX**  
??? manuscripts  
in ??? families!




**ELLIPSOID**  
??? manuscripts  
in ??? families!



**ZONOTOPE**  
??? manuscripts  
in ??? families!



**POLYTOPE**  
??? manuscripts  
in ??? families!



**LATTICE**  
??? manuscripts  
in ??? families!



**MANIFOLD**  
??? manuscripts  
in ??? families!



**GRAPH**  
??? manuscripts  
in ??? families!



**METRIC SPACE**  
??? manuscripts  
in ??? families!



**AND MANY MORE!**  
??? manuscripts  
in ??? families!



9 781234 567898

**SAME EDGE LENGTHS. MANY SHAPES.  
A SURPRISING AMOUNT OF PAPER.  
COLLECT THEM ALL — IF YOU HAVE THE SHELF SPACE!**

**MATHEMATICS  
NEVER ENDS!**