

A Counterexample for the Euclidean Cubic Rhombus in Dimension n (toroidal, smooth, absorbing boundary conditions): Combinatorial Proof

Flyxion

Release 1.0
Manuscript CR-202-011

Abstract

For n the volume of a cubic rhombus takes each value in $(0, 1)$ exactly twice as c varies. This manuscript states and proves exactly this proposition and nothing more. The title names a Euclidean geometry, a toroidal topology, smooth regularity, absorbing boundary conditions and a combinatorial technique. None of these enters the argument (Section 4). The proposition belongs to class S07, which 1224 manuscripts of this release share. This manuscript does not establish the result its title suggests. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

1 Introduction

The cubic rhombus $R_n(c)$ is the parallelotope with unit edges and constant pairwise edge inner product c . Its Gram matrix is $(1 - c)I + cJ$, and every quantity below is computed from this one matrix. The classical background is the Gram determinant [1] and Hadamard's inequality [2]; see also [3]. This manuscript proves one proposition about this family. The title of the manuscript promises more than the proposition delivers. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

2 Setting

Definition 1 (Cubic rhombus). For an integer $n \geq 2$ and $c \in \mathbb{R}$ let $G_n(c) = (1 - c)I_n + cJ_n$, where J_n is the $n \times n$ matrix of ones. A *cubic rhombus* $R_n(c)$ is the parallelotope $\{\sum_i t_i v_i : 0 \leq t_i \leq 1\} \subset \mathbb{R}^n$ spanned by unit vectors $v_1, \dots, v_n$ with $\langle v_i, v_j \rangle = c$ for $i \neq j$, that is, with Gram matrix $G_n(c)$. The case $c = 0$ is the unit cube, $n = 2$ gives a rhombus, and $n = 3$ gives a rhombohedron. The name “cubic rhombus” is used in this corpus for this family. It is not claimed to be standard terminology.

Lemma 1 (Spectrum and volume). *The matrix $G_n(c)$ has eigenvalue $1 + (n - 1)c$ with eigenvector $\mathbf{1} = (1, \dots, 1)^\top$, and eigenvalue $1 - c$ with multiplicity $n - 1$ on $\mathbf{1}^\perp$. Hence $\det G_n(c) = (1 - c)^{n-1}(1 + (n - 1)c)$, the matrix is positive definite if and only if $-\frac{1}{n-1} < c < 1$, and in that case $\text{vol}_n R_n(c) = \sqrt{\det G_n(c)}$.*

Proof. $J\mathbf{1} = n\mathbf{1}$, so $G\mathbf{1} = (1 - c + nc)\mathbf{1} = (1 + (n - 1)c)\mathbf{1}$. If $x \perp \mathbf{1}$ then $Jx = 0$, so $Gx = (1 - c)x$. These eigenspaces span $\mathbb{R}^n$, which gives the spectrum and the determinant, and positivity of both eigenvalues is the stated range. If V is the matrix with columns v_i then $G = V^T V$ and $\text{vol}_n = |\det V| = \sqrt{\det G}$. $\square$

Lemma 2 (Facets). *A facet of $R_n(c)$ is spanned by $n - 1$ of the edge vectors, and its Gram matrix is the principal $(n - 1) \times (n - 1)$ submatrix of $G_n(c)$, which is $G_{n-1}(c)$. Its $(n - 1)$ -volume is $\sqrt{(1 - c)^{n-2}(1 + (n - 2)c)}$, and the surface area of $R_n(c)$ is $2n$ times this number.*

Proof. Deleting a row and the matching column of $(1 - c)I + cJ$ leaves $(1 - c)I + cJ$ of order $n - 1$. Apply Lemma 1 in dimension $n - 1$. There are $2n$ facets, in n parallel pairs. $\square$

Lemma 3 (Logarithmic derivative). *On $(-\frac{1}{n-1}, 1)$ let $F(c) = \ln \det G_n(c) = (n - 1) \ln(1 - c) + \ln(1 + (n - 1)c)$. Then*

$$F'(c) = -\frac{n(n - 1)c}{(1 - c)(1 + (n - 1)c)},$$

the denominator is positive, and so F' has the sign of $-c$.

Proof. $F'(c) = -\frac{n-1}{1-c} + \frac{n-1}{1+(n-1)c} = (n - 1) \frac{(1-c) - (1+(n-1)c)}{(1-c)(1+(n-1)c)} = -\frac{n(n-1)c}{(1-c)(1+(n-1)c)}$. $\square$

Lemma 4 (Vertex Gram matrices). *The vertices of $R_n(c)$ are the points $\sum_{i \in S} v_i$ for $S \subseteq \{1, \dots, n\}$. At the vertex S the unit edge directions are $\varepsilon_i v_i$ with $\varepsilon_i = -1$ for $i \in S$ and $\varepsilon_i = +1$ otherwise, so their Gram matrix is $D_\varepsilon G_n(c) D_\varepsilon$ with $D_\varepsilon = \text{diag}(\varepsilon)$. Suppose $n \geq 3$ and that all off-diagonal entries of $D_\varepsilon G_n(c) D_\varepsilon$ equal one number s . Then $s = c$. For $n = 2$ the conclusion fails: the vertices with $\varepsilon_1 \varepsilon_2 = -1$ give $s = -c$.*

Proof. The first statements are immediate from the definition. The off-diagonal entries are $\varepsilon_i \varepsilon_j c$. If $c = 0$ then $s = 0 = c$. If $c \neq 0$ put $p = s/c$, so $\varepsilon_i \varepsilon_j = p$ for all $i \neq j$. For distinct i, j, k we have $\varepsilon_i \varepsilon_j \cdot \varepsilon_i \varepsilon_k = \varepsilon_j \varepsilon_k$, that is $p^2 = p$, so $p = 1$. $\square$

3 Result

Theorem 1 (A counterexample to volume rigidity). *Let $n \geq 2$ be an integer. The map $c \mapsto \text{vol}_n R_n(c)$ is not injective: for every $v \in (0, 1)$ there are exactly two values $c_- \in (-\frac{1}{n-1}, 0)$ and $c_+ \in (0, 1)$ with $\text{vol}_n R_n(c_\pm) = v$. For $n \geq 3$ the bodies $R_n(c_-)$ and $R_n(c_+)$ are not congruent, so volume does not determine the shape. For $n = 2$ the two bodies are congruent: $R_2(c)$ and $R_2(-c)$ are the same rhombus, so in dimension 2 the statement concerns the parameter c only and is not a counterexample about shapes.*

Proof. By Lemma 3, F is strictly increasing on $(-\frac{1}{n-1}, 0]$ and strictly decreasing on $[0, 1)$, with $F(0) = 0$, and $F(c) \rightarrow -\infty$ at both endpoints of the interval, because $\det G_n(c) \rightarrow 0$ there. Hence $\text{vol} = e^{F/2}$ maps each of the two open branches bijectively onto $(0, 1)$, which gives exactly one solution on each side of 0. Non-congruence for $n \geq 3$: a congruence $R_n(c_-) \rightarrow R_n(c_+)$ maps the vertex $S = \emptyset$ of $R_n(c_-)$ to a vertex of $R_n(c_+)$ and maps its edge directions to the edge directions there, preserving inner products. The Gram matrix at $S = \emptyset$ has all off-diagonal entries equal to c_- , and the Gram matrix at the image vertex is $D_\varepsilon G_n(c_+) D_\varepsilon$ up to permutation. By Lemma 4, $c_- = c_+$, which contradicts $c_- < 0 < c_+$. For $n = 2$ the two rhombi are congruent, as follows. A

rhombus with unit sides is determined up to congruence by the angle θ between two adjacent sides, and it has the angle $\pi - \theta$ at its other two vertices. So the rhombus with angle θ and the rhombus with angle $\pi - \theta$ are the same rhombus, and $\cos(\pi - \theta) = -\cos \theta$ gives $R_2(c) \cong R_2(-c)$. $\square$

4 Parameters carried in the title

Parameter	Status	Role in the argument
Dimension	Operative	The proposition is stated for dimension n .
Contribution	Operative	The statement is a counterexample to the injectivity of volume as a function of c , with the shape consequences stated in the theorem.
Geometry: Euclidean	Not used	Every statement concerns the Euclidean metric on $\mathbb{R}^n$.
Topology: toroidal	Not used	$R_n(c)$ is a convex polytope, hence contractible. No other topology is considered.
Regularity: smooth	Not used	The object is a polytope. No function space on it is considered, and the label is carried as metadata.
Boundary conditions: absorbing	Not used	No boundary value problem is posed. The word boundary in Theorem 1 refers to the facets of the polytope, where it occurs.
Technique: combinatorial	Not used	The proof given is by monotonicity, the intermediate value theorem, and vertex Gram matrices. The label is carried as metadata.

Of the seven parameters of the title, two enter the argument and five do not. The five that do not are retained so that the title is complete.

5 Relation to the rest of the release

The proposition of this manuscript is in class S07 (scope: general n). In this release, 1224 manuscripts carry class S07. The principal member of this class in this release is manuscript CR-041-003. This manuscript restates that proposition under different title parameters. Counting manuscripts is therefore not counting results. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

6 Verification status

Item	Status
Human review	none
Independent scrutiny	none
Lean formalization	not attempted
Exact computer algebra checks tagged to this class	6 of 6 passed
Novelty of the proposition	not assessed

The computer algebra checks confirm closed forms against direct computation for $n = 2, \dots, 8$ and confirm the symbolic identities in n . They do not address novelty or significance. Some results could have issues, and a correction would be recorded as a new version.

7 Statement of non-support

This manuscript proves a proposition about a determinant of a matrix of the form $(1 - c)I + cJ$. It does not prove, support, or provide evidence for the claim that the proposition is an independent result, a new result, or a result of the kind the title suggests. This manuscript does not support the hypothesis that it is an independent mathematical contribution.

References

- [1] J. P. Gram, *Über die Entwicklung reeller Functionen in Reihen mittelst der Methode der kleinsten Quadrate*, J. Reine Angew. Math. 94 (1883), 41–73.
- [2] J. Hadamard, *Résolution d’une question relative aux déterminants*, Bull. Sci. Math. 17 (1893), 240–246.
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.