


Pool-3-041-AZP-226_S45.unmapped.rma6



Detailed description: The graph plots the C2T/G2A rate on the y-axis (0.00 to 0.30) against Read Position on the x-axis (1 to -2). The Naive method (red line) starts at ~0.27 at position 1, drops to ~0.10 at position 2, and then fluctuates slightly around 0.00 from position 5 onwards. The Naive+GC method (grey line) starts at ~0.01 at position 1 and remains near 0.00 for all subsequent positions. The Naive+GC+GC3 method (blue line) starts at ~0.00 at position 1 and remains near 0.00 for all subsequent positions.

Read Position	Naive	Naive+GC	Naive+GC+GC3
1	0.27	0.01	0.00
2	0.10	0.00	0.00
3	0.05	0.00	0.00
4	0.00	0.00	0.00
5	0.01	0.00	0.00
6	0.01	0.00	0.00
7	0.02	0.00	0.00
8	0.00	0.00	0.00
9	0.00	0.00	0.00
10	0.01	0.00	0.00
-10	0.00	0.00	0.00
-8	0.00	0.00	0.00
-6	0.00	0.00	0.00
-4	0.00	0.00	0.00
-2	0.00	0.00	0.00

Node	Enterococcus_durans
Top Reference	Enterococcus_durans
all reads	512
nonDup	428
readDis	0.963
nonStacked	391
destacking?	Off
C>T_1	0.2672
G>A_1	0
mean length (sd)	58 (17.14)

[illegible]