

2 **Goodness-of-fit for multi-distribution neutrino** 3 **cross-section measurements with shared events**

4 **F. Martínez López and T. Mohayai**

5 *Department of Physics, Indiana University,*
6 *727 E. Third St., Bloomington, IN, USA, 47405*

7 *E-mail:* frmart@iu.edu

8 **ABSTRACT:** When multiple differential cross-section distributions are extracted from a common event
9 sample with correctly evaluated inter-distribution correlations, the statistical covariance matrix is
10 generically singular. Exact linear constraints from event sharing across measurements, such as equal
11 distribution totals when all selected events populate bins in every distribution, produce directions
12 with zero statistical variance. Systematic uncertainties restore formal invertibility but leave near-
13 singular directions that yield unphysically inflated χ^2 values in global goodness-of-fit tests. We
14 present a projection method that restricts the test to the subspace carrying independent statistical
15 information, yielding a χ^2 distributed with $N_{\text{bins}} - N_{\text{null}}$ degrees of freedom, where N_{null} is the
16 number of independent constraints. The method is connected to established results on hypothesis
17 testing with singular covariance matrices and validated with both an analytical toy model and a
18 realistic simulated neutrino–argon cross-section measurement including unfolding and parametrized
19 multi-source systematic uncertainties.

20 **KEYWORDS:** Analysis and statistical methods

21 **ARXIV EPRINT:** [1234.56789](https://arxiv.org/abs/1234.56789)

22 Contents

23	1 Introduction	1
24	2 Rank deficiency of the statistical covariance	2
25	2.1 Statistical covariance from shared events	3
26	2.2 Combination matrix and null space	3
27	2.3 Determination of the structural null space	4
28	2.4 Effect of systematic uncertainties	6
29	3 Range-projected χ^2 statistic	8
30	3.1 Projected chi-squared	8
31	3.2 Proof sketch / justification	8
32	3.3 Relation to alternative approaches	8
33	A Algorithm	8

34 1 Introduction

35 Modern neutrino cross-section literature is experiencing an increase of published measurements
36 reporting multiple differential distributions extracted from the same event sample. MicroBooNE
37 has reported simultaneous measurements of cross sections for final states with and without protons
38 [1, 2], multi-differential QE-like cross section measurements in kinematic imbalance variables
39 [3, 4], and triple-differential inclusive results [5]. Other relevant examples include the multiple
40 final-state kinematic projections from T2K [6, 7], and double-differential inclusive results from
41 NOvA [8–10]. In each of these analyses, the same selected events populate bins in multiple
42 reported distributions, leading to strong inter-measurement statistical correlations. Separate but
43 related efforts have produced simultaneous measurements across different nuclear targets [11–13]
44 and combined neutrino-antineutrino results [14], where correlations arise from shared systematic
45 uncertainties rather than shared events.

46 When multiple distributions are extracted from a common event sample, the event-sharing
47 structure imposes exact linear constraints among the bin counts. If every selected event populates
48 one bin in each distribution, then the total number of events in all of them is exactly identical. More
49 generally, the pattern of shared bin occupancy across distributions determines a set of constraints that
50 reduce the effective dimensionality of the measurement below the total bin count. These constraints
51 are a property of the measurement itself, independent of any particular statistical treatment.

52 The blockwise unfolding framework of Ref. [15] provides a rigorous method for computing
53 the full cross-distribution covariance matrices in such measurements. In this approach, all bins
54 corresponding to the same measured observable in the multi-distribution analysis are organized
55 in blocks, each corresponding to a single differential cross-section distribution. Each bin block

is unfolded independently using any standard method (e.g. iterative D’Agostini unfolding [16] or Wiener-SVD [17]), while the full covariance matrix, encoding the correlations between bins from different distributions, is constructed and propagated through the cross section extraction procedure. The statistical covariance for bins in different blocks is determined by the number of events they share. The constraints described above can manifest as a rank deficiency in the statistical covariance matrix, as certain directions in bin space carry identically zero variance. The number of these degenerate directions is exactly predictable from the binning geometry.

A natural use of these combined measurements is computing global goodness-of-fit metrics, such as a χ^2 statistic, that test a model against all distributions simultaneously. Compared to evaluating each distribution independently, a global test that accounts for all inter-block correlations provides the maximum discriminating power available from the data. Such global metrics are essential for interaction model tuning efforts [18–20], where measurements of multiple kinematic distributions provide complementary sensitivity to different aspects of the underlying physics. However, a global χ^2 that does not account for the constrained directions yields a test with the wrong effective number of degrees of freedom. When the statistical covariance matrix is constructed under the Poisson assumption, the constraints appear as exact null directions that prevent direct matrix inversion. Systematic uncertainties may formally restore invertibility, but the resulting near-singular directions produce unphysically inflated χ^2 values. In either case, the test is unreliable unless the constrained directions are properly identified and excluded.

We present a method to correctly compute a global goodness-of-fit test statistic for multi-distribution measurements with shared events. The approach is based on identifying the constrained directions from the bin structure of the measurement and projecting the statistic onto the subspace where independent statistical information is available. We refer to the resulting test statistic as the range-projected χ^2 . We demonstrate the method using a pedagogical toy example and connect it to the established statistical literature on hypothesis testing with singular covariance matrices.

2 Rank deficiency of the statistical covariance

Consider a neutrino cross-section analysis simultaneously measuring B different flux-integrated differential cross sections, each with N_b true bins. If the measurement is extracted from a common event sample then each selected event will populate exactly one bin in each distribution it belongs to. Recent MicroBooNE analyses [21] have adopted the blockwise unfolding approach presented in Ref. [15] to simultaneous cross-section measurements.

This method provides a systematic framework for reporting the correlations between the different extracted differential cross sections. It provides a general prescription to construct statistical covariance matrices that respect the correlations between bins in different distributions or blocks. The full inter-block covariance matrix constructed in reconstructed space is propagated through unfolding using a block-diagonal unfolding matrix, obtained as a direct sum of the individual unfolding matrices computed for each block. The result of the combined measurement is a vector of $N_{\text{bins}} = \sum_{b=1}^B N_b$ cross-section values with an $N_{\text{bins}} \times N_{\text{bins}}$ covariance matrix.

2.1 Statistical covariance from shared events

The core idea behind the treatment of the statistical correlations in the blockwise unfolding approach is the partition of any arbitrary pair of overlapping kinematic bins X and Y into a set of three non-overlapping bins u , v , and w . The events in the intersection $X \cap Y$ are assigned to bin v , while bin u contains the events in the relative complement of Y in X (i.e. $X \setminus Y$), and bin w corresponds to the relative complement of X in Y (i.e. $Y \setminus X$). Denoting the number of predicted events in each bin as n_i , and assuming these follow independent Poisson distributions, the covariance between the counts in bins X and Y is given by [15]:

$$\begin{aligned} \text{Cov}(n_X, n_Y) &= \text{Cov}(n_u + n_v, n_v + n_w) \\ &= \text{Cov}(n_u, n_v) + \text{Cov}(n_u, n_w) + \text{Cov}(n_v, n_v) + \text{Cov}(n_v, n_w) \\ &= \text{Var}(n_v), \end{aligned} \quad (2.1)$$

as by definition bins u , v , and w have no events in common. This implies that the covariance between any two bins can be estimated from the variance of the events they share.

The variance for a set of Monte Carlo (MC) events can be estimated from the sum of their squared weights. This gives us a general prescription to calculate the elements of the statistical covariance matrix as:

$$(C_{\text{stat}})_{ij} = \sum_{e \in (i \cap j)} w_e^2, \quad (2.2)$$

where w_e is the weight associated to event e belonging to the intersection of bins i and j . In the case of measured data the weights are 1, and the covariance element simplifies to the number of events shared between the bins n_{ij} .

After propagation through the block-diagonal unfolding, the statistical covariance matrix in unfolded space inherits the same rank structure, as the unfolding matrices for each individual block are invertible. This rank structure has a direct interpretation in terms of the number of distinct event-sharing patterns across distributions, which we discuss next.

2.2 Combination matrix and null space

The previous discussion shows that the statistical covariance between any two bins is determined solely by the events they share. We now formalize this observation to derive an exact formula for the rank of C_{stat} , and therefore for the number of directions in bin space that carry zero statistical variance.

For each event e , we can define the indicator vector $\phi_e := (\phi_{e,i})_{i=1}^{N_{\text{bins}}}$ such that $\phi_{e,i} = 1$ if $i \in \text{bins}(e)$ and $\phi_{e,i} = 0$ if $i \notin \text{bins}(e)$, where $\text{bins}(e)$ is the set of bins event e populates across all distributions. The indicator vector acts as an identifier of each event in the combined measurement. For example, in a two-block measurement with 3 bins each, an event populating bin 2 of block 1 and bin 1 of block 2 has indicator vector $\phi_e = (0, 1, 0, 1, 0, 0)$. The statistical covariance matrix can then be re-written in terms of indicator vectors as:

$$C_{\text{stat}} = \sum_e w_e^2 \phi_e \phi_e^T. \quad (2.3)$$

This outer-product decomposition makes the rank structure of C_{stat} clear. The rank of the statistical covariance matrix equals the dimension of $\text{span}\{\phi_e\}$, which depends only on the distinct

indicator vectors present, not on their weights. Since all weights are $w_e > 0$ no indicator vector is removed from the sum, and the column space of C_{stat} is exactly $\text{span}\{\phi_e\}$.

We define the bin-combination matrix Ω as the matrix whose rows are the unique indicator vectors among $\{\phi_e\}$. Each row of Ω represent a distinct pattern of bin occupancy across the blocks. By construction the row space of Ω equals $\text{span}\{\phi_e\}$, and therefore:

$$\text{rank}(C_{\text{stat}}) = \text{rank}(\Omega). \quad (2.4)$$

The number of degenerate directions is then:

$$N_{\text{null}} = N_{\text{bins}} - \text{rank}(\Omega). \quad (2.5)$$

Physically, each of the N_{null} directions corresponds to a linear combination of bins along which the measurement has identically zero variance. Any component of the residual vector along such a direction is not perceived by the statistical covariance; it is a direction that the measurement has no statistical power to constrain. Importantly, both Ω and N_{null} are directly computable from the MC event-to-bin mapping alone, before constructing any covariance matrix.

A vector $\mathbf{x}$ is in the null space of C_{stat} if and only if, for every event e :

$$\sum_{i \in \text{bins}(e)} x_i = 0, \quad (2.6)$$

or equivalently $\Omega \mathbf{x} = 0$. Each unique indicator vector acts as a linear constraint on the null-space vectors, requiring that the components of $\mathbf{x}$ sum to zero over the bins populated by the different events. The null space directions can be divided into two physically distinct categories: structural constraints that follow from the block topology alone, and kinematic constraints that depend on which regions of phase space are populated.

2.3 Determination of the structural null space

The combination matrix Ω presented previously is constructed from MC events. One determines which events populate the different bins and collects the unique indicator vectors. However, the indicator vectors that can appear are constrained by two distinct mechanisms. The first is purely structural, the geometry of the binning scheme forbids certain bin combinations. The second is kinematic, correlations between the different variables render some combinations unreachable even though they are geometrically allowed. These constraints are imposed by the underlying physics effects, mainly the differential cross section and the detector acceptance. In the following, we show that the structural constraints can be determined a priori, without generating MC events, from the bin edges used.

The variables used in a measurement are drawn from a set $\{v_1, \dots, v_D\}$. Each block b uses a subset of these variables, $\text{vars}(b) \subseteq \{v_1, \dots, v_D\}$. Restricting the discussion to combinations of single- and double-differential measurements, we have $|\text{vars}(b)| \in \{1, 2\}$ for every b . For each variable v used in a given block b , the binning scheme defines a set of bin edges $E_v^{(b)}$ that partition the range of v into intervals. In the case of a two-dimensional block, the edges in the binning variable may be slice-dependent, but nevertheless these can be collected into a single set of edges

per variable across slices. For each variable $v \in \bigcup_b \text{vars}(b)$, we can define the refined edges:

$$\mathcal{E}_v = \bigcup_{b: v \in \text{vars}(b)} E_v^{(b)}, \quad (2.7)$$

with $M_v = |\mathcal{E}_v| - 1$ denoting the number of refined bins for variable v . Since $\mathcal{E}_v$ contains all the bin edges from every block that uses that variable, each refined bin is contained entirely within one original bin of every such block.

The common refinement of all blocks is the Cartesian product of the refined bins across all variables:

$$\mathcal{R} = \bigotimes_v \{1, \dots, M_v\}. \quad (2.8)$$

Each cell $c \in \mathcal{R}$ specifies a refined bin index for every variable in the measurement. For every block b , there exists a unique mapping $\beta_b(c)$ between each cell c and an original bin in said block. The elements of the indicator vector $\phi_c := (\phi_{c,i})_{i=1}^{N_{\text{bins}}}$ associated to cell c can be written as:

$$\phi_{c,i} = \begin{cases} 1 & \text{if } i = \beta_b(c) \text{ for any block } b, \\ 0 & \text{otherwise.} \end{cases} \quad (2.9)$$

If block b does not use variable v then β_b is independent of the refined bin index in v . Therefore, multiple cells may produce identical indicator vectors.

The structural combination matrix Ω_{struct} is defined as the matrix whose rows are the unique indicator vectors from the cells of the common refinement $\{\phi_c : c \in \mathcal{R}\}$. By construction, every indicator vector that appears in the Ω matrix obtained from the MC events is also present in Ω_{struct} , therefore:

$$\text{rank}(\Omega) \leq \text{rank}(\Omega_{\text{struct}}) \leq N_{\text{bins}}. \quad (2.10)$$

This hierarchy motivates a decomposition of the total number of degenerate directions:

$$\begin{aligned} N_{\text{null}} &= N_{\text{bins}} - \text{rank}(\Omega_{\text{struct}}) + \text{rank}(\Omega_{\text{struct}}) - \text{rank}(\Omega) \\ &= N_{\text{null}}^{\text{struct}} + N_{\text{null}}^{\text{kinem}}. \end{aligned} \quad (2.11)$$

The first term, $N_{\text{null}}^{\text{struct}}$, counts null directions that arise purely from the binning geometry and is computable a priori from the bin edges alone. The second term, $N_{\text{null}}^{\text{kinem}}$, accounts for additional null directions introduced by phase-space restrictions that exclude geometrically allowed combinations and can only be determined from Monte Carlo. A description of the algorithm used to compute Ω_{struct} , including an optimization that restricts the enumeration to shared variables, is presented in Appendix A.

The structure of the null space of Ω_{struct} depends on how the bin edges for the different blocks overlap. In the simplest case, when blocks do not share variables, every combination of one bin per block is allowed. The resulting null space is $(B - 1)$ -dimensional, with a single constraint that requires all block totals to be equal. When blocks share one or more variables, the allowed combinations are restricted and the dimension of the null space can be different from $B - 1$. The number of null directions depends on how shared variable bin edges are aligned between blocks. Aligned slice edges decouple the measurement into different sub-problems, adding additional sum

rules and increasing $N_{\text{null}}^{\text{struct}}$. On the other hand, misaligned edges couple adjacent distributions, potentially reducing the number of constraints. In the general case, the interplay between shared variables, bin edge alignment, and block topology makes a closed-form expression impractical, but the computation of $\text{rank}(\Omega_{\text{struct}})$ from the bin edge specifications is straightforward.

The structural and kinematic components of the null space respond differently to systematic uncertainties. Whether these cause the rank deficiency of C_{stat} to persist, partially resolve, or fully resolve in the total covariance C_{total} depends on the nature of the systematic treatment, which we discuss next.

2.4 Effect of systematic uncertainties

Previously, we established that C_{stat} is generically singular, with $N_{\text{null}} = N_{\text{bins}} - \text{rank}(\Omega)$ degenerate directions that can be separated into structural and kinematic contributions. Whether the total covariance $C_{\text{total}} = C_{\text{stat}} + C_{\text{syst}}$ inherits the rank deficiency depends on which of these null directions receive support from the systematic uncertainties. Now, we examine the mechanisms by which different systematic treatments do or do not populate the null space of C_{stat} .

A systematic variation s produces a modified vector of expected events $\mathbf{n}_s$, whose departure from the central-value prediction can be expressed in terms of a shift vector:

$$\delta_s = \mathbf{n}_s - \mathbf{n}_{\text{CV}}. \quad (2.12)$$

The different contributions to C_{syst} are constructed from the outer product of these shift vectors. A null direction $\mathbf{x}$ of C_{stat} remains a null direction of C_{total} if and only if $\mathbf{x} \cdot \delta_s = 0$ for every systematic variation in the uncertainty budget. The question is therefore whether the shift vectors have nonzero projections onto the structural and kinematic null spaces.

Consider first the simplest case, where a systematic s acts by reweighting the reconstructed event counts directly. Each event e is assigned a modified weight $w_e^{(s)}$ while preserving its bin assignments. The shift vector can then be written as:

$$\delta_s = \sum_e (w_e^{(s)} - w_e) \phi_e. \quad (2.13)$$

For any null vector $\mathbf{x}$ which satisfies the null-space condition of Eq. 2.6, the projection of $\mathbf{x}$ onto the shift vector gives:

$$\begin{aligned} \mathbf{x} \cdot \delta_s &= \sum_e (w_e^{(s)} - w_e) (\phi_e \cdot \mathbf{x}) \\ &= \sum_e (w_e^{(s)} - w_e) \sum_{i=1}^{N_{\text{bins}}} \phi_{e,i} x_i \\ &= 0, \end{aligned} \quad (2.14)$$

since the inner sum vanishes for every event. Direct reweighting therefore preserves the entire null space, adding no rank to C_{total} . Flux uncertainties, which modify the neutrino energy spectrum through per-event weights, are the classic example. Cross-section model variations and hadronic reinteraction uncertainties, when implemented as per-event reweighting, behave identically.

218 A qualitatively different situation arises when the systematic uncertainties are propagated
 219 through the response matrix rather than changing the event counts per bin directly. In this approach,
 220 adopted as the standard prescription in MicroBooNE for neutrino interaction uncertainties [15], a
 221 modified response matrix R'_s , mapping true bins to reconstructed bins under the given variation,
 222 is constructed for each systematic uncertainty. This matrix is obtained either by reweighting the
 223 events used to build the nominal response matrix or from an independent MC sample with varied
 224 simulation parameters. In this case, the predicted reconstructed counts for systematic variation s
 225 are:

$$\mathbf{n}_s = R'_s \mathbf{t}_{\text{CV}}, \quad (2.15)$$

226 where $\mathbf{t}_{\text{CV}}$ is the central-value truth prediction. The shift vector is then:

$$\delta_s = (R'_s - R) \mathbf{t}_{\text{CV}}. \quad (2.16)$$

227 Since R is block-diagonal, the difference $R'_s - R$ acts independently on each block, projecting
 228 the nominal truth vector through block-specific changes in efficiency and migration. The per-event
 229 coupling that makes direct reweighting respect all null directions is absent in this prescription. When
 230 the weight of an event changes, it affects simultaneously every block it populates; the response-
 231 matrix method, in contrast, processes each block independently through its own sub-matrix. The
 232 projection of δ_s onto any null vector, whether structural or kinematic, therefore involves differences
 233 in how each block responds to the same truth-level prediction, and in general is nonzero. Response-
 234 matrix variations thus break both types of null directions, regardless of how R'_s is obtained.

235 A third mechanism applies when systematic uncertainties are evaluated by directly comparing
 236 reconstructed event counts from independent MC samples simulated with nominal and varied
 237 parameters. In this case, the shift vector takes the form:

$$\delta_s = \sum_{e'} w_{e'}^{(s)} \phi_{e'}^{(s)} - \sum_e w_e \phi_e, \quad (2.17)$$

238 where the first sum runs over the events in the varied-model sample and the second sum over
 239 the nominal sample. Since each event in both samples populates exactly one bin per block,
 240 their difference projects to zero onto the structural null directions by an analogous argument to
 241 that used for the direct reweighting. Kinematic null directions, however, can be lifted if the
 242 variation scatters events into reconstructed-bin combinations absent in the nominal sample. In that
 243 case, new indicator vectors appear that are not rows of Ω , potentially breaking the corresponding
 244 constraints. Detector systematic uncertainties, which usually require independent samples with
 245 modified simulation parameters that cannot be captured by event reweighting, are the typical
 246 example where this prescription is applied.

247 In practice, whether the eigenvalues of C_{total} along the formerly null directions are exactly zero
 248 or merely smaller than the physical eigenvalues, including these directions in a χ^2 test via standard
 249 matrix inversion produces inflated values with the wrong effective number of degrees of freedom.
 250 A valid goodness-of-fit test must therefore be restricted to the subspace where the measurement
 251 carries genuine sensitivity, which is the role of the projection method developed in the following
 252 section.

3 Range-projected χ^2 statistic

The analysis of Sec. 2 established that the statistical covariance matrix C_{stat} is in general rank-deficient, with N_{null} directions along which the measurement carries no statistical sensitivity. A global goodness-of-fit must therefore be formulated in the subspace where independent statistical information is available.

The rank structure discussed previously was established at the level of reconstructed event counts. The blockwise unfolding procedure of Ref. [15] propagates the full covariance matrix through the block-diagonal unfolding matrix, which is invertible by construction.

the SVD decomposition of the statistical covariance matrix as:

$$C_{\text{stat}} = U \Sigma U^T, \quad (3.1)$$

and that $r = \text{rank}(C_{\text{stat}})$. partitionings of $U = [U_r | \bar{U}_r]$

$$\begin{aligned} U_r U_r^T &= \text{projection on to } \text{ran}(C_{\text{stat}}) \\ \bar{U}_r \bar{U}_r^T &= \text{projection on to } \text{null}(C_{\text{stat}}) \end{aligned} \quad (3.2)$$

3.1 Projected chi-squared

3.2 Proof sketch / justification

3.3 Relation to alternative approaches

Acknowledgments

A Algorithm

The naive counting of the common refinement $\mathcal{R}$ has cost $\prod_v M_v$, which grows exponentially with the number of kinematic variables. A typical multi-distribution measurement involves $O(5)$ variables each with $O(10)$ refined bins, giving $O(10^5)$ cells. The key insight is that only shared variables (i.e. the ones appearing in two or more blocks) generate the cross-block constraints. Dividing the variable set into shared (V_s) and private (V_p) variables, one has to enumerate only the reduced refinement $\prod_{v \in V_s} M_v$. For each of the shared-variable cells, the bins from blocks that depend on the shared variable are determined by lookup, while blocks whose binning variable is private contribute their bins independently via a local Cartesian product. If the number of shared variables is small the computational cost is dramatically reduced.

References

- [1] MICROBooNE collaboration, *First Simultaneous Measurement of Differential Muon-Neutrino Charged-Current Cross Sections on Argon for Final States with and Without Protons Using MicroBooNE Data*, *Phys. Rev. Lett.* **133** (2024) 041801 [[2402.19281](#)].
- [2] MICROBooNE collaboration, *Inclusive cross section measurements in final states with and without protons for charged-current ν_μ -Ar scattering in MicroBooNE*, *Phys. Rev. D* **110** (2024) 013006 [[2402.19216](#)].

- [3] MICROBooNE collaboration, *First Double-Differential Measurement of Kinematic Imbalance in Neutrino Interactions with the MicroBooNE Detector*, *Phys. Rev. Lett.* **131** (2023) 101802 [2301.03706].
- [4] MICROBooNE collaboration, *Multidifferential cross section measurements of $\nu\mu$ -argon quasielasticlike reactions with the MicroBooNE detector*, *Phys. Rev. D* **108** (2023) 053002 [2301.03700].
- [5] MICROBooNE collaboration, *Measurement of three-dimensional inclusive muon-neutrino charged-current cross sections on argon with the MicroBooNE detector*, *Phys. Lett. B* **870** (2025) 139939 [2307.06413].
- [6] T2K collaboration, *Characterization of nuclear effects in muon-neutrino scattering on hydrocarbon with a measurement of final-state kinematics and correlations in charged-current pionless interactions at T2K*, *Phys. Rev. D* **98** (2018) 032003 [1802.05078].
- [7] T2K collaboration, *First T2K measurement of transverse kinematic imbalance in the muon-neutrino charged-current single- π^+ production channel containing at least one proton*, *Phys. Rev. D* **103** (2021) 112009 [2102.03346].
- [8] NOvA collaboration, *Measurement of the double-differential muon-neutrino charged-current inclusive cross section in the NOvA near detector*, *Phys. Rev. D* **107** (2023) 052011 [2109.12220].
- [9] NOvA collaboration, *Measurement of $d^2\sigma/d|\vec{q}|dE_{avail}$ in charged current ν_μ -nucleus interactions at $\langle E_\nu \rangle = 1.86$ GeV using the NOvA Near Detector*, *Phys. Rev. D* **111** (2025) 052009 [2410.05526].
- [10] NOvA collaboration, “Measurement of the double-differential cross section of muon-neutrino charged-current interactions with low hadronic energy in the NOvA Near Detector.” 10, 2024.
- [11] T2K collaboration, *Simultaneous measurement of the muon neutrino charged-current cross section on oxygen and carbon without pions in the final state at T2K*, *Phys. Rev. D* **101** (2020) 112004 [2004.05434].
- [12] MINERvA collaboration, *Simultaneous Measurement of ν_μ Quasielasticlike Cross Sections on CH, C, H₂O, Fe, and Pb as a Function of Muon Kinematics at MINERvA*, *Phys. Rev. Lett.* **130** (2023) 161801 [2301.02272].
- [13] MINERvA collaboration, *Simultaneous Measurement of Muon Neutrino $\nu\mu$ Charged-Current Single π^+ Production in CH, C, H₂O, Fe, and Pb Targets in MINERvA*, *Phys. Rev. Lett.* **131** (2023) 011801 [2209.07852].
- [14] T2K collaboration, *First combined measurement of the muon neutrino and antineutrino charged-current cross section without pions in the final state at T2K*, *Phys. Rev. D* **101** (2020) 112001 [2002.09323].
- [15] S. Gardiner, “Mathematical methods for neutrino cross-section extraction.” 2024.
- [16] G. D’Agostini, *A Multidimensional unfolding method based on Bayes’ theorem*, *Nucl. Instrum. Meth. A* **362** (1995) 487.
- [17] W. Tang, X. Li, X. Qian, H. Wei and C. Zhang, *Data Unfolding with Wiener-SVD Method*, *JINST* **12** (2017) P10002 [1705.03568].
- [18] C. Wilkinson et al., *Testing charged current quasi-elastic and multinucleon interaction models in the NEUT neutrino interaction generator with published datasets from the MiniBooNE and MINERvA experiments*, *Phys. Rev. D* **93** (2016) 072010 [1601.05592].

- 325 [19] MINERvA collaboration, *Tuning the GENIE Pion Production Model with MINERvA Data*, *Phys. Rev.*
326 *D* **100** (2019) 072005 [[1903.01558](#)].
- 327 [20] GENIE collaboration, *Neutrino-nucleus $CC0\pi$ cross-section tuning in GENIE v3*, *Phys. Rev. D* **106**
328 (2022) 112001 [[2206.11050](#)].
- 329 [21] MICROBooNE collaboration, *Measurement of single- and double-differential cross sections for*
330 *mesonless charged-current muon neutrino interactions on argon with final-state protons using the*
331 *MicroBooNE detector*, *Phys. Rev. D* **112** (2025) 112004 [[2403.19574](#)].