

2 **Goodness-of-fit for multi-distribution neutrino** 3 **cross-section measurements with shared events**

4 **F. Martínez López and T. Mohayai**

5 *Department of Physics, Indiana University,*
6 *727 E. Third St., Bloomington, IN, USA, 47405*

7 *E-mail:* frmart@iu.edu

8 **ABSTRACT:** When multiple differential cross-section distributions are extracted from a common event
9 sample with correctly evaluated inter-distribution correlations, the statistical covariance matrix is
10 generically singular. Exact linear constraints from event sharing across measurements, such as equal
11 distribution totals when all selected events populate bins in every distribution, produce directions
12 with zero statistical variance. Systematic uncertainties restore formal invertibility but leave near-
13 singular directions that yield unphysically inflated χ^2 values in global goodness-of-fit tests. We
14 present a projection method that restricts the test to the subspace carrying independent statistical
15 information, yielding a χ^2 distributed with $N_{\text{bins}} - N_{\text{null}}$ degrees of freedom, where N_{null} is the
16 number of independent constraints. The method is connected to established results on hypothesis
17 testing with singular covariance matrices and validated with both an analytical toy model and a
18 realistic simulated neutrino–argon cross-section measurement including unfolding and parametrized
19 multi-source systematic uncertainties.

20 **KEYWORDS:** Analysis and statistical methods

21 **ARXIV EPRINT:** [1234.56789](https://arxiv.org/abs/1234.56789)

22 Contents

23	1 Introduction	1
24	2 Rank deficiency of the statistical covariance	2
25	2.1 Statistical covariance from shared events	2
26	2.2 Combination matrix and null space	3
27	2.3 Determination of the structural null space	4
28	2.4 Effect of systematic uncertainties	6
29	3 Range-projected χ^2 statistic	7
30	3.1 Projected chi-squared	8
31	3.2 Proof sketch / justification	8
32	3.3 Relation to alternative approaches	8
33	A Algorithm	8

34 1 Introduction

35 Modern neutrino cross-section literature is experiencing an increase of published measurements
36 reporting multiple differential distributions extracted from the same event sample. MicroBooNE
37 has reported simultaneous measurements of cross sections for final states with and without protons
38 [1, 2], multi-differential QE-like cross section measurements in kinematic imbalance variables
39 [3, 4], and triple-differential inclusive results [5]. Other relevant examples include the multiple
40 final-state kinematic projections from T2K [6, 7], and double-differential inclusive results from
41 NOvA [8–10]. In each of these analyses, the same selected events populate bins in multiple
42 reported distributions, leading to strong inter-measurement statistical correlations. Separate but
43 related efforts have produced simultaneous measurements across different nuclear targets [11–13]
44 and combined neutrino-antineutrino results [14], where correlations arise from shared systematic
45 uncertainties rather than shared events.

46 The blockwise unfolding framework [15] provides a rigorous method for computing the full
47 cross-distribution covariance matrices in all such measurements. In this approach, all bins cor-
48 responding to the same measured observable in the multi-distribution analysis are organized in
49 blocks, each corresponding to a single differential cross-section distribution. Each bin block is
50 unfolded independently using any standard method (e.g. iterative D’Agostini unfolding [16] or
51 Wiener-SVD [17]), while the full covariance matrix, encoding the correlations between bins from
52 different distributions, is constructed and propagated through the cross section extraction proce-
53 dure. The statistical covariance for bins in different blocks is determined by the number of events
54 they share. The result is a single combined covariance matrix covering all reported bins across all
55 distributions.

A natural use of these combined measurements is computing global goodness-of-fit metrics, such as a χ^2 statistic, that test a model against all distributions simultaneously. Compared to evaluating each distribution independently, a global test that accounts for all inter-block correlations provides the maximum discriminating power available from the data. Such global metrics are essential for interaction model tuning efforts [18–20], where measurements of multiple kinematic distributions provide complementary sensitivity to different aspects of the underlying physics. However, as we show in this paper, the full inter-block statistical covariance is generically singular when it correctly accounts for correlations from shared events. The rank deficiency arises from exact linear constraints imposed by the event-sharing structure of the binning scheme: certain linear combinations of bins, such as the difference in total event counts between two distributions of the same events, have identically zero variance. While systematic uncertainties formally restore invertibility, the resulting near-singular directions produce unphysically inflated χ^2 values, yielding a global test with the wrong effective number of degrees of freedom.

We present a method based on projecting onto the range (column space) of the statistical covariance matrix that yields a well-defined chi-squared with the correct number of degrees of freedom. We demonstrate the method using a pedagogical toy example and connect it to the established statistical literature on hypothesis testing with singular covariance matrices.

Brief roadmap of sections.

2 Rank deficiency of the statistical covariance

Consider a neutrino cross-section analysis simultaneously measuring B different flux-integrated differential cross sections, each with N_b true bins. If the measurement is extracted from a common event sample then each selected event will populate exactly one bin in each distribution it belongs to. Recent MicroBooNE analyses [21] have adopted the blockwise unfolding approach presented in Ref. [15] to simultaneous cross-section measurements.

This method provides a systematic framework for reporting the correlations between the different extracted differential cross sections. It provides a general prescription to construct statistical covariance matrices that respect the correlations between bins in different distributions or blocks. The full inter-block covariance matrix constructed in reconstructed space is propagated through unfolding using a block-diagonal unfolding matrix, obtained as a direct sum of the individual unfolding matrices computed for each block. The result of the combined measurement is a vector of $N_{\text{bins}} = \sum_{b=1}^B N_b$ cross-section values with an $N_{\text{bins}} \times N_{\text{bins}}$ covariance matrix.

2.1 Statistical covariance from shared events

The core idea behind the treatment of the statistical correlations in the blockwise unfolding approach is the partition of any arbitrary pair of overlapping kinematic bins X and Y into a set of three non-overlapping bins u , v , and w . The events in the intersection $X \cap Y$ are assigned to bin v , while bin u contains the events in the relative complement of Y in X (i.e. $X \setminus Y$), and bin w corresponds to the relative complement of X in Y (i.e. $Y \setminus X$). Denoting the number of predicted events in each bin as n_i , and assuming these follow independent Poisson distributions, the covariance between the

counts in bins X and Y is given by [15]:

$$\begin{aligned} \text{Cov}(n_X, n_Y) &= \text{Cov}(n_u + n_v, n_v + n_w) \\ &= \text{Cov}(n_u, n_v) + \text{Cov}(n_u, n_w) + \text{Cov}(n_v, n_v) + \text{Cov}(n_v, n_w) \\ &= \text{Var}(n_v), \end{aligned} \quad (2.1)$$

as by definition bins u , v , and w have no events in common. This implies that the covariance between any two bins can be estimated from the variance of the events they share.

The variance for a set of Monte Carlo (MC) events can be estimated from the sum of their squared weights. This gives us a general prescription to calculate the elements of the statistical covariance matrix as:

$$(C_{\text{stat}})_{ij} = \sum_{e \in (i \cap j)} w_e^2, \quad (2.2)$$

where w_e is the weight associated to event e belonging to the intersection of bins i and j . In the case of measured data the weights are 1, and the covariance element simplifies to the number of events shared between the bins n_{ij} .

After propagation through the block-diagonal unfolding, the statistical covariance matrix in unfolded space inherits the same rank structure, as the unfolding matrices for each individual block are invertible. This rank structure has a direct interpretation in terms of the number of distinct event-sharing patterns across distributions, which we discuss next.

2.2 Combination matrix and null space

The previous discussion shows that the statistical covariance between any two bins is determined solely by the events they share. We now formalize this observation to derive an exact formula for the rank of C_{stat} , and therefore for the number of directions in bin space that carry zero statistical variance.

For each event e , we can define the indicator vector $\phi_e := (\phi_{e,i})_{i=1}^{N_{\text{bins}}}$ such that $\phi_{e,i} = 1$ if $i \in \text{bins}(e)$ and $\phi_{e,i} = 0$ if $i \notin \text{bins}(e)$, where $\text{bins}(e)$ is the set of bins event e populates across all distributions. The indicator vector acts as an identifier of each event in the combined measurement. For example, in a two-block measurement with 3 bins each, an event populating bin 2 of block 1 and bin 1 of block 2 has indicator vector $\phi_e = (0, 1, 0, 1, 0, 0)$. The statistical covariance matrix can then be re-written in terms of indicator vectors as:

$$C_{\text{stat}} = \sum_e w_e^2 \phi_e \phi_e^T. \quad (2.3)$$

This outer-product decomposition makes the rank structure of C_{stat} clear. The rank of the statistical covariance matrix equals the dimension of $\text{span}\{\phi_e\}$, which depends only on the distinct indicator vectors present, not on their weights. Since all weights are $w_e > 0$ no indicator vector is removed from the sum, and the column space of C_{stat} is exactly $\text{span}\{\phi_e\}$.

We define the bin-combination matrix Ω as the matrix whose rows are the unique indicator vectors among $\{\phi_e\}$. Each row of Ω represent a distinct pattern of bin occupancy across the blocks. By construction the row space of Ω equals $\text{span}\{\phi_e\}$, and therefore:

$$\text{rank}(C_{\text{stat}}) = \text{rank}(\Omega). \quad (2.4)$$

125 The number of degenerate directions is then:

$$N_{\text{null}} = N_{\text{bins}} - \text{rank}(\Omega). \quad (2.5)$$

126 Physically, each of the N_{null} directions corresponds to a linear combination of bins along which
 127 the measurement has identically zero variance. Any component of the residual vector along such a
 128 direction is not perceived by the statistical covariance; it is a direction that the measurement has no
 129 statistical power to constrain. Importantly, both Ω and N_{null} are directly computable from the MC
 130 event-to-bin mapping alone, before constructing any covariance matrix.

131 A vector $\mathbf{x}$ is in the null space of C_{stat} if and only if, for every event e :

$$\sum_{i \in \text{bins}(e)} x_i = 0, \quad (2.6)$$

132 or equivalently $\Omega \mathbf{x} = 0$. Each unique indicator vector acts as a linear constraint on the null-space
 133 vectors, requiring that the components of $\mathbf{x}$ sum to zero over the bins populated by the different
 134 events. The null space directions can be divided into two physically distinct categories: structural
 135 constraints that follow from the block topology alone, and kinematic constraints that depend on
 136 which regions of phase space are populated.

137 2.3 Determination of the structural null space

138 The combination matrix Ω presented previously is constructed from MC events. One determines
 139 which events populate the different bins and collects the unique indicator vectors. However, the
 140 indicator vectors that can appear are constrained by two distinct mechanisms. The first is purely
 141 structural, the geometry of the binning scheme forbids certain bin combinations. The second is
 142 kinematic, correlations between the different variables render some combinations unreachable even
 143 though they are geometrically allowed. These constraints are imposed by the underlying physics
 144 effects, mainly the differential cross section and the detector acceptance. In the following, we show
 145 that the structural constraints can be determined a priori, without generating MC events, from the
 146 bin edges used.

147 The variables used in a measurement are drawn from a set $\{v_1, \dots, v_D\}$. Each block b uses
 148 a subset of these variables, $\text{vars}(b) \subseteq \{v_1, \dots, v_D\}$. Restricting the discussion to combinations of
 149 single- and double-differential measurements, we have $|\text{vars}(b)| \in \{1, 2\}$ for every b . For each
 150 variable v used in a given block b , the binning scheme defines a set of bin edges $E_v^{(b)}$ that partition
 151 the range of v into intervals. In the case of a two-dimensional block, the edges in the binning
 152 variable may be slice-dependent, but nevertheless these can be collected into a single set of edges
 153 per variable across slices. For each variable $v \in \bigcup_b \text{vars}(b)$, we can define the refined edges:

$$\mathcal{E}_v = \bigcup_{b: v \in \text{vars}(b)} E_v^{(b)}, \quad (2.7)$$

154 with $M_v = |\mathcal{E}_v| - 1$ denoting the number of refined bins for variable v . Since $\mathcal{E}_v$ contains all the
 155 bin edges from every block that uses that variable, each refined bin is contained entirely within one
 156 original bin of every such block.

157 The common refinement of all blocks is the Cartesian product of the refined bins across all
 158 variables:

$$\mathcal{R} = \bigtimes_v \{1, \dots, M_v\}. \quad (2.8)$$

159 Each cell $c \in \mathcal{R}$ specifies a refined bin index for every variable in the measurement. For every block
 160 b , there exists a unique mapping $\beta_b(c)$ between each cell c and an original bin in said block. The
 161 elements of the indicator vector $\phi_c := (\phi_{c,i})_{i=1}^{N_{\text{bins}}}$ associated to cell c can be written as:

$$\phi_{c,i} = \begin{cases} 1 & \text{if } i = \beta_b(c) \text{ for any block } b, \\ 0 & \text{otherwise.} \end{cases} \quad (2.9)$$

162 If block b does not use variable v then β_b is independent of the refined bin index in v . Therefore,
 163 multiple cells may produce identical indicator vectors.

164 The structural combination matrix Ω_{struct} is defined as the matrix whose rows are the unique
 165 indicator vectors from the cells of the common refinement $\{\phi_c : c \in \mathcal{R}\}$. By construction, every
 166 indicator vector that appears in the Ω matrix obtained from the MC events is also present in Ω_{struct} ,
 167 therefore:

$$\text{rank}(\Omega) \leq \text{rank}(\Omega_{\text{struct}}) \leq N_{\text{bins}}. \quad (2.10)$$

168 This hierarchy motivates a decomposition of the total number of degenerate directions:

$$\begin{aligned} N_{\text{null}} &= N_{\text{bins}} - \text{rank}(\Omega_{\text{struct}}) + \text{rank}(\Omega_{\text{struct}}) - \text{rank}(\Omega) \\ &= N_{\text{null}}^{\text{struct}} + N_{\text{null}}^{\text{kinem}}. \end{aligned} \quad (2.11)$$

169 The first term, $N_{\text{null}}^{\text{struct}}$, counts null directions that arise purely from the binning geometry and is
 170 computable a priori from the bin edges alone. The second term, $N_{\text{null}}^{\text{kinem}}$, accounts for additional null
 171 directions introduced by phase-space restrictions that exclude geometrically allowed combinations
 172 and can only be determined from Monte Carlo. A description of the algorithm used to compute
 173 Ω_{struct} , including an optimization that restricts the enumeration to shared variables, is presented in
 174 Appendix A.

175 The structure of the null space of Ω_{struct} depends on how the bin edges for the different blocks
 176 overlap. In the simplest case, when blocks do not share variables, every combination of one bin
 177 per block is allowed. The resulting null space is $(B - 1)$ -dimensional, with a single constraint
 178 that requires all block totals to be equal. When blocks share one or more variables, the allowed
 179 combinations are restricted and the dimension of the null space can be different from $B - 1$. The
 180 number of null directions depends on how shared variable bin edges are aligned between blocks.
 181 Aligned slice edges decouple the measurement into different sub-problems, adding additional sum
 182 rules and increasing $N_{\text{null}}^{\text{struct}}$. On the other hand, misaligned edges couple adjacent distributions,
 183 potentially reducing the number of constraints. In the general case, the interplay between shared
 184 variables, bin edge alignment, and block topology makes a closed-form expression impractical, but
 185 the computation of $\text{rank}(\Omega_{\text{struct}})$ from the bin edge specifications is straightforward.

186 The structural and kinematic components of the null space respond differently to systematic
 187 uncertainties. Whether these cause the rank deficiency of C_{stat} to persist, partially resolve, or fully
 188 resolve in the total covariance C_{total} depends on the nature of the systematic treatment, which we
 189 discuss next.

2.4 Effect of systematic uncertainties

Previously, we established that C_{stat} is generically singular, with $N_{\text{null}} = N_{\text{bins}} - \text{rank}(\Omega)$ degenerate directions that can be separated into structural and kinematic contributions. Whether the total covariance $C_{\text{total}} = C_{\text{stat}} + C_{\text{syst}}$ inherits the rank deficiency depends on which of these null directions receive support from the systematic uncertainties. Now, we examine the mechanisms by which different systematic treatments do or do not populate the null space of C_{stat} .

A systematic variation s produces a modified vector of expected events $\mathbf{n}_s$, whose departure from the central-value prediction can be expressed in terms of a shift vector:

$$\delta_s = \mathbf{n}_s - \mathbf{n}_{\text{CV}}. \quad (2.12)$$

The different contributions to C_{syst} are constructed from the outer product of these shift vectors. A null direction $\mathbf{x}$ of C_{stat} remains a null direction of C_{total} if and only if $\mathbf{x} \cdot \delta_s = 0$ for every systematic variation in the uncertainty budget. The question is therefore whether the shift vectors have nonzero projections onto the structural and kinematic null spaces.

Consider first the simplest case, where a systematic s acts by reweighting the reconstructed event counts directly. Each event e is assigned a modified weight $w_e^{(s)}$ while preserving its bin assignments. The shift vector can then be written as:

$$\delta_s = \sum_e (w_e^{(s)} - w_e) \phi_e. \quad (2.13)$$

For any null vector $\mathbf{x}$ which satisfies the null-space condition of Eq. 2.6, the projection of $\mathbf{x}$ onto the shift vector gives:

$$\begin{aligned} \mathbf{x} \cdot \delta_s &= \sum_e (w_e^{(s)} - w_e) (\phi_e \cdot \mathbf{x}) \\ &= \sum_e (w_e^{(s)} - w_e) \sum_{i=1}^{N_{\text{bins}}} \phi_{e,i} x_i \\ &= 0, \end{aligned} \quad (2.14)$$

since the inner sum vanishes for every event. Direct reweighting therefore preserves the entire null space, adding no rank to C_{total} . Flux uncertainties, which modify the neutrino energy spectrum through per-event weights, are the classic example. Cross-section model variations and hadronic reinteraction uncertainties, when implemented as per-event reweighting, behave identically.

A qualitatively different situation arises when the systematic uncertainties are propagated through the response matrix rather than changing the event counts per bin directly. In this approach, adopted as the standard prescription in MicroBooNE for neutrino interaction uncertainties [15], a modified response matrix R'_s , mapping true bins to reconstructed bins under the given variation, is constructed for each systematic uncertainty. This matrix is obtained either by reweighting the events used to build the nominal response matrix or from an independent MC sample with varied simulation parameters. In this case, the predicted reconstructed counts for systematic variation s are:

$$\mathbf{n}_s = R'_s \mathbf{t}_{\text{CV}}, \quad (2.15)$$

where $\mathbf{t}_{\text{CV}}$ is the central-value truth prediction. The shift vector is then:

$$\delta_s = (R'_s - R) \mathbf{t}_{\text{CV}}. \quad (2.16)$$

Since R is block-diagonal, the difference $R'_s - R$ acts independently on each block, projecting the nominal truth vector through block-specific changes in efficiency and migration. The per-event coupling that makes direct reweighting respect all null directions is absent in this prescription. When the weight of an event changes, it affects simultaneously every block it populates; the response-matrix method, in contrast, processes each block independently through its own sub-matrix. The projection of δ_s onto any null vector, whether structural or kinematic, therefore involves differences in how each block responds to the same truth-level prediction, and in general is nonzero. Response-matrix variations thus break both types of null directions, regardless of how R'_s is obtained.

A third mechanism applies when systematic uncertainties are evaluated by directly comparing reconstructed event counts from independent MC samples simulated with nominal and varied parameters. In this case, the shift vector takes the form:

$$\delta_s = \sum_{e'} w_{e'}^{(s)} \phi_{e'}^{(s)} - \sum_e w_e \phi_e, \quad (2.17)$$

where the first sum runs over the events in the varied-model sample and the second sum over the nominal sample. Since each event in both samples populates exactly one bin per block, their difference projects to zero onto the structural null directions by an analogous argument to that used for the direct reweighting. Kinematic null directions, however, can be lifted if the variation scatters events into reconstructed-bin combinations absent in the nominal sample. In that case, new indicator vectors appear that are not rows of Ω , potentially breaking the corresponding constraints. Detector systematic uncertainties, which usually require independent samples with modified simulation parameters that cannot be captured by event reweighting, are the typical example where this prescription is applied.

In practice, whether the eigenvalues of C_{total} along the formerly null directions are exactly zero or merely smaller than the physical eigenvalues, including these directions in a χ^2 test via standard matrix inversion produces inflated values with the wrong effective number of degrees of freedom. A valid goodness-of-fit test must therefore be restricted to the subspace where the measurement carries genuine sensitivity, which is the role of the projection method developed in the following section.

3 Range-projected χ^2 statistic

The analysis of Sec. 2 established that the statistical covariance matrix C_{stat} is in general rank-deficient, with N_{null} directions along which the measurement carries no statistical sensitivity. A global goodness-of-fit must therefore be formulated in the subspace where independent statistical information is available.

The rank structure discussed previously was established at the level of reconstructed event counts. The blockwise unfolding procedure of Ref. [15] propagates the full covariance matrix through the block-diagonal unfolding matrix, which is invertible by construction.

the SVD decomposition of the statistical covariance matrix as:

$$C_{\text{stat}} = U \Sigma U^T, \quad (3.1)$$

and that $r = \text{rank}(C_{\text{stat}})$. partitionings of $U = [U_r | \bar{U}_r]$

$$\begin{aligned} U_r U_r^T &= \text{projection on to } \text{ran}(C_{\text{stat}}) \\ \bar{U}_r \bar{U}_r^T &= \text{projection on to } \text{null}(C_{\text{stat}}) \end{aligned} \quad (3.2)$$

3.1 Projected chi-squared

3.2 Proof sketch / justification

3.3 Relation to alternative approaches

Acknowledgments

A Algorithm

The naive counting of the common refinement $\mathcal{R}$ has cost $\prod_v M_v$, which grows exponentially with the number of kinematic variables. A typical multi-distribution measurement involves $\mathcal{O}(5)$ variables each with $\mathcal{O}(10)$ refined bins, giving $\mathcal{O}(10^5)$ cells. The key insight is that only shared variables (i.e. the ones appearing in two or more blocks) generate the cross-block constraints. Dividing the variable set into shared (V_s) and private (V_p) variables, one has to enumerate only the reduced refinement $\prod_{v \in V_s} M_v$. For each of the shared-variable cells, the bins from blocks that depend on the shared variable are determined by lookup, while blocks whose binning variable is private contribute their bins independently via a local Cartesian product. If the number of shared variables is small the computational cost is dramatically reduced.

References

- [1] MICROBooNE collaboration, *First Simultaneous Measurement of Differential Muon-Neutrino Charged-Current Cross Sections on Argon for Final States with and Without Protons Using MicroBooNE Data*, *Phys. Rev. Lett.* **133** (2024) 041801 [[2402.19281](#)].
- [2] MICROBooNE collaboration, *Inclusive cross section measurements in final states with and without protons for charged-current ν_μ -Ar scattering in MicroBooNE*, *Phys. Rev. D* **110** (2024) 013006 [[2402.19216](#)].
- [3] MICROBooNE collaboration, *First Double-Differential Measurement of Kinematic Imbalance in Neutrino Interactions with the MicroBooNE Detector*, *Phys. Rev. Lett.* **131** (2023) 101802 [[2301.03706](#)].
- [4] MICROBooNE collaboration, *Multidifferential cross section measurements of $\nu\mu$ -argon quasielasticlike reactions with the MicroBooNE detector*, *Phys. Rev. D* **108** (2023) 053002 [[2301.03700](#)].
- [5] MICROBooNE collaboration, *Measurement of three-dimensional inclusive muon-neutrino charged-current cross sections on argon with the MicroBooNE detector*, *Phys. Lett. B* **870** (2025) 139939 [[2307.06413](#)].

- [6] T2K collaboration, *Characterization of nuclear effects in muon-neutrino scattering on hydrocarbon with a measurement of final-state kinematics and correlations in charged-current pionless interactions at T2K*, *Phys. Rev. D* **98** (2018) 032003 [[1802.05078](#)].
- [7] T2K collaboration, *First T2K measurement of transverse kinematic imbalance in the muon-neutrino charged-current single- π^+ production channel containing at least one proton*, *Phys. Rev. D* **103** (2021) 112009 [[2102.03346](#)].
- [8] NOvA collaboration, *Measurement of the double-differential muon-neutrino charged-current inclusive cross section in the NOvA near detector*, *Phys. Rev. D* **107** (2023) 052011 [[2109.12220](#)].
- [9] NOvA collaboration, *Measurement of $d^2\sigma/d|\vec{q}|dE_{avail}$ in charged current ν_μ -nucleus interactions at $\langle E_\nu \rangle = 1.86$ GeV using the NOvA Near Detector*, *Phys. Rev. D* **111** (2025) 052009 [[2410.05526](#)].
- [10] NOvA collaboration, “Measurement of the double-differential cross section of muon-neutrino charged-current interactions with low hadronic energy in the NOvA Near Detector.” 10, 2024.
- [11] T2K collaboration, *Simultaneous measurement of the muon neutrino charged-current cross section on oxygen and carbon without pions in the final state at T2K*, *Phys. Rev. D* **101** (2020) 112004 [[2004.05434](#)].
- [12] MINERvA collaboration, *Simultaneous Measurement of ν_μ Quasielasticlike Cross Sections on CH, C, H₂O, Fe, and Pb as a Function of Muon Kinematics at MINERvA*, *Phys. Rev. Lett.* **130** (2023) 161801 [[2301.02272](#)].
- [13] MINERvA collaboration, *Simultaneous Measurement of Muon Neutrino ν_μ Charged-Current Single π^+ Production in CH, C, H₂O, Fe, and Pb Targets in MINERvA*, *Phys. Rev. Lett.* **131** (2023) 011801 [[2209.07852](#)].
- [14] T2K collaboration, *First combined measurement of the muon neutrino and antineutrino charged-current cross section without pions in the final state at T2K*, *Phys. Rev. D* **101** (2020) 112001 [[2002.09323](#)].
- [15] S. Gardiner, “Mathematical methods for neutrino cross-section extraction.” 2024.
- [16] G. D’Agostini, *A Multidimensional unfolding method based on Bayes’ theorem*, *Nucl. Instrum. Meth. A* **362** (1995) 487.
- [17] W. Tang, X. Li, X. Qian, H. Wei and C. Zhang, *Data Unfolding with Wiener-SVD Method*, *JINST* **12** (2017) P10002 [[1705.03568](#)].
- [18] C. Wilkinson et al., *Testing charged current quasi-elastic and multinucleon interaction models in the NEUT neutrino interaction generator with published datasets from the MiniBooNE and MINERvA experiments*, *Phys. Rev. D* **93** (2016) 072010 [[1601.05592](#)].
- [19] MINERvA collaboration, *Tuning the GENIE Pion Production Model with MINERvA Data*, *Phys. Rev. D* **100** (2019) 072005 [[1903.01558](#)].
- [20] GENIE collaboration, *Neutrino-nucleus $CC0\pi$ cross-section tuning in GENIE v3*, *Phys. Rev. D* **106** (2022) 112001 [[2206.11050](#)].
- [21] MICROBooNE collaboration, *Measurement of single- and double-differential cross sections for mesonless charged-current muon neutrino interactions on argon with final-state protons using the MicroBooNE detector*, *Phys. Rev. D* **112** (2025) 112004 [[2403.19574](#)].