


¹ ADVANCING NEUTRINO
² DETECTION AND TRIGGERING IN
³ DUNE



⁵ Francisco Martínez López

⁶ Submitted in partial fulfillment of the requirements
⁷ of the Degree of Doctor of Philosophy

⁸ School of Physical and Chemical Sciences

⁹ Queen Mary University of London

¹⁰ December 2023

¹¹ Statement of originality

- ¹² I, [insert name as recorded in QM records], confirm that the research included within
¹³ this thesis is my own work or that where it has been carried out in collaboration with, or
¹⁴ supported by others, that this is duly acknowledged below and my contribution indicated.
¹⁵ Previously published material is also acknowledged below.
- ¹⁶ I attest that I have exercised reasonable care to ensure that the work is original, and
¹⁷ does not to the best of my knowledge break any UK law, infringe any third party's
¹⁸ copyright or other Intellectual Property Right, or contain any confidential material.
- ¹⁹ I accept that the College has the right to use plagiarism detection software to check the
²⁰ electronic version of the thesis.
- ²¹ I confirm that this thesis has not been previously submitted for the award of a degree
²² by this or any other university.
- ²³ The copyright of this thesis rests with the author and no quotation from it or information
²⁴ derived from it may be published without the prior written consent of the author.
- ²⁵ Signature: [can be digital signature]
- ²⁶ Date:
- ²⁷ Details of collaboration and publications:
²⁸ [insert details here if applicable]

²⁹ Abstract

³⁰ Hello, here is some text without a meaning. This text should show what a printed text
³¹ will look like at this place. If you read this text, you will get no information. Really? Is
³² there no information? Is there a difference between this text and some nonsense like
³³ “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about
³⁴ the selected font, how the letters are written and an impression of the look. This text
³⁵ should contain all letters of the alphabet and it should be written in of the original
³⁶ language. There is no need for special content, but the length of words should match
³⁷ the language.

O time, thou must untangle this, not I.
It is too hard a knot for me to untie!

Twelfth Night

SHAKESPEARE

³⁸ Acknowledgements

³⁹ In principio fu la Maestra Manuela.

⁴⁰ A heartfelt thank you to . . . for accepting to be the referees of this thesis, and for
⁴¹ the thought-provoking discussions we shared. Their comments brought this manuscript
⁴² to a significant improvement.

⁴³ I feel privileged to have this many eminent researchers in my committee. A very big
⁴⁴ thank you for accepting to be part of it.

⁴⁵ Thank you to . . .

⁴⁶ Un ringraziamento va anche agli amici triestini di nascita, di adozione, o di passaggio,
⁴⁷ matematici, sballerine, o altro, oggi sparpagliati per il mondo a formare una famiglia
⁴⁸ grazie alla quale uno non si sente mai troppo solo e lontano da tutti. Tanto siamo
⁴⁹ dappertutto.

⁵⁰ In particolare grazie a . . .

⁵¹ Prima di giungere ai ringraziamenti più personali, . . .

⁵² Contents

⁵³	Statement of originality	3
⁵⁴	Abstract	5
⁵⁵	Acknowledgements	9
⁵⁶	List of Figures	15
⁵⁷	List of Tables	29
⁵⁸	List of Abbreviations	31
⁵⁹	1 Introduction	33
⁶⁰	2 Neutrino physics	35
⁶¹	2.1 Neutrinos in the SM	35
⁶²	2.2 Neutrino oscillations	36
⁶³	2.2.1 Oscillations in vacuum	37
⁶⁴	2.2.2 Oscillations in matter	38
⁶⁵	2.2.3 Current status of neutrino oscillations	39
⁶⁶	2.3 Open questions in the neutrino sector	40
⁶⁷	3 The Deep Underground Neutrino Experiment	43
⁶⁸	3.1 Overview	43
⁶⁹	3.2 Physics goals of DUNE	45
⁷⁰	3.3 Far Detector	47

CONTENTS

71	3.3.1	Horizontal Drift	48
72	3.3.2	Vertical Drift	51
73	3.3.3	FD Data Acquisition System	53
74	3.4	Near Detector	54
75	3.4.1	ND-LAr	56
76	3.4.2	TMS/ND-GAr	57
77	3.4.3	PRISM	58
78	3.4.4	SAND	60
79	3.5	LBNF beamline	60
80	4	ND-GAr	63
81	4.1	Requirements	63
82	4.2	Reference design	64
83	4.2.1	HPgTPC	64
84	4.2.2	ECal	65
85	4.2.3	Magnet	66
86	4.2.4	Muon system	67
87	4.3	GArSoft	67
88	4.3.1	Event generation	67
89	4.3.2	Detector simulation	68
90	4.3.3	Reconstruction	69
91	5	FWTPG offline software	73
92	6	Matched Filter approach to induction wire Trigger Primitives	75
93	6.1	Motivation	75
94	6.2	Signal-to-noise ratio definition	77
95	6.3	Low-pass FIR filter design	79
96	6.4	Matched filters	82
97	6.5	Using simulated samples	88

CONTENTS

98	6.5.1 Angular dependence	94
99	6.5.2 Distortion and peak asymmetry	96
100	6.5.3 Hit sensitivity	99
101	7 DM searches with neutrinos from the Sun	109
102	7.1 Motivation	109
103	7.2 Gravitational capture of DM by the Sun	109
104	7.3 Neutrino flux from DM annihilations	116
105	7.4 Computing limits from solar neutrino fluxes	117
106	7.5 Example: Kaluza-Klein Dark Matter	121
107	7.6 High energy DM neutrino signals	125
108	7.6.1 DIS events	127
109	7.6.2 Single proton QEL events	132
110	7.6.3 Results	135
111	7.7 Example: Leptophilic Dark Matter	137
112	8 Particle ID in GArSoft	143
113	8.1 dE/dx measurement in the TPC	144
114	8.1.1 Energy calibration	146
115	8.1.2 Truncated dE/dx mean	156
116	8.1.3 Mean dE/dx parametrisation	159
117	8.1.4 Particle identification	163
118	8.2 Muon and pion separation in the ECal and MuID	163
119	8.2.1 Track-ECal matching	166
120	8.2.2 Classification strategy	170
121	8.2.3 Feature selection and importance	173
122	8.2.4 Hyperparameter optimisation	176
123	8.2.5 Probability calibration	176
124	8.2.6 Performance	176
125	8.3 ECal time-of-flight	176

CONTENTS

126	8.3.1 Arrival time estimations	176
127	8.3.2 Proton and pion separation	176
128	8.4 Charged pion decay in flight	176
129	8.4.1 Track breakpoints	183
130	8.5 Neutral particle identification	183
131	8.5.1 ECal clustering	183
132	8.5.2 π^0 reconstruction	187
133	9 Event selection in ND-GAr	191
134	9.1 CAFs and CAFAna	191
135	9.2 Event selection	191
136	9.2.1 ν_μ CC selection	191
137	9.2.2 Charged pion multiplicity	191
138	10 Conclusions	193
139	A An appendix	195
140	Bibliography	197

List of Figures

142	3.1 Schematic diagram of the DUNE experiment and the LBNF beamline [1].	44
143	3.2 Schematic diagram showing the operating principle of a LArTPC with	
144	wire readout	48
145	3.3 Proposed design for the FD-1 and FD-2 modules following the HD principle.	49
146	3.4 Schematic representation of an APA frames showing the U, V, X and G	
147	wires.	50
148	3.5 A PDS module containing 24 X-ARAPUCAs and the location of the	
149	modules on the APAs.	50
150	3.6 Proposed design for the FD-3 module following the VD principle.	51
151	3.7 Schematic representation of the electrode strip configuration for a top	
152	and bottom CRU.	52
153	3.8 Detailed diagram of the DUNE FD DAQ system. Figure taken from Ref.	
154	[2].	53
155	3.9 Representation of the ND hall in Phase II, showing the different subcomponents.	55
156	3.10 Schematic representation of the external components of ND-LAr, including	
157	the cryostat and the PRISM movable system and detailed drawing of one	
158	ArgonCube module.	56
159	3.11 Schematic view of the TMS detector, highlighting its main parts.	57
160	3.12 Cross section of the ND-GAr geometry, showing the HPgTPC, ECal and	
161	magnet.	58

LIST OF FIGURES

162	3.13	Predicted beam muon neutrino flux at the ND location for different off-axis positions.	59
163			
164	3.14	Schematic longitudinal section of the LBNF beamline at Fermilab.	60
165	3.15	Predicted neutrino fluxes at the FD in FHC mode and RHC mode.	61
166	4.1	Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers.	65
167			
168	4.2	Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers.	66
169			
170	6.1	<i>Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G.</i>	76
171			
172			
173			
174	6.2	<i>Left panel: Zoomed unfiltered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code> (blue line). The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{noise}}$.</i>	77
175			
176			
177			
178			
179			
180			
181	6.3	<i>Left panel: Zoomed filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code> (blue line). The filter used was the current implementation of the low-pass FIR filter in <code>dtp-firmware</code>. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$</i>	79
182			
183			
184			
185			
186			
187			
188			

LIST OF FIGURES

189	6.4 Power spectrum in decibels for the current implementation of the low-pass	
190	FIR filter in <code>dtp-firmware</code> (blue line), compared to the response of an	
191	optimal filter obtained using the Parks-McClellan algorithm for the same	
192	pass-band (red line). Also for comparison I include the spectrum of the	
193	optimal filter when taking only the integer part of the coefficients (red	
194	dashed line)	80
195	6.5 Relative change in the S/N for the ProtoDUNE-SP raw data capture	
196	<code>felix-2020-07-17-21:31:44</code> , using different values of the cutoff frequency	
197	f_c and the transition width δf . The optimal Chebyshev filters were applied	
198	using just the integer part of the coefficients given by the Parks-McClellan	
199	algorithm.	81
200	6.6 Distribution of the relative change of the S/N on the different wire planes	
201	from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code>	
202	after the optimal Chebyshev filter was applied. The filter was computed	
203	with the Parks-McClellan algorithm using a cutoff of $f_c = 0.068 \text{ ticks}^{-1}$	
204	and a transition width $\delta f = 0.010 \text{ ticks}^{-1}$	82
205	6.7 Left panel: Zoomed match filtered waveform corresponding to channel 7840	
206	from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code>	
207	(blue line). The filter used was directly extracted from the data, being	
208	the 32 values around the first peak in the original waveform. The green	
209	dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is	
210	also shown (red line). Top right panel: ADC distribution for channel 7840	
211	after match filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$.	
212	Bottom right panel: noise ADC distribution for channel 7840 after match	
213	filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$	83
214	6.8 Relative improvement in the S/N for the raw data capture <code>felix-2020-07-17-21:31:44</code> ,	
215	using the matched filter following the parametrisation in Eq. (6.17). The	
216	black crosses in both panels denote the location of the maximum ratio value.	86

LIST OF FIGURES

217	6.9 Left panel: Optimal matched filter coefficients for the U (blue line) and V (red line) planes. The filters were computed with our parametrisation in Eq. (6.17) for the parameter values $\delta = 0.035$, $\sigma = 0.191$ and $\delta = 0.018$, $\sigma = 0.191$ respectively. Right panel: Distribution of the relative change of the S/N on the two induction wire planes from the ProtoDUNE-SP raw data capture <i>felix-2020-07-17-21:31:44</i> after their respective optimal matched filters were applied.	89
224	6.10 Left panel: distributions of the particles track length in the liquid argon for the generated $E_k = 100$ MeV monoenergetic samples, electrons (blue), muons (red), protons (green) and neutral pions (purple). Right panel: distribution of the length of the longest photon in the neutral pion sample after the decay process $\pi^0 \rightarrow \gamma\gamma$	90
229	6.11 Left panel: schematic representation of the two new rotated reference frames used in this analysis (denoted as prime and double prime), viewed from the yz plane. The magenta stack of lines represent the wires in the U plane, whereas the green lines correspond to the wires in the V plane. Right panel: 3D representation of the momentum of one of the generated monoenergetic muons (red arrow) in the original reference frame (black lines), along with the new reference frame used for the U plane waveforms (blue lines). In the yz plane I added the projection of these three.	92
237	6.12 Distributions of the mean S/N improvement per event for the corresponding sample after applying the matched filters. Here I separated the change in the U plane (blue) and the V plane (red) channels. From top left to the right: muon, electron, proton and neutral pion. All the events have a fixed kinetic energy of $E_k = 100$ MeV.	93

LIST OF FIGURES

LIST OF FIGURES

262	6.16 Left panel: peak asymmetry distribution for the case of the monoenergetic	
263	$E_k = 100$ MeV muon sample. Each value corresponds to a single bipolar	
264	signal peak from a channel in any event. The blue distribution represents	
265	the peaks on U plane channels, whereas the red corresponds to signal peaks	
266	in V wires. Right panel: relation between the mean peak asymmetry per	
267	event with the S/N for U channel waveforms from the $E_k = 100$ MeV	
268	muon sample. The top subplot shows the decimal logarithm of the mean	
269	S/N for the raw (red) and the matched filtered (blue) waveforms. The	
270	bottom subplot contains the mean S/N improvement ratio after the matched	
271	filter was applied.	98
272	6.17 Raw data display in the plane time (in firmware ticks) vs. offline channel	
273	number for an $E_k = 100$ MeV electron event. The produced true hits are	
274	superimposed (black boxes) as well as the hits comming from the standard	
275	hit finder chain (blue circles) and the hit finder using the matched filter	
276	(green triangles).	100
277	6.18 Dependence of the precision (blue), sensitivity (red) and F_1 (green) scores	
278	on the threshold values used in the hit finder, for the FIR (left panel)	
279	and matched filter (right panel) cases. The results were obtained after	
280	matching the hits to the true hits in the case of the isotropic muon sample	
281	with kinetic energy in the range 5 to 100 MeV, taking only into account	
282	the induction plane channels. The points represent the mean value while	
283	the error bars indicate one standard deviation around that mean value. . .	102

LIST OF FIGURES

- 284 6.19 *Dependence of the averaged hit sensitivity on the kinetic energy of the*
 285 *events for the matched filter (blue) and standard (red) hits, for the case of*
 286 *the muon (left panel) and electron (right panel) samples, separated between*
 287 *U (top plots) and V (bottom plots) induction wire planes. The top subplots*
 288 *contain the hit sensitivities for the two hit finder alternatives, while the*
 289 *bottom subplots show the ratio between the two. The horizontal lines sit at*
 290 *the mean value and represent the size of the energy bins, while the vertical*
 291 *error bars indicate one standard deviation around that mean value.* 104
- 292 6.20 *Distributions of the hit sensitivity in the U (top panels) and V (bottom*
 293 *panels) planes versus the hit sensitivity in the X plane, both for the*
 294 *standard hits (left panels) and the matched filter hits (right panels), in the*
 295 *case of the electron sample and a threshold of 30 ADC.* 105
- 296 6.21 *Top panels: standard residual plots of the hit sensitivities between the X*
 297 *and U planes. Bottom panels: quantile-quantile plots of the hit sensitivity*
 298 *standard residuals between the X and U planes. In all cases, the left*
 299 *panel corresponds to the standard hits while the right panel represents the*
 300 *matched filter case, all from the electron sample with a 30 ADC threshold.* 106
- 301 7.1 *Input solar parameters used in our capture rate computation as functions*
 302 *of the Sun's radius, from left to right: temperature (with respect to the*
 303 *temperature at the core), mass (in solar masses) and electron number*
 304 *density (with respect to the electron density at the core). All quantities*
 305 *shown correspond to the standard solar model BS2005-OP [3].* 113
- 306 7.2 *Capture rates as a function of the DM mass for the DM-electron interactions*
 307 *(red lines), SD DM-nucleons interactions (green lines) and SI DM-nucleons*
 308 *interactions (blue lines). Solid lines represent the values computed in this*
 309 *work while the dashed lines are the one given in Ref. [4]. All the rates*
 310 *are shown for a choice of scattering cross section of $\sigma_i = 10^{-40} \text{ cm}^2$.* . . 114

LIST OF FIGURES

311	7.3 <i>NuWro computed $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section as a function of the neutrino energy E_μ. The black line shows to the total cross section, whereas the others correspond to the different contributions (in red quasi-elastic scattering, in green resonant pion exchange, in blue deep inelastic scattering and in purple meson exchange current).</i>	118
316	7.4 <i>Expected atmospheric neutrino flux as a function of the neutrino energy E_ν at Homestake at solar minimum, taken from Ref. [5]. The blue solid (dashed) line correspond to muon neutrinos (antineutrinos) and the red solid (dashed) line correspond to electron neutrinos (antineutrinos)</i>	120
320	7.5 Feynman diagrams for B^1B^1 annihilation into SM fermions.	122
321	7.6 Feynman diagrams for B^1B^1 annihilation into a Higgs boson pair.	122
322	7.7 <i>Computed spectra of muon neutrinos at the DUNE FD site from B^1 annihilations in the Sun for three different values of M_{LKP}, plotted in relative energy units for legibility.</i>	123
325	7.8 <i>Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin- dependent B^1-proton scattering cross section as a function of M_{LKP} (green dots). I also show the previous limits from IceCube [6] (blue line) and Antares [7] (red line) on the LKP cross section. The shaded area represents the disfavoured region (at 95% confidence level) on the mass of the LKP from LHC data [8]</i>	124
331	7.9 <i>Computed spectra of muon neutrinos at the DUNE FD site from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilations in the Sun for the DM masses $m_{\text{DM}} = 10$ GeV (red line), 50 GeV (green line) and 100 GeV (blue line), plotted in relative energy units.</i>	125
335	7.10 <i>Distribution of the muon neutrino energies from the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilation channels, for $m_{\text{DM}} = 10$ GeV, separated by CC interaction type: QEL (blue), MEC (orange), RES (green) and DIS (red)</i>	127

LIST OF FIGURES

<p>339 7.11 <i>Distributions of θ_μ (left panel), θ_j (central panel) and θ_{plane} (right panel)</i> 340 <i>for the $b\bar{b}$ sample with $m_{DM} = 10$ GeV (blue) and the atmospheric</i> 341 <i>background (red).</i></p> <p>342 7.12 <i>Left panel: signal efficiencies (blue lines) and background rejections (red</i> 343 <i>lines) for events passing the cuts $\theta < \theta_{cut}$ for the jet (solid lines) and</i> 344 <i>muon (dashed lines) angles. Right panel: signal efficiency (blue line) and</i> 345 <i>background rejection (red line) for events passing the cut $\theta_{plane} < \theta_{cut}$ for</i> 346 <i>the momentum conservation plane deviation.</i></p> <p>347 7.13 <i>Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) DIS samples</i> 348 <i>as functions of the DM mass, m_{DM}, obtained by applying the optimal</i> 349 <i>angular cuts $\theta_\mu < 27^\circ$, $4^\circ < \theta_j < 26^\circ$ and $\theta_{plane} < 3.5^\circ$.</i></p> <p>350 7.14 <i>Distributions of $\cos \theta_\mu$ (left panel), $\cos \theta_p$ (central panel) and $\cos \theta_N$</i> 351 <i>(right panel) for the $\tau^+\tau^-$ QEL sample with $m_{DM} = 5$ GeV (blue) and</i> 352 <i>the atmospheric background (red).</i></p> <p>353 7.15 <i>Left panel: value of the loss function for the training sample (blue line)</i> 354 <i>and accuracy for the validation sample (red line) versus the number of</i> 355 <i>iterations for the MLP classifier training. Right panel: distributions of the</i> 356 <i>predicted probabilities assigned by the MLP classifier to the test sample</i> 357 <i>for the $\tau^+\tau^-$ QEL signal with $m_{DM} = 5$ GeV (blue) and the atmospheric</i> 358 <i>background (red).</i></p> <p>359 7.16 <i>Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) single proton</i> 360 <i>QEL samples as functions of the DM mass, m_{DM}, obtained by requiring a</i> 361 <i>minimum predicted probability from the MLP classifier of 0.97 in order to</i> 362 <i>achieve a background rejection greater than 99.8%.</i></p>	<p>128</p> <p>129</p> <p>131</p> <p>132</p> <p>133</p> <p>134</p>
---	---

LIST OF FIGURES

363	7.17 Projected 90% confidence level upper limit for DUNE (400 kT yr) on 364 the spin-dependent DM-nucleon scattering cross section as a function of 365 m_{DM} , for the annihilation channels $\tau^+\tau^-$ (blue) and $b\bar{b}$ (red) separated 366 by interaction type (up triangles denote DIS interactions whereas down 367 triangles represent QEL interactions). I also show the previous limits 368 from IceCube [9] (solid lines) and the projected sensitivities for Pingu [10] 369 (dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines), as well 370 as the direct detection limits from PICASSO [12] (solid green line) and 371 PICO-60 C ₃ F ₈ [13] (dashed green line).	136
372	7.18 Left panel: Projected 90% confidence level sensitivity of DUNE (400 kT 373 yr) to the scale Λ of an EFT containing only leptophilic DM axial-axial 374 interactions (blue line). Right panel: . In both cases the corresponding 375 limits from DarkSide-50 [14] (dotted green line) and XENON1T [15] 376 (dashed red line) are also shown, together with the configurations for which 377 the correct relic density is achieved (black line), all for the coupling values 378 $c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$	140
379	8.1 Distribution of the fraction of energy deposits with residual range less 380 than 20% of the total track length, and distribution of the ionisation per 381 unit length after removing the tracks with less than 30% of their energy 382 deposits in the last 20% of the track.	147
383	8.2 Distribution of the reconstructed ionisation charge per unit length for 384 different reclustering values, and distribution of the median change in 385 dQ/dx per track for the $N_{\text{group}} = 4$ reclustering.	148
386	8.3 Distribution of the Geant4-simulated energy losses per unit length versus 387 residual range for the stopping proton sample.	150
388	8.4 Fitted most probable dQ/dx values for each dE/dx bin, together with 389 best fit to the logarithmic calibration function.	151

LIST OF FIGURES

390	8.5 Fitted most probable dQ/dx values for each dE/dx bin for three different	
391	ADC bit limits.	153
392	8.6 Area normalised dE/dx distributions for the true and the reconstructed	
393	energy deposits in the stopping proton sample, both after applying the	
394	calibration and the calibration and the normalisation correction.	154
395	8.7 Fractional residuals between the true and the corrected dE/dx means	
396	and the 60% truncated means, and fractional residuals between the true	
397	and the uncorrected, corrected and uncalibrated dE/dx 60% truncated	
398	means.	156
399	8.8 Estimated values of the mean dE/dx bias and resolution for the stopping	
400	proton sample at different values of the truncation factor.	157
401	8.9 Examples of the truncated mean dE/dx LanGauss fits for various $\beta\gamma$	
402	bins, from a simulated FHC neutrino sample.	159
403	8.10 Resulting one and two dimensional projections of the posterior probability	
404	distributions of the ALEPH $\langle dE/dx \rangle$ parameters obtained by fitting the	
405	60% truncated mean dE/dx values from a FHC neutrino sample.	160
406	8.11 Truncated mean dE/dx obtained for the FHC neutrino sample as a	
407	function of the $\beta\gamma$ product, together with the fitted most probable values	
408	for each $\beta\gamma$ bin and the best fit obtained using the ALEPH parametrisation.	161
409	8.12 Distribution of the 60% truncated mean dE/dx versus reconstructed	
410	momentum for the FHC neutrino sample.	162
411	8.13 Estimated values of the mean dE/dx bias and resolution obtained for the	
412	true protons in a FHC neutrino sample.	163
413	8.14 True momentum distribution for the primary muon in ν_μ CC $N\pi^\pm$	
414	interactions inside the fiducial volume of ND-GAr, compared to the	
415	post FSI charged pion spectrum.	164
416	8.15 Distributions of energy deposits in the ECal for a muon and a charged	
417	pion with similar momenta.	165

LIST OF FIGURES

418	8.16 Left panel: comparison between the precision (blue), sensitivity (yellow) and F_1 score (red) obtained for the default (horizontal lines) and new algorithms, both with the χ^2 -based direction estimator (squares) and cheating the directions (circles), for different values of the χ^2 cut. Right panel: comparison of the performance of the new algorithm when applying the cluster t_0 correction (squares) and when (circles).	167
424	8.17 Schematics of a possible option to deal with track-ECal associations in non-zero t_0 neutrino interaction events, trying to correct for the drift direction uncertainty in a cluster-by-cluster basis using the cluster time, $t_{cluster}$	169
428	8.18 Momentum distribution for the reconstructed muons (top panel) and charged pions (bottom panel) in a FHC neutrino sample, together with the fraction of them reaching the ECal (red) and MuID (blue). Each entry corresponds to a reconstructed track, backtracked to a true muon or pion which has not produced any other reconstructed track.	171
433	8.19 Predicted truncated mean dE/dx versus momentum, for electrons, muons, charged pions and protons, obtained using the ALEPH parametrisation. The vertical dashed lines represent the boundaries of the six regions used for the muon and pion classification training.	172
437	8.20 Left panel: number of non-decaying, decaying and decaying in the fiducial volume pions for a MC sample of 10^5 , $p = 500$ MeV isotropic positively charged pions inside the TPC. Right panel: event display for a positive pion decaying inside the fiducial volume, with a single reconstructed track for the pion and muon system.	177
442	8.21 Values of $\chi_k^{2(FB)}$ (top left panel), F_k (top right panel), $D_k^{1/R}$ (bottom left panel) and D_k^ϕ (bottom right panel) versus position along the drift direction for a reconstructed track in a positive pion decay event. The vertical red dashed line indicates the true location of the decay point. . .	178

LIST OF FIGURES

446	8.22 Fractional residual distributions of the true and reconstructed decay position along the drift coordinate, using the position of the maximum of $\chi_k^2(FB)$ (left panel) and F_k (right panel) as estimates of the decay position. Also shown are double Gaussian fits to these points (red lines).	180
447		
448		
449		
450	8.23 Distributions of the extreme values of $\chi_k^2(FB)$ (top left panel), F_k (top right panel), $D_k^{1/R}$ (bottom left panel) and D_k^ϕ (bottom right panel) for non-decaying reconstructed pion tracks (blue) and tracks which include the decay inside the fiducial volume (red).	181
451		
452		
453		
454	8.24 Left panel: distributions of the predicted probabilities assigned by the BDT classifier to a test sample of decaying pion+muon tracks (blue) and non-decaying pion tracks (red). Left: signal efficiency versus background acceptance (ROC curve) obtained from the BDT for the test sample.	182
455		
456		
457		
458	8.25 Left panel: dependence of the decay position finding resolution on the true value of the decay angle for the $\chi_k^2(FB)$ (red) and F_k (blue) methods. Right panel: signal efficiency (blue line) and background rejection (red line) from the BDT classifier versus true decay angle.	182
459		
460		
461		
462	8.26 Mean values of the F_1 -score marginal distributions for the different free parameters of the new clustering algorithm, with the error bars representing one standard deviation around the mean. The F_1 -score values were computed for the 6561 possible parameter configurations using 1000 ν_μ CC interaction events.	185
463		
464		
465		
466		
467	8.27 Left panel: distributions of the number of ECal clusters per photon from π^0 decays for the standard (red) and new (blue) clustering algorithms. Right panel: reconstructed invariant mass distributions for photon pairs from single π^0 events using the standard (red) and new (blue) ECal clustering algorithms.	187
468		
469		
470		
471		

List of Tables

473	2.1	Summary of neutrino oscillation parameters determined in the Neutrino	
474		Global Fit of 2020 [16].	40
475	3.1	Summary of the two-phased plan for DUNE	45
476	3.2	Exposure and time required to achieve the different physics milestones of	
477		the two phases	46
478	6.1	<i>Characteristic parameters of the two monoenergetic muon events selected, relative to the U plane: projected angles in the xz' and $y'z'$ planes, S/N values for the raw and filtered waveforms, mean improvement of the S/N and peak asymmetry.</i>	97
482	8.1	Calibration parameters obtained from the fit of the ND-GAr simulated stopping proton sample to the calibration function, for different ADC limits.	153
484	8.2	Momentum ranges and description of the PID approach assumed for the muon and pion classification task.	173
486	8.3	Summary of parameters and sampled values used in the optimisation of the clustering algorithm.	186
488			

⁴⁸⁹ List of Abbreviations

ALEPH Apparatus for LEP PHysics.

ALICE A Large Ion Collider Experiment.

BDT Boosted Decision Tree.

DUNE Deep Underground Neutrino Experiment.

⁴⁹⁰ Chapter 1

⁴⁹¹ Introduction

492 Chapter 2

493 Neutrino physics

494 Ever since they were postulated in 1930 by Wolfgang Pauli to explain the continuous
495 β decay spectrum [17] and later found by Reines and Cowan at the Savannah River
496 reactor in 1953 [18], neutrinos have had a special place among all other elementary
497 particles. They provide a unique way to probe a wide range of quite different physics,
498 from nuclear physics to cosmology, from astrophysics to colliders. Moreover, there is
499 compelling evidence to believe that the study of neutrinos may be key to unveil different
500 aspects of physics beyond the SM, difficult to test elsewhere.

501 In this Chapter I will review the basics of neutrino physics, from its role within the
502 SM to the main open questions related to the neutrino sector, paying special attention
503 to the phenomenology of neutrino oscillations.

504 2.1 Neutrinos in the SM

505 By definition, in the SM there are no right-handed neutrino fields. A direct implication
506 of this fact is that neutrinos are strictly massless within the SM. This follows from the
507 experimental observation that all neutrinos produced via weak interactions are pure
508 left-handed helicity states (and similarly antineutrinos are pure right-handed states).
509 The hypothetical existence of right-handed neutrinos could be indirectly inferred from
510 the observation of non-zero neutrino masses, nevertheless the existence neutrino masses

Chapter 2. Neutrino physics

511 is not a sufficient condition for the existence of such fields.

512 In the SM neutrinos appear in three flavours, namely ν_e , ν_μ and ν_τ . These are
513 associated with the corresponding charged leptons e , μ and τ , in such a way that the
514 charged current part of the Lagrangian coupling them is diagonal. As in the electroweak
515 theory neutrinos are coupled to the Z boson in a universal way, by measuring the so-called
516 invisible decay width of the Z we have an estimate of the number of light (i.e. lighter
517 than the Z boson) neutrino flavours. This number was measured by LEP in a combined
518 analysis of $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow$ hadrons to be $N_\nu = 2.9840 \pm 0.0082$ [19].

519 2.2 Neutrino oscillations

520 The evidence for neutrino oscillation [20], and therefore the existence of non-zero neutrino
521 masses, constitutes one of the groundbreaking discoveries of modern Physics and has
522 acted as driving force for Beyond the Standard Model (BSM) Physics. The minimal
523 extension of the Standard Model (SM) we can do to address these phenomena is
524 introducing distinct masses for at least two of the neutrinos. This way, we are left with
525 three neutrino mass eigenstates ν_1 , ν_2 and ν_3 , with masses m_1 , m_2 and m_3 respectively,
526 which in general will not coincide with the flavour eigenstates ν_e , ν_μ and ν_τ .

527 The way to relate these two sets of neutrino eigenstates is via a 3×3 unitary matrix,
528 called the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [21, 22], as:

$$|\nu_\alpha\rangle = \sum_{i=1}^3 U_{\alpha i}^* |\nu_i\rangle, \quad (2.1)$$

529 where the Greek index α denotes the flavour $\{e, \mu, \tau\}$ and the Latin index i the associated
530 masses $\{1, 2, 3\}$. This leptonic mixing matrix may be parametrized in terms of 6
531 parameters, 3 of which are mixing angles θ_{12} , θ_{13} and θ_{23} , one CP-violating phase δ_{CP}

2.2. Neutrino oscillations

532 and 2 Majorana phases α and β :

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{CP}} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta_{CP}} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\beta} \end{pmatrix}, \quad (2.2)$$

533 where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$. This matrix is analogous to the Cabibbo-Kobayashi-
534 Maskawa (CKM) matrix in the quark sector. If neutrinos are Dirac fermions, we can
535 drop the Majorana phases in the PMNS matrix. But, in any case, these phases play no
536 role on the neutrino oscillations.

537 2.2.1 Oscillations in vacuum

538 Consider the case where a neutrino of flavour α is produced at $t = 0$, and then it
539 propagates through vacuum. Such a state will evolve in time according to the relation:

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^3 U_{\alpha i}^* e^{-iE_i t} |\nu_i(t=0)\rangle, \quad (2.3)$$

540 as the mass eigenstates are also eigenstates of the free Hamiltonian. Now, if we express
541 the mass eigenstates as a superposition of flavour eigenstates, the last expression can be
542 rewritten as:

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^3 U_{\beta i} e^{-iE_i t} U_{\alpha i}^* |\nu_\beta\rangle. \quad (2.4)$$

543 This way, the probability for the neutrino to transition from flavour α to flavour β
544 will be given by:

$$P(\nu_\alpha \rightarrow \nu_\beta) = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{i=1}^3 U_{\beta i} e^{-iE_i t} U_{\alpha i}^* \right|^2. \quad (2.5)$$

545 A usual approximation to take at this point is to consider ultra-relativistic neutrinos,
546 i.e. $E \approx |\vec{p}|$, so we can write the dispersion relations as:

$$E_i = \sqrt{p^2 + m_i^2} \approx E + \frac{m_i^2}{2E}, \quad (2.6)$$

Chapter 2. Neutrino physics

547 so we can write the oscillation probability as:

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\beta) &= \sum_{i,j} U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^* e^{-i \frac{\Delta m_{ij}^2}{2E} t} \\ &= \delta_{\alpha\beta} - 4 \sum_{i < j} \Re [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin^2 \left(\frac{\Delta m_{ij}^2}{4E} t \right) \\ &\quad + 2 \sum_{i < j} \Im [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin \left(\frac{\Delta m_{ij}^2}{2E} t \right), \end{aligned} \quad (2.7)$$

548 where Δm_{ij}^2 is the difference of the squared masses of the j th and i th neutrino mass
 549 eigenvalues. At this point, it is usual to write the phase responsible for the oscillations
 550 as (under the approximate assumption $t \approx L$):

$$\Delta_{ij} \equiv \frac{\Delta m_{ij}^2}{4E} L \simeq 1.27 \frac{\Delta m_{ij}^2}{(\text{eV}^2)} \frac{L}{(\text{km})} \frac{(\text{GeV})}{E}. \quad (2.8)$$

551 Notice that, in the case of antineutrinos the only difference would be the sign of the
 552 last term in the oscillation probability. This way, one can write the CP asymmetry as:

$$\begin{aligned} A_{CP}^{\alpha\beta} &= P(\nu_\alpha \rightarrow \nu_\beta) - P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) \\ &= 4 \sum_{i < j} \Im [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin 2\Delta_{ij}. \end{aligned} \quad (2.9)$$

553 2.2.2 Oscillations in matter

554 When neutrinos propagate through matter, their oscillation can be affected in mainly
 555 two ways. First, neutrinos can inelastically scatter with nuclei, thus destroying the
 556 coherent propagation of their quantum state. Nevertheless, in most cases this effect is
 557 negligible (even in very dense mediums like the core of the Sun). Second, neutrinos can
 558 also experience coherent or forward scatterings, that can affect their oscillation but not
 559 lose the coherent propagation of the state.

560 The first proposed model to account for neutrino oscillations in matter was proposed
 561 by Mikhaev, Smirnov and Wolfenstein (MSW) [23]. It relies on the fact that, as the
 562 only charged lepton present in ordinary matter is the electron, electron neutrinos can

2.2. Neutrino oscillations

563 undergo both charged and neutral-current interactions with matter whereas for muon
564 and tau neutrinos just neutral currents are possible.

565 2.2.3 Current status of neutrino oscillations

566 A wide range of neutrino experiments provide experimental input to the neutrino
567 oscillation framework, both using natural or synthetic neutrino sources. The results
568 from one of the neutrino global fit analyses, shown in Tab. 2.1¹, summarise well our
569 current understanding of the different oscillation parameters.

570 **Solar neutrino experiments** detect neutrinos produced in thermonuclear reactions
571 inside the Sun, mainly from the so-called *pp* chain and the CNO cycle. These neutrinos
572 have a typical energy in the range from 0.1 to 20 MeV. These experiments (Homestake
573 [24], GALLEX [25], SAGE [26], Borexino [27], Super-Kamiokande [28] and SNO [29])
574 provide the best sensitivities to θ_{12} and Δm_{21}^2 .

575 **Atmospheric neutrino experiments** detect the neutrino flux produced when
576 cosmic rays scatter with particles in Earth's atmosphere. These collisions generate particle
577 showers that eventually produce electron and muon neutrinos (and antineutrinos). Their
578 energies range from few MeV to about 10^9 GeV. Experiments, like Super-Kamiokande
579 [30] and IceCube [31] use atmospheric neutrinos to measure oscillations and are specially
580 sensitive to θ_{23} and Δm_{32}^2 .

581 **Reactor neutrino experiments** look for the $\bar{\nu}_e$ spectrum produced by nuclear
582 reactors, with energies in the MeV scale. Depending on the distance to the source,
583 long-baseline experiments like KamLAND [32] are sensitive to the solar mass splitting
584 Δm_{21}^2 whereas much shorter baseline experiment such as RENO [33] or DayaBay [34]
585 measure θ_{13} and Δm_{31}^2 .

586 **Accelerator experiments** measure neutrino fluxes generated in particle accelerators.
587 Usually mesons are produced in the accelerator to be focused into a beam, then some
588 decay to muon neutrinos and the rest are absorbed by a target. Depending on the

¹These are the results reported during M. Tórtola's talk at Neutrino 2024 (see this link). I need to keep an eye and see if they publish these or other updated results in the near future.

Chapter 2. Neutrino physics

Table 2.1: Summary of neutrino oscillation parameters determined in the Neutrino Global Fit of 2020 [16].

Parameter	Best fit $\pm 1\sigma$	3σ range
Δm_{21}^2 [eV $^2 \times 10^{-5}$]	$7.55^{+0.22}_{-0.20}$	6.98 – 8.19
$ \Delta m_{31}^2 $ [eV $^2 \times 10^{-3}$] (NO)	$2.51^{+0.02}_{-0.03}$	2.43 – 2.58
$ \Delta m_{31}^2 $ [eV $^2 \times 10^{-3}$] (IO)	$2.41^{+0.03}_{-0.02}$	2.34 – 2.49
$\sin^2 \theta_{12}/10^{-1}$	3.04 ± 0.16	2.57 – 3.55
$\sin^2 \theta_{23}/10^{-1}$ (NO)	$5.64^{+0.15}_{-0.21}$	4.23 – 6.04
$\sin^2 \theta_{23}/10^{-1}$ (IO)	$5.64^{+0.15}_{-0.18}$	4.27 – 6.03
$\sin^2 \theta_{13}/10^{-2}$ (NO)	$2.20^{+0.05}_{-0.06}$	2.03 – 2.38
$\sin^2 \theta_{13}/10^{-2}$ (IO)	$2.20^{+0.07}_{-0.04}$	2.04 – 2.38
δ_{CP}/π (NO)	$1.12^{+0.16}_{-0.12}$	0.76 – 2.00
δ_{CP}/π (IO)	$1.50^{+0.13}_{-0.14}$	1.11 – 1.87

589 configuration one can obtain a beam made of mostly neutrinos or antineutrinos. The
 590 typical energies of these neutrinos are in the GeV range. Experiments such as NOvA
 591 [35], T2K [36], MINOS [37], OPERA [?] and K2K [38] (and in the future DUNE [39])
 592 are primarily sensitive to θ_{13} , θ_{23} and Δm_{32}^2 . Also, in the coming years DUNE [39] and
 593 Hyper-Kamiokande [40] will be sensitive to δ_{CP} .

594 2.3 Open questions in the neutrino sector

595 A crucial question that remains open these days, and is of vital importance for oscillation
 596 phenomena, is whether the mass eigenvalue ν_3 is the heaviest (what we call normal
 597 ordering) or the lightest (referred to as inverted ordering) of the mass eigenstates. In
 598 other words, this means that we do not know the sign of Δm_{32}^2 , so we can either have
 599 $m_1 < m_2 < m_3$ (NO) or $m_3 < m_1 < m_2$ (IO).

600 Another big puzzle is related to the value of δ_{CP} . Nowadays it is poorly constrained,
 601 with all values between π and 2π being consistent with data. A prospective measurement
 602 different from $\delta_{CP} = 0, \pi$ will predict CP-violation in the leptonic sector, and thus

2.3. Open questions in the neutrino sector

603 contribute along with the one measured in the quark sector to the total amount of
604 CP-violation. Although it is true that these two contributions by themselves are not
605 enough to explain the matter anti-matter asymmetry in our universe, the amount of
606 CP-violation in the leptonic sector can be key to explain such imbalance.

607 Both of these questions, because of their nature, could be understood thanks to
608 future oscillation experiments.

609 Notwithstanding, there are other mysteries that can not be unveiled just by conducting
610 oscillation experiments, as certain quantities do not influence these phenomena. Among
611 these there is the question of the absolute values of the neutrino masses. Depending
612 on the value of the lightest of the neutrino masses we can have different mass spectra,
613 from hierarchical $m_1 \ll m_2 < m_3$ (NO) or $m_3 \ll m_1 < m_2$ (IO) to quasi-degenerate
614 $m_1 \simeq m_2 \simeq m_3$.

615 Other open question concerns the nature itself of the neutrinos. If neutrinos are Dirac
616 particles then their mass term can be generated through the usual Higgs mechanism
617 by adding right-handed neutrino fields. However, if they are Majorana particles and
618 therefore their own antiparticles, there is no need to add extra fields to have the mass
619 term in the Lagrangian. Experiments like SuperNEMO [?], SNO+ [?] and NEXT [?],
620 which search for neutrino-less double beta decay, will be able to determine whether
621 neutrinos are Dirac or Majorana.

622 Chapter 3

623 The Deep Underground Neutrino
624 Experiment

625 The Deep Underground Neutrino Experiment (DUNE) is a next generation long-baseline
626 neutrino experiment [1]. It will aim to address several questions in neutrino physics,
627 study neutrinos from astrophysical sources and search for beyond the standard model
628 physics.

629 This chapter reviews the main goals of the DUNE experiment, the design of the far
630 detector modules and their data acquisition (DAQ) system, and the role that the near
631 detector plays in the physics program of DUNE.

632 3.1 Overview

633 The main physics goals of DUNE are:

- 634 • measure the neutrino mass hierarchy, the amount of CP violation in the leptonic
635 sector and the θ_{23} octant,
- 636 • detect rare low energy neutrino events, like neutrinos from supernova bursts, and
- 637 • search for proton decay and other beyond the standard model phenomena.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.1: Schematic diagram of the DUNE experiment and the LBNF beamline [1].

The design of DUNE has been tailored with these goals in mind. It will consist of two neutrino detectors. A near detector (ND) complex will be placed in Fermilab, 574 m downstream of the neutrino production point, whereas a larger far detector (FD) will be built in the Sandford Underground Research Facility (SURF), South Dakota, approximately 1300 km away. Fig. 3.1 shows a simplified view of the various components of DUNE (not to scale).

The beam neutrinos to be used in DUNE will be provided by the LBNF beamline, the multi-megawatt wide-band neutrino beam planned for Fermilab. First, an intense proton beam is extracted from the Fermilab Main Injector. Then, these protons with energies between 60 GeV and 120 GeV collide with a high-power production target and produce charged mesons. Two magnetic horns allow to focus the mesons and perform a sign selection (thus having the capability to switch between neutrino and antineutrino mode). Soon after that, the mesons decay and produce neutrinos (or antineutrinos) which are then aimed to SURF.

Before arriving to the FD, the neutrino beam meets the ND complex, which serves as the experiment's control. Its role is to measure the unoscillated neutrino energy spectra. From these we can predict the unoscillated spectra at the FD, which can be compared to the spectra measured at the FD in order to extract the oscillation parameters. Therefore, the design of the DUNE ND is mainly driven by the needs of the oscillation physics program.

The liquid Argon time projection chamber (LArTPC) technology has been chosen for

3.2. Physics goals of DUNE

Table 3.1: Summary of the two-phased plan for DUNE. Adapted from Ref. [41].

Parameter	Phase I	Phase II	Benefit
FD mass	20 kt fiducial	40 kt fiducial	FD statistics
Beam power	up to 1.2 MW	2.4 MW	FD statistics
ND config.	ND-LAr, TMS, SAND	ND-LAr, ND-GAr, SAND	Systematic constraints

the FD modules of DUNE. Its four modules will record neutrino interactions from the accelerator-produced beam arriving at predictable times. As it also aims at recording rare events, the FD requires trigger schemes which can deal with both kinds of physics, and also maximum uptime.

DUNE is planned to be built using a staged approach consisting on two phases, which are summarised in Tab. 3.1. Phase I consists of a FD with 50% of the total fiducial mass, a reduced version of the ND complex and a 1.2 MW proton beam. It will be sufficient to achieve some early physics goals, like the determination of the neutrino mass ordering. For its Phase II, DUNE will feature the full four FD modules, a more capable ND and a 2.4 MW proton beam. The physics milestones for the two phases are given in Tab. 3.2, in a staging scenario which assumes that Phase II is completed after 6 years of operation.

A summary of the DUNE science program can be found in the DUNE FD Technical Design Report (TDR) Volume I [1]. For a detailed discussion on the two-phased approach the reader is referred to the DUNE Snowmass 2021 report [41].

3.2 Physics goals of DUNE

As noted in the literature (see for instance Ref. [16] for a review), the parameter space of the neutrino oscillation phenomena within the three-flavour picture is quite constrained by current experimental data. However, there are still crucial open questions, like the mass ordering, the value of δ_{CP} or the θ_{13} octant. One of the main goals of DUNE is to shed some light on the values of these parameters [42].

To address these questions DUNE can look to the subdominant oscillation channel

Chapter 3. The Deep Underground Neutrino Experiment

Table 3.2: Exposure and time required to achieve the different physics milestones of the two phases. The predictions assume a Phase II staging scenario where FD modules 3 and 4 are deployed in years 4 and 6 and both the beam and ND are upgraded after 6 years. Adapted from Ref. [41].

Stage	Physics milestone	Exposure (kt-MW-years)	Years (staged)
Phase I	5σ MO ($\delta_{CP} = -\pi/2$)	16	1-2
	5σ MO (100% of the δ_{CP} values)	66	3-5
	3σ CPV ($\delta_{CP} = -\pi/2$)	100	4-6
Phase II	5σ CPV ($\delta_{CP} = -\pi/2$)	334	7-8
	δ_{CP} resolution of 10 degrees ($\delta_{CP} = 0$)	400	8-9
	5σ CPV (50% of the δ_{CP} values)	646	11
	3σ CPV (75% of the δ_{CP} values)	936	14
	$\sin^2(2\theta_{13})$ resolution of 0.004	1079	16

681 $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) and study the energy dependence of the ν_e ($\bar{\nu}_e$) appearance probability.

682 When we focus on the antineutrino channel $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ there is a change in the sign of δ_{CP} ,
 683 thus introducing CP-violation. Moreover, due to the fact that there are no positrons in
 684 the composition of Earth, there is a sign difference for the matter effect contribution
 685 when looking to the antineutrino channel. This asymmetry is proportional to the baseline
 686 length L and is sensitive to the sign of Δ_{31} , and thus to the neutrino mass ordering.

687 Another of the main physics goals of DUNE is the search for baryon-number violating
 688 processes. Specifically, it will try to answer the question of whether protons are stable
 689 or not. There is no symmetry argument that forbids protons from decaying, but its
 690 apparent stability seems to suggest that baryon number is conserved [43]. However,
 691 proton decay is a usual feature of grand-unified theories, where electromagnetic, weak
 692 and strong interactions are unified above a certain energy scale [44].

693 As the energy deposition scale for this kind of searches is nearly the same as the one
 694 for long-baseline neutrino oscillations, DUNE will be able to look for them. It has several
 695 advantages over other experiments, such as excellent imaging and particle identification,
 696 which can be translated to lower backgrounds.

697 The last of the main objectives of DUNE is the detection of neutrinos originated in
 698 supernovae explosions, what is called a supernova neutrino burst (SNB). These neutrinos
 699 carry with them information about the core-collapse process, from the progenitor to the

3.3. Far Detector

explosion and the remnant; but also may have information about new exotic physics. So far, the only neutrino events ever recorded from such a process were a few dozens of $\bar{\nu}_e$ events from the 1987A supernova located in the Magellanic Cloud, 50 kpc away from Earth [45, 46].

DUNE aims to collect also some SNB events. Although these are quite rare, as the expected supernovae explosion events are about one every few decades for our galaxy and Andromeda, the long lifetime of the experiment (around a few decades as well) makes it reasonable to expect some. Nowadays the main sensitivity to SNB of most experiments is to the $\bar{\nu}_e$ through inverse beta decay. One of the advantages of DUNE is its expected sensitivity to ν_e , since the dominant channel will be ν_e CC scattering.

Moreover, due to the stringent requirements that the main physics goals set for DUNE, it will allow also to perform searches for all kind of BSM physics. Among others, DUNE will be able to look for: active-sterile neutrino mixing, non-unitarity of the PMNS matrix, non-standard interactions, Lorentz and CPT violations, neutrino trident production, light-mass DM, boosted DM and heavy neutral leptons. The reader is referred to the DUNE FD TDR Volume II [42] for a full discussion of the physics scope of DUNE.

3.3 Far Detector

The so-called DUNE FD complex will sit 1.5 km underground at SURF, South Dakota. Two caverns will host the four FD modules, two of them per cavern, each embedded in cryostats of dimensions 18.9 m (w) \times 17.8 m (h) \times 65.8 m (l). A central, smaller cavern will host the cryogenic system.

Three out of the four modules will be liquid argon (LAr) time projection chamber detectors, often refer to as LArTPCs, with a LAr fiducial mass of at least 10 kt each. The first and second FD modules, FD-1 and FD-2, will use a Horizontal Drift (HD) technology, whereas the third module, FD-3, will have a Vertical Drift (VD) direction. The technology for the fourth module is still to be decided,

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.2: Schematic diagram showing the operating principle of a LArTPC with wire readout. Figure taken from Ref. [1].

For each event, with energies ranging from a few MeV to several GeV, these detectors collect both the scintillation light and the ionisation electrons created when the charged particles produced in neutrino-nucleus interactions ionise the argon nuclei. In both HD and VD designs the characteristic 128 nm scintillation light of argon is collected by a photon detection system (PDS). This light will indicate the time at which electrons start to drift, thus enabling reconstruction over the drift coordinate when compared to the time when the first ionisation electron arrives to the anode. Reconstruction of the topology in the transverse direction is achieved using the charge readout. Fig. 3.2 illustrates the detection principle described, for the case of a HD detector with a wire readout.

3.3.1 Horizontal Drift

Within the HD design the ionisation electrons produced as charged particles traverse the LAr drift horizontally towards the anode planes, made out of three layers of wire readout, due to the effect of an electric field. This design, previously known as single-phase (SP), was tested by the ProtoDUNE-SP detector at CERN. The prototype collected data from

3.3. Far Detector




Figure 3.3: Proposed design for the FD-1 and FD-2 modules following the HD principle. Figure taken from Ref. [1].

742 a hadron beam and cosmic rays, providing high-quality data sets for calibration studies
 743 and proving the excellent performance of this design.

744 Each FD HD detector module is divided in four drift regions, with a maximum drift
 745 length of 3.5 m, by alternating anode and cathode walls. The surrounding field cage
 746 ensures the uniformity of the 500 V/cm horizontal electric field across the drift volumes.
 747 The three anode walls, which constitute the charge readout of the detector, are built by
 748 stacking anode plane assemblies (APAs), 2 high times 25 wide. The design of the HD
 749 modules is shown in Fig. 3.3.

750 Each APA is made of 2560 active wires arranged in three layers, plus an extra grid
 751 layer, wrapped around a metal frame. The two induction wire planes, U and V, sit at
 752 $\pm 35.7^\circ$ to the vertical on each side of the APA. The collection and shielding plane wires,
 753 X and G, run parallel to the vertical direction. The ionisation electrons drift past the
 754 induction planes, generating bipolar signals on those wires, and are collected by the
 755 collection plane, producing a monopolar positive signal. The spacing between the wires
 756 is ~ 5 mm, and it defines the spatial resolution of the APA.

757 The front-end readout electronics, or cold electronics as they are immerse in the LAr,

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.4: Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G. Figure taken from Ref. [1].

758 are attached to the top of the up APAs and the bottom of the down APAs. Mounted on
 759 the front-end mother boards we have a series of ASICs that digitize the signals from the
 760 collection and induction planes. Each wire signal goes to a charge-sensitive amplifier,
 761 then there is a pulse-shaping circuit and this is followed by the analogue-to-digital
 762 converter. This part of the process happens inside the LAr to minimise the number of
 763 cables penetrating the cryostat. The digitised signals come out finally via a series of
 764 high-speed serial links to the warm interface boards (WIBs), from where the data is sent
 765 to the back-end DAQ through optical fibers.




Figure 3.5: A PDS module containing 24 X-ARAPUCAs (left) and the location of the modules on the APAs (right). Figure taken from Ref. [1].

3.3. Far Detector




Figure 3.6: Proposed design for the FD-3 module following the VD principle. Figure adapted from Ref. [47].

The PDS uses modules of X-ARAPUCA devices, mounted on the APA frames between the wire planes. Each X-ARAPUCA consists of layers of dichroic filter and wavelength-shifter. They shift the VUV scintillation light into the visible spectrum, sending then the visible photons to silicon photomultiplier (SiPM) devices. The PDS modules are $209\text{ cm} \times 12\text{ cm} \times 2\text{ cm}$ bars, containing 24 X-ARAPUCAs. There are 10 of these PDS modules per APA. Fig. 3.5 shows a PDS module (left) and the placement of the modules on the APAs (right).

3.3.2 Vertical Drift

In the VD case the ionisation electrons will drift vertically until they meet a printed circuit board-based (PCB) readout plane. It is based on the original dual-phase (DP) design deployed at CERN, known as ProtoDUNE-DP, used a vertical drift design with an additional amplification of the ionization electrons using a gaseous argon (GAr) layer above the liquid phase. The VD module incorporates the positive features of the DP design without the complications of having the LAr-GAr interface.

The current design of the FD VD module counts with two drift chambers with a maximum drift distance of 6.5 cm. A cathode plane splits the detector volume along the

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.7: Schematic representation of the electrode strip configuration for a top (left) and bottom (right) CRU. Figure taken from Ref. [47].

drift direction while the two anode planes are connected to the bottom and top walls of the detector. The layout of the VD module is shown in Fig. 3.6. Compared with the HD design, the VD option offers a slightly larger instrumented volume and a more cost-effective solution for the charge readout.

As in the HD design, each drift volume features a 500 V/cm electric field and a field cage that ensures its uniformity. The anode planes are arrays of $3.4\text{ m} \times 3\text{ m}$ charge-readout planes (CRPs). These are formed by a pair of charge-readout units (CRUs), which are built from two double-sided perforated PCBs, with their perforations aligned. The perforations allow the drift electrons to pass between the layers.

The PCB face opposite to the cathode has a copper guard plane which acts as shielding, while its reverse face is etched with electrode strips forming the first induction plane. The outer PCB has electrode strips on both faces, the ones facing the inner PCB form the second induction plane while the outermost ones form the collection plane. Fig. 3.7 shows the layout of the electrode strips for the top (left) and bottom (right) CRUs.

3.3. Far Detector




Figure 3.8: Detailed diagram of the DUNE FD DAQ system. Figure taken from Ref. [2].

796 The magenta and blue lines represent the first and second induction planes respectively,
 797 and the green lines correspond to the collection plane.

798 The PDS in the VD module will use the same X-ARAPUCA technology developed
 799 for the HD design. The plan is to place the PDS modules on the cryostat walls and on
 800 the cathode, in order to maximise the photon yield.

801 **3.3.3 FD Data Acquisition System**

802 The task of the data acquisition (DAQ) system is to receive, process and store data from
 803 the detector modules. In the case of DUNE the DAQ architecture is designed to work
 804 for all FD modules interchangeably, except some aspects of the upstream part which
 805 may depend on the specific module technology.

806 The enormous sample rate and the number of channels in TPC and PD readouts
 807 will produce a very large volume of data. These pose really strong requirements and
 808 challenges to the DUNE FD DAQ architecture. It will be required to read out data of
 809 the order of ten thousand or more channels at rates of a few MHz. In order to cope
 810 with the huge data volume, segmented readouts and compression algorithms are used to
 811 reduce the data rate to manageable levels.

Chapter 3. The Deep Underground Neutrino Experiment

812 The DAQ system of the DUNE FD is composed of five different subsystems. The
813 first one is the upstream DAQ, which receives the raw data from the detector, buffers it
814 and perform some low-level pre-processing. The minimally processed data is then fed
815 into a hierarchical data selection system, which then performs a module level trigger
816 decision. In case of a positive decision a trigger command is produced and executed by
817 the data flow orchestrator, located in the back-end (BE) DAQ subsystem. Subsequently
818 the DAQ BE retrieves the relevant data from the buffers located in the upstream DAQ,
819 adds all the data into a cohesive record and saves it to permanent storage. Watching
820 over all the other subsystems we also have the control, configuration and monitoring
821 subsystem and the time and synchronization subsystem. Fig. 3.8 shows a schematic
822 diagram of the DAQ system, showing the different subsystems and their relations.

823 A notorious challenge for the DUNE DAQ system comes from its broad physics
824 goals. We must be prepared to process events spanning a wide range of time windows
825 (from 5 ms in the case of beam and cosmic neutrinos and nucleon decay to 100 s in the
826 case of SNBs) and therefore this requires a continuous readout of the detector modules.
827 Moreover, because of the off-beam measurements we need to ensure the capabilities
828 of online data processing and self-triggering. Having this into account, together with
829 the technical constraints, the DUNE FD DAQ faces a series of challenges: it needs to
830 be fault tolerant and redundant to reduce downtime, accommodate new components
831 while it keeps serving the operational modules, have large upstream buffers to handle
832 SNB physics, be able to support a wide range of readout windows and last reduce the
833 throughput of data to permanent storage to be at most 30 PB/year.

834 3.4 Near Detector

835 In order to estimate the oscillation parameters we measure the neutrino energy spectra
836 at the FD. This reconstructed energy arises from a convolution of the neutrino flux, cross
837 section, detector response and the oscillation probability. Using theoretical and empirical
838 models to account for the other effects, one can extract the oscillation probability using

3.4. Near Detector




Figure 3.9: Representation of the ND hall in Phase II, showing the different subcomponents. From right to left, in the direction of the beam, we have ND-LAr, ND-GAr and SAND. Figure taken from Ref. [48].

839 the measurement. However, these models have associated a number of uncertainties that
 840 are then propagated to the oscillation parameters.

841 One of the main roles of the ND is to measure the neutrino interaction rates before
 842 the oscillation effects become relevant, i.e. close to the production point. By measuring
 843 the ν_μ and ν_e energy spectra, and that of their corresponding antineutrinos, at the ND
 844 we can constrain the model uncertainties. A complete cancellation of the uncertainties
 845 when taking the ratio between the FD and ND measurements is not possible, as that
 846 would require both detectors to have identical designs and the neutrino fluxes to be
 847 the same. Because of the distance, the flux probed by the FD will have a different
 848 energy and flavour composition than that at the ND, as neutrinos oscillate and the beam
 849 spreads. The differences in the flux also determine the design of the detectors, therefore
 850 the ND is limited in its capability to match the FD design.

851 Nevertheless, having a highly capable ND DUNE can minimise the systematic
 852 uncertainties affecting the observed neutrino energy. The ND data can be used to
 853 tune the model parameters by comparison with the prediction. Then, one uses the
 854 tuned model to predict the unoscillated FD spectra. Comparing the prediction with the

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.10: Schematic representation of the external components of ND-LAr, including the cryostat and the PRISM movable system (left) and detailed drawing of one ArgonCube module (right). Figure adapted from Ref. [1].

measured spectra it is possible to extract the oscillation parameters.

Additionally, the ND will have a physics program of its own. In particular, it will measure neutrino cross sections that will then be used to constrain the model used in the long-baseline oscillation analysis. It will also be used to search for BSM phenomena such as heavy neutral leptons, dark photons, millicharged particles, etc.

The DUNE ND can be divided in three main components, a LArTPC known as ND-LAr, a magnetised muon spectrometer, which will be the Temporary Muon Spectrometer (TMS) in Phase I and ND-GAr in Phase II, and the System for on-Axis Neutrino Detection (SAND). The layout of the Phase II DUNE ND can be seen in Fig. 3.9. The first two components of the ND will be able to move off-axis, in what is called the Precision Reaction-Independent Spectrum Measurement (PRISM) concept. More details on the purpose and design of the ND can be found in the DUNE ND Conceptual Design Report (CDR) [48].

3.4.1 ND-LAr

ND-LAr is a LArTPC, as the ND needs a LAr component in order to reduce cross section and detector systematic uncertainties in the oscillation analysis. However, its

3.4. Near Detector




Figure 3.11: Schematic view of the TMS detector, highlighting its main parts. Figure adapted from Ref. [1].

871 design differs significantly from those proposed for the FD modules. Because of the
 872 high event rates at the ND, approximately 55 neutrino interaction events per $10 \mu\text{s}$ spill,
 873 ND-LAr will be built in a modular way. Each of the modules, based on the ArgonCube
 874 technology, is a fully instrumented, optically isolated TPC with a pixelated readout.
 875 The pixelisation allows for a fully 3D reconstruction and the optical isolation reduces
 876 the problems due to overlapping interactions. Fig. 3.10 shows a representation of the
 877 external parts of ND-LAr (left) and a detailed diagram of an ArgonCube module (right).

878 With a fiducial mass of 67 t and dimensions $7 \text{ m} (\text{w}) \times 3 \text{ m} (\text{h}) \times 5 \text{ m} (\text{l})$, ND-LAr
 879 will be able to provide high statistics and contain the hadronic systems from the beam
 880 neutrino interactions, but muons with a momentum higher than 0.7 GeV will exit the
 881 detector.

882 **3.4.2 TMS/ND-GAr**

883 In order to accurately estimate the neutrino energy, the momentum of the outgoing
 884 muons needs to be determined. That is the reason why a muon spectrometer is needed
 885 downstream of ND-LAr.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.12: Cross section of the ND-GAr geometry, showing the HPgTPC, ECal and magnet. Figure adapted from Ref. [1].

886 In Phase I that role will be fulfilled by TMS. It is a magnetised sampling calorimeter,
887 with alternating steel and plastic scintillator layers. Fig. 3.11 shows a schematic view of
888 the TMS detector. The magnetic field allows a precise measurement of the sign of the
889 muon, so one can distinguish between neutrino and antineutrino interactions.

890 After the Phase II upgrade, TMS will be replaced with ND-GAr. This detector is
891 a magnetised, high-pressure GAr TPC (often denoted as HPgTPC) surrounded by an
892 electromagnetic calorimeter (ECal) and a muon tagger. A cross section of its geometry
893 can be seen in Fig. 3.12. ND-GAr will be able to measure the momenta of the outgoing
894 muons while also detect neutrino interactions inside the GAr volume. This allows
895 ND-GAr to constrain the systematic uncertainties even further, as it will be able to
896 accurately measure neutrino interactions at low energies thanks to the lower tracking
897 thresholds of GAr.

898 3.4.3 PRISM

899 In general, the observed peak neutrino energy of a neutrino beam decreases as the
900 observation angle with respect to the beam direction increases. This feature has been
901 used in other long-baseline neutrino experiments, like T2K (2.5° off-axis) and NOvA

3.4. Near Detector




Figure 3.13: Predicted beam muon neutrino flux at the ND location for different off-axis positions. Figure taken from Ref. [48].

902 (0.8° off-axis), in order to achieve narrower energy distributions. The DUNE PRISM
 903 concept exploits this effect using a movable ND. Within PRISM both ND-LAr and the
 904 muon spectrometer (TMS in Phase I and ND-GAr in Phase II) can be moved up to 3.2°
 905 off-axis, equivalent to move the detectors 30.5 m laterally through the ND hall.

906 This allows to record additional data samples with different energy compositions.
 907 Fig. 3.13 compares the on-axis muon neutrino flux at the ND with the fluxes at different
 908 off-axis positions. As the off-axis position increases the neutrino flux becomes closer to
 909 a monoenergetic beam with a lower peak energy. These samples can be used to perform
 910 a data-driven determination of the relation between true and reconstructed neutrino
 911 energy, in order to reduce the dependence on the interaction model. The off-axis samples
 912 are linearly combined to produce a narrow Gaussian energy distribution centered on
 913 a target true energy. From the combination coefficients one can build a sample of
 914 reconstructed neutrino events that will determine the energy mapping.

915 The PRISM samples can also be used to form a flux at the ND location similar in
 916 shape to the oscillated flux measured by the FD. This method can be used to extract
 917 the oscillation parameters with minimal input from the neutrino interaction model.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.14: Schematic longitudinal section of the LBNF beamline at Fermilab (not to scale). Figure taken from Ref. [49].

918 3.4.4 SAND

919 The role of SAND is to monitor the beam stability by measuring the on-axis neutrino
 920 energy spectra. As the PRISM program requires that ND-LAr and its downstream
 921 muon spectrometer spend about half of the time in off-axis positions, it is not possible
 922 to monitor the stability with the movable detectors. Moreover, for the success of PRISM
 923 it is essential to have a stable beam configuration, or, at least, a quick assessment and
 924 modeling of the distortions.

925 The SAND detector is magnetised, and it counts with an inner low density tracker,
 926 a LAr target with optical readout and surrounding sampling calorimeter.

927 3.5 LBNF beamline

928 The Long-Baseline Neutrino Facility (LBNF) project is responsible for producing the
 929 neutrino beam for the DUNE detectors. A detailed discussion of the LBNF program
 930 can be found in the DUNE/LBNF CDR Volume III [49].

931 The LBNF beamline will provide a high-intensity neutrino beam within the adequate
 932 energy range in order to meet the long-baseline oscillation physics goals of DUNE. A
 933 schematic diagram of the longitudinal section of the LBNF beamline is shown in Fig.
 934 3.14. First, a beam of $60 - 120$ GeV protons is extracted from the Fermilab Main

3.5. LBNF beamline

935 Injector. This beam is aimed towards the target area, where it collides with a cylindrical
 936 graphite target to produce pions and kaons.

937 The diffuse, secondary beam of particles is focused by a pair of magnetic horns.
 938 These select the positively charged particles when operated in Forward Horn Current
 939 (FHC) mode, or the negatively charged ones when the current is reversed, also known as
 940 Reverse Horn Current (RHC) mode. The focused secondary beam then enters a 194 m
 941 decay pipe where the pions and kaons will predominantly produce $\mu^+\nu_\mu$ pairs when in
 942 FHC mode (or $\mu^-\bar{\nu}_\mu$ in RHC mode).

943 At the end of the decay pipe a hadron absorber removes the undecayed hadrons and
 944 muons from the beam, which reduces the ν_e ($\bar{\nu}_e$) and $\bar{\nu}_\mu$ (ν_μ) contamination coming
 945 from the μ^+ (μ^-) decays. The resulting neutrino flux at the FD is shown in Fig. 3.15,
 946 both for FHC (left) and RHC (right) modes. These predictions show the intrinsic $(\bar{\nu}_e)$
 947 contamination and wrong sign component from wrong sign and neutral meson decays,
 948 as well as muons decaying before reaching the absorber.




Figure 3.15: Predicted neutrino fluxes at the FD in FHC mode (left panel) and RHC mode (right panel). Figures taken from Ref. [42].

949 **Chapter 4**

950 **ND-GAr**

951 ND-GAr is a magnetised, high-pressure gaseous argon TPC (HPgTPC), surrounded by
952 an electromagnetic calorimeter (ECal) and a muon detector (commonly refer to as μ ID).
953 A detailed discussion on the requirements, design, performance and physics of ND-GAr
954 can be found in the DUNE ND CDR [48] and the ND-GAr whitepaper (cite).

955 In DUNE Phase II ND-GAr will fulfill the role of TMS, measuring the momentum
956 and sign of the charged particles exiting ND-LAr. Additionally, it will be able to measure
957 neutrino interactions inside the HPgTPC, achieving lower energy thresholds than those
958 of the ND and FD LArTPCs. By doing so ND-GAr will allow to constrain the relevant
959 systematic uncertainties for the LBL analysis even further.

960 The goal of the present chapter is to review the requirements that the physics program
961 of DUNE impose on ND-GAr, present the current status of its design and describe the
962 GArSoft package, its simulation and reconstruction software.

963 **4.1 Requirements**

964 The primary requirement for ND-GAr is to the measure the momentum and charge of
965 muons from ν_μ and $\bar{\nu}_\mu$ CC interactions in ND-LAr, in order to measure their energy
966 spectrum. To achieve the sensitivity to the neutrino oscillation parameters described
967 in the DUNE FD TDR Volume II [42] ND-GAr should be able to constrain the muon

Chapter 4. ND-GAr

968 energy within a 1% uncertainty or better. The main constraint will come from the
969 calibration of the magnetic field, performed using neutral kaon decays in the HPgTPC.

970 Another requirement for ND-GAr is the precise measurement of neutrino interactions
971 on argon for the energies relevant to the neutrino oscillation program. The goal is to
972 constrain the cross section systematic uncertainties in the regions of phase space that
973 are not accessible to ND-LAr. This requires the kinematic acceptance for muons in
974 ND-GAr to exceed that of ND-LAr, being comparable to the one observed in the FD.

975 ND-GAr should also be able to the relationship between true and reconstructed energy
976 from neutrino interactions on argon with low thresholds, being sensitive to particles that
977 are not observed or may be misidentified in ND-LAr. In particular, ND-GAr needs to
978 have low tracking thresholds in order to measure the spectrum of pions and protons
979 produced in final-state interactions (FSI). It also must be able to accurately measure
980 the pion multiplicity in 1, 2 and 3 pions final states, to inform the pion mass correction
981 in the LArTPCs.

982 4.2 Reference design

983 The final design of ND-GAr is still under preparation. However, a preliminary baseline
984 design was in place at the time of the ND CDR. This section summarises the main
985 features of that design, as it is also the one used for the default geometry in our simulation.
986 A DUNE Phase II whitepaper, discussing the different options under consideration for
987 the ND-GAr design, is in progress.

988 4.2.1 HPgTPC

989 The reference design for the ND-GAr HPgTPC follow closely that of the ALICE TPC.
990 It is a cylinder with a central high-voltage cathode, generating the electric field for
991 the two drift volumes, with a maximum drift distance of 2.5 m each. The anodes will
992 be instrumented with charge readout chambers. The original design repurposed the
993 multi-wire proportional readout chambers of ALICE, however the current R&D efforts

4.2. Reference design




Figure 4.1: Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers. Figure taken from Ref. [1].

994 focus on a gas electron multiplier option instead. Fig. 4.1 shows a schematic diagram of
 995 the ALICE TPC design. The basic ND-GAr geometry will resemble this, except for the
 996 inner field cage.

997 It will use a 90-10 molar fraction argon-CH₄ mixture at 10 bar. With this baseline
 998 gas mixture light collection is not possible, as the quenching gas absorbs most of the
 999 VUV photons. Additional R&D efforts are underway, to understand if different mixtures
 1000 allow for the light signal to be used to provide a t_0 while maintaining stable charge gain.

1001 4.2.2 ECal

1002 The main role of the ND-GAr ECal is the calorimetric measurement of the electron
 1003 energies and the reconstruction of photons, in particular those from neutral pion decays.
 1004 Also, the ECal is able to provide a t_0 timestamp for neutrino interactions, by associating
 1005 its activity to the tracks in the HPgTPC. The ECal will also be able to perform
 1006 neutron reconstruction using time of flight and reject external backgrounds, thanks to
 1007 its sub-nanosecond time resolution.

1008 The ECal design features three independent subdetectors, two end caps at each side
 1009 and a barrel surrounding the HPgTPC. Each of the detectors is divided in modules,

Chapter 4. ND-GAr




Figure 4.2: View of the 12-sided ECal barrel and outer muon tagger geometries (left) and layout of the ECal tile layers for the 2 mm Cu, 10 mm scintillator option (right). Figure adapted from Ref. [1].

which combine alternating layers of plastic scintillator and absorber material readout by SiPMs. The inner scintillator layers consist of $2.5 \times 2.5 \text{ cm}^2$ high-granularity tiles, whereas the outer ones are made out of 4 cm wide cross-strips spanning the whole module length. The current barrel geometry consists of 8 tile layers and 34 strip layers, while the end caps feature 6 and 36 respectively. The thickness of the scintillator layers is 7 mm and 5 mm for the Pb absorber layers. The 12-sided geometry of the ECal barrel (left) and the layout of the tile layers (left)¹ can be seen in Fig. 4.2.

4.2.3 Magnet

The ND-GAr magnet design, known as the Solenoid with Partial Yoke (SPY), consists of two coupled solenoids with an iron return yoke. The idea behind the design is to have a solenoid as thin as possible, as well as a return yoke mass distribution that minimises the material budget between ND-LAr and ND-GAr. The magnet needs to provide a 0.5 T field in the direction perpendicular to the beam, parallel to the drift electric field. It needs to host the pressure vessel and the surrounding ECal, which points to an inner diameter of $\sim 6.4 \text{ m}$.

The solenoid is a single layer coil, based on niobium titanium superconducting

¹The figure shows the layout of the tile layers for a previous design with 2 mm Cu absorber and 10 mm plastic scintillator, as mentioned in the text the current choice is 5 mm Pb absorber and 7 mm scintillator.

4.3. GArSoft

1026 Rutherford cable. The total length of the coil is 7.5 m. The bobbin will be split in four
1027 segments grouped in pairs with two identical cryostats, connected in series. The iron
1028 yoke features an aperture in the upstream side to allow the muons coming from ND-LAr.
1029 Still, its material will be enough to reduce the magnetic field reaching SAND, and also
1030 stop the charged pions produced inside the HPgTPC.

1031 4.2.4 Muon system

1032 The design of the ND-GAr muon system is still in a preliminary stage. Its role is to
1033 distinguish between muons and pions punching through the ECal. This is especially
1034 important for wrong-sign determination, to separate these from neutral current events.

1035 In its current form, the muon system consists of three layers of longitudinal sampling
1036 structures. It alternates 10 cm Fe absorber slabs with 2 cm plastic scintillator strips.

1037 The transverse granularity required is still under study.

1038 4.3 GArSoft

1039 GArSoft is a software package developed for the simulation and reconstruction of events
1040 in ND-GAr. It is inspired by the LArSoft toolkit used for the simulation of LArTPC
1041 experiments, like the DUNE FD modules. It is based on `art`, the framework for event
1042 processing in particle physics experiments [?]. Other of its main dependencies are `ROOT`,
1043 `NuTools`, `GENIE` and `Geant4`. It allows the user to run all the steps of a generation-
1044 simulation-reconstruction workflow using FHiCL configuration files.

1045 4.3.1 Event generation

1046 The standard generator FHiCLs in GArSoft run the event generation and particle
1047 propagation simulation (i.e. Geant4) in the same job by default. However, it is possible
1048 to split them up if needed. The current version of GArSoft provides five different event
1049 generators, each of them producing `simb::MCTruth` products defined in `NuTools`. The
1050 available modules are:

Chapter 4. ND-GAr

- 1051 • **SingleGen**: particle gun generator. It produces the specified particles with a given
1052 distribution of momenta, initial positions and angles.
 - 1053 • **TextGen**: text file generator. The input file must follow the `hepevt` format², the
1054 module simply copies this to `simb::MCTruth` data products.
 - 1055 • **GENIEGen**: GENIE neutrino event generator. The module runs the neutrino-nucleus
1056 interaction generator using the options specified in the driver FHiCL file (flux file,
1057 flavour composition, number of interactions per event, t_0 distribution, ...). Current
1058 default version is `v3_04_00`.
 - 1059 • **RadioGen**: radiological generator. It produces a set list of particles to model
1060 radiological decays. Not tested.
 - 1061 • **CRYGen**: cosmic ray generator. The module runs the CRY event generator with a
1062 configuration specified in the FHiCL file (latitude and altitude of detector, energy
1063 threshold, ...). Not tested.
- 1064 The module `GArG4` searches for all the generated `simb::MCTruth` data products, using
1065 them as inputs to the Geant4 simulation with the specified detector geometry. A constant
1066 0.5 T magnetic field along the drift coordinate is assumed. The main outputs of this step
1067 are `simb::MCParticle` objects for the generated Geant4 particles, `gar::EnergyDeposit`
1068 data products for the energy deposits in the HPgTPC and `gar::CaloDeposit` data
1069 products for the energy deposits in the ECal and muon system.

1070 4.3.2 Detector simulation

- 1071 The standard detector simulation step in GArSoft is all run with a single FHiCL, but
1072 the different modules can be run independently as well. First the `IonizationReadout`

²In brief, each event contains at least two lines. The first line contains two entries, the event number and the number of particles in the event. Each following line contains 15 entries to describe each particle. The entries are: status code, pdg code for the particle, entry of the first mother for this particle, entry of the second mother for this particle, entry of the first daughter for this particle, entry of the second daughter for this particle, x component of the particle momentum, y component of the particle momentum, z component of the particle momentum, energy of the particle, mass of the particle, x component of the particle initial position, y component of the particle initial position, z component of the particle initial position and time of the particle production.

4.3. GArSoft

1073 module simulates the charge readout of the HPgTPC, and later the `SiPMReadout` module
1074 runs twice, once for the ECal and then for the muon system, with different configurations.

1075 The `IonizationAndScintillation` module collects all the `gar::EnergyDeposit`
1076 data products, to compute the equivalent number of ionization electrons for each energy
1077 deposit. The `ElectronDriftAlg` module simulates the electron diffusion numerically
1078 both in the longitudinal and transverse directions and applies an electron lifetime
1079 correction factor. The induced charge on the nearest and neighbouring readout pads
1080 is modeled using the provided pad response functions. The digitisation of the data is
1081 then simulated with the `TPCReadoutSimAlg` module. By default, the ADC sampling
1082 rate used is 50.505 MHz. The resulting raw waveforms for each channel are stored with
1083 zero-suppression, in order to save memory and CPU time. The algorithms keep blocks
1084 of ADC values above a certain threshold, plus some adjustable additional early and late
1085 tick counts. The results of these three steps are `gar::raw::RawDigit` data products.

1086 For the ECal and the muon system the `SiPMReadout` module calls either the
1087 `ECALReadoutSimStandardAlg` or `MuIDReadoutSimStandardAlg` modules. These take
1088 all the `gar::CaloDeposit` data products in the corresponding detector and do the
1089 digitisation depending on whether the hit was in a tile or strip layer. They include single
1090 photon statistics, electronic noise, SiPM saturation and time smearing. The resulting
1091 objects are `gar::raw::CaloRawDigit` data products.

1092 4.3.3 Reconstruction

1093 The reconstruction in GArSoft is also run as a single job by default. It first runs the hit
1094 finding, clustering, track fitting and vertex identification in the HPgTPC, followed by
1095 the hit finding and clustering in the ECal and muon system. After those it produces the
1096 associations between the associations between the tracks and the ECal clusters.

1097 Focusing first on the HPgTPC reconstruction, the `CompressedHitFinder` module
1098 takes the zero-suppressed ADCs from the `gar::raw::RawDigit` data products. The
1099 reconstructed hits largely correspond to the above threshold blocks, however the hit
1100 finder identifies waveforms with more than one maximum, diving them in multiple hits

Chapter 4. ND-GAr

1101 if they dip below a certain threshold. The data products produced are of the form
1102 `gar::rec::Hit`. These are the inputs to the clustering of hits in the `TPCHitCluster`
1103 module. Hits close in space and time are merged, and the resulting centroids are found.
1104 This module outputs `gar::rec::TPCClusters` objects and associations to the input
1105 hits.

1106 The following step prior to the track fitting is pattern recognition. The module
1107 called `tpcvecchitfinder2` uses the `gar::rec::TPCClusters` data products to find track
1108 segments, typically called vector hits. They are identified by performing linear 2D fits
1109 to the positions of the clusters in a 10 cm radius, one fit for each coordinate pair. A
1110 3D fit defines the line segment of the vector hit, using as independent variable the one
1111 whose sum of (absolute value) slopes in the 2D fits is the smallest. The clusters are
1112 merged to a given vector hit if they are less than 2 cm away from the line segment. The
1113 outputs are `gar::rec::VecHit` data products, as well as associations to the clusters. The
1114 `tpcpatrec2` module takes the `gar::rec::VecHit` objects to form the track candidates.
1115 The vector hits are merged together if their direction matches, their centers are within
1116 60 cm and their direction vectors point roughly to their respective centers. Once
1117 the clusters of vector hits are formed they are used to make a first estimation of the
1118 track parameters, simply taking three clusters along the track. The module produces
1119 `gar::rec::Track` data products and associations between these tracks and the clusters
1120 and vector hits.

1121 The track is fitted by means of a Kalman filter in the `tpctrackfit2` module, using
1122 the position along the drift direction as the independent variable. Two different fits are
1123 performed per track, a forward and a backwards fit, each starting from one of the track
1124 ends. The Kalman filter state vector ($y, z, R, \phi, \tan\lambda$) is estimated at each point along
1125 the track using a Bayesian update. The track parameters reported in the forward and
1126 backwards fits are the ones computed at the opposite end where the fit started. The
1127 main outputs of the track fit are the `gar::rec::Track` objects. Additionally, the module
1128 stores the fitted 3D positions along the track in the `gar::rec::TrackTrajectory` data
1129 products and the total charge and step sizes for each point also get stored in the form of

4.3. GArSoft

1130 `gar::rec::TrackIonization` objects.

1131 After the tracking step, the `vertexfinder1` module looks at the reconstructed
1132 `gar::rec::Track` products, creating vertex candidates with the track ends that are
1133 within 12 cm of each other. The vertices are then fitted using linear extrapolations from
1134 the different track ends associated. The results are `gar::rec::Vertex` data products,
1135 and associations to the tracks and corresponding track ends.

1136 For the ECal and muon tagger, the `SiPMHitFinder` module runs twice with different
1137 configurations, adapted to the particular capabilities of both. The module simply takes
1138 the `gar::raw::CaloRawDigit` products, applies a calibration factor to convert the ADC
1139 counts to MeV and for the strip layer hits it calculates the position along the strip using
1140 the times recorded of both SiPMs. This module produces `gar::rec::CaloHit` data
1141 products. Next, these objects are used as inputs to the `CaloClustering` module. It
1142 merges the hits based on a simple nearest neighbours (NN) algorithm. For the resulting
1143 clusters it also computes the total energy and position of the centroid. The results are
1144 stored as `gar::rec::Cluster` data products, with associations to the hits.

1145 The last step in the reconstruction is associating the reconstructed tracks in the
1146 HPgTPC to the clusters formed in the ECal and muon system. The `TPCECALAssociation`
1147 module checks first the position of the track end points, considering only the points
1148 that are at least 215 cm away from the cathode or have a radial distance to the center
1149 greater than 230 cm. The candidates are propagated up to the radial position, in the
1150 case of clusters in the barrel, or the drift coordinate position, for the end cap cluster, of
1151 the different clusters in the collection using the track parameters computed at the end
1152 point. The end point is associated to the cluster if certain proximity criteria are met.
1153 This module creates associations between the tracks, the end points and the clusters.
1154 The criteria for the associations are slightly different for the ECal and the muon tagger.

₁₁₅₅ Chapter 5

₁₁₅₆ FWTPG offline software

1157 Chapter 6

1158 Matched Filter approach to
1159 induction wire Trigger Primitives

1160 6.1 Motivation

1161 The filter implemented in the firmware of the upstream DUNE FD DAQ is a 32nd-order
1162 low-pass finite impulse-response (FIR) filter. The output of such filter for a discrete
1163 system can be written as:

$$y[i] = \sum_{j=0}^N h[j]x[i-j], \quad (6.1)$$

1164 where N is the order of the filter, y is the output sequence, x is the input sequence and h
1165 is the set of coefficients of the filter. The current implementation within `dtp-firmware`
1166 [50] uses a set of 16 non-zero integer coefficients.

1167 Filtering is a vital step in the hit finder chain. It helps to suppress the noise and
1168 enhance the signal peaks with respect to the noiseless baseline. A good filtering strategy
1169 allows us to use lower thresholds when forming the trigger primitives (TPs) and thus
1170 increasing the sensitivity of our detector to low energy physics events. In such events,
1171 the hits produced by the ionisation electrons tend to have lower amplitudes than those
1172 of interest to the baseline physics programme of the DUNE experiment.

1173 This is particularly important for the induction planes. In general, signal peaks in

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.1: Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G.

the induction wires have smaller amplitude than the ones in the induction plane. This, together with the fact that the pulse shapes are bipolar, reduces our capacity to detect the hits on these channels. The inefficiency of detecting TPs in the induction planes (denoted as U and V planes) lead trigger algorithms to focus mainly on the TPs from the collection plane (so-called X plane). As a result, the possibility of making trigger decisions based on the coincidence of TPs across the three wire planes remains nowadays unexploited in DUNE. Fig. 6.1 shows a schematic view of an anode plane assembly (APA), with the different wire plane orientations highlighted.

A possible improvement of the current hit finder chain could require optimising the existing or choosing a new filter implementation. A filter strategy which improves the induction signals may be able to enhance the detection efficiency of TPs from the induction planes and ideally make it comparable to that of the collection plane.

The goal is to implement a better finite-impulse response filter design and to evaluate its performance relative to the current filter. To do so, we need to take into account the limitations of the firmware: the FIR filter shall have maximum 32 coefficients (so-called taps) whose values are 12-bit unsigned integers. Although it is technically possible to include non-integer coefficients, it would be a technical challenge as we have 40 FIR

6.2. Signal-to-noise ratio definition




Figure 6.2: Left panel: Zoomed unfiltered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44` (blue line). The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{noise}}$.

1191 instances per APA, as there are 4 FIR per optical link and 10 optical links per APA.
 1192 With these restrictions, the task is to provide a set of 32 coefficients which yield an
 1193 optimal filter performance for the induction wires.

1194 6.2 Signal-to-noise ratio definition

1195 I introduce the signal to noise ratio (S/N) as a measure of the FIR filter performance
 1196 and demonstrate how to extract its value for a set of ProtoDUNE-SP data. The S/N
 1197 metrics allow us to compare different filter implementations and serve as a basis for more
 1198 detailed studies presented later in this document. Specifically, I use the ADC capture
 1199 `felix-2020-07-17-21:31:44` (data capture taken for firmware validation purposes). I
 1200 defined S/N as the height of the signal peaks relative to the size of the noise peaks.
 1201 To quantify this quantity channel by channel one first need to estimate the standard
 1202 deviation of the ADC data for each channel, σ_{ADC} . Then, I define the corresponding

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

noise waveform to be the ADC values in the range $\pm 3\sigma_{ADC}$. From this new noise data one can estimate again the mean and standard deviation, μ_{noise} and σ_{noise} , so I can write the S/N for any given channel as:

$$S/N = \frac{\max [ADC] - \mu_{noise}}{\sigma_{noise}}, \quad (6.2)$$

where $\max [ADC]$ is simply the maximum ADC value found in the corresponding channel.

One can apply this definition of the S/N with a waveform from one of the channels of the data capture¹. Fig. 6.2 shows a zoomed region of the waveform corresponding to channel 7840 (blue line), where one can clearly see three signal peaks and continuous additive noise (we actually see 6 peaks, 3 positive and 3 negative, but, because by design for induction channels the expected signal pulse shapes are bipolar, I treat them as a collection of 3 individual signal peaks). I estimated the standard deviation of this raw waveform to be $\sigma_{raw} = 7.84$ ADC, so I am able to define the noise waveform (red line) as the ADC values in the range ± 23.52 ADC. This way one obtains $\mu_{noise} = 0$ and $\sigma_{noise} = 5.58$ ADC, which gives $S/N = 30.84$.

We can repeat this calculation now for the corresponding filtered waveform (using the current firmware FIR filter). In Fig. 6.3 I plotted the same time window for the filtered waveform from channel 7840 (blue line). In this case, the standard deviation of the waveform is larger than before, giving $\sigma_{raw} = 10.99$ ADC. The resulting noise waveform (red line) results from selection the ADC values in the range ± 32.91 ADC, giving now $\mu_{noise} = -0.47$ ADC and $\sigma_{noise} = 7.03$ ADC. Finally, one obtains $S/N = 24.68$. Notice that the value of S/N decreases after the filtering. Clearly, one can see that the noise baseline has increased by a factor of 1.35 when we applied the FIR filter and at the same time the amplitude of the signal peaks has remained almost unchanged, leading to this poorer S/N value.

¹All the original work was done within the `dtp-simulation` package [51], which offers a variety of tools to read raw data and emulate the TPG block (pedestal subtraction, filtering and hit finder). However, the results shown in this report were re-worked later using the C++ based `dtpemulator` package [52]. Its main purpose is the emulation of the TPG block and, in the same way as its predecessor, it has been cross-checked against the current firmware implementation.

6.3. Low-pass FIR filter design




Figure 6.3: Left panel: Zoomed filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44` (blue line). The filter used was the current implementation of the low-pass FIR filter in `dtp-firmware`. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$

1226 6.3 Low-pass FIR filter design

1227 In general, when one uses a method to optimize the frequency response of a digital filter,
 1228 such as the Parks-McClellan algorithm, one finds a set of N real coefficients that give
 1229 the best response for the specified pass-band and order of the filter [53].

1230 In our case, as the sampling frequency is defined as 1 ticks^{-1} , the Nyquist frequency
 1231 will simply be $1/2 \text{ ticks}^{-1}$. The current implementation of the filter seems to have as
 1232 pass-band the range $[0, 0.1] \text{ ticks}^{-1}$. This can be seen in Fig. 6.4, where I show the
 1233 power spectrum, in decibels, of such filter implementation (blue solid line). For instance,
 1234 the Park-McClellan algorithm finds the optimal Chebyshev FIR filter taking as input
 1235 the boundaries of the target pass-band and stop-band, which can be written in the form:

$$\left\{ \begin{array}{l} [0, f_c] \\ [f_c + \delta f, f_N] \end{array} \right. , \quad (6.3)$$

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

where f_c is the cut-off frequency, δf is the transition width and f_N is the aforementioned Nyquist frequency. A similar behaviour to the one in the current filter can be obtained by setting $f_c = 0$ and $\delta f = 0.1 \text{ ticks}^{-1}$. The response of the resulting filter is also shown in Fig. 6.4 (blue solid line). Notice that the suppression of the stop-band is enhanced for this optimal filter. For comparison I included the power response of the filter obtained by taking the integer part of the coefficients resulting from the Parks-McClellan method (red dashed line). One can see that it does not suppress that much the stop-band, in a similar way to the current implementation of the filter.




Figure 6.4: Power spectrum in decibels for the current implementation of the low-pass FIR filter in `dtp-firmware` (blue line), compared to the response of an optimal filter obtained using the Parks-McClellan algorithm for the same pass-band (red line). Also for comparison I include the spectrum of the optimal filter when taking only the integer part of the coefficients (red dashed line).

At this point, I tried to improve the performance of the FIR filter using the Parks-McClellan method, i.e. maximize the overall S/N, using the available data captures. I did so by varying the values of the two quantities that parametrize the pass-band and stop-band, the cut-off frequency f_c and the transition width δf .

Fig. 6.5 shows the average relative change in the S/N (i.e. the ratio between the value of the S/N after and before the filtering) for capture `felix-2020-07-17-21:31:44`,

6.3. Low-pass FIR filter design




Figure 6.5: Relative change in the S/N for the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44`, using different values of the cutoff frequency f_c and the transition width δf . The optimal Chebyshev filters were applied using just the integer part of the coefficients given by the Parks-McClellan algorithm.

when using filters designed with the Parks-McClellan algorithm for the specified values of the cut-off frequency f_c and the transition width δf , restricted to integer values for the filter coefficients. One can clearly distinguish different regions where we get an improvement of up to a factor of 1.35 for the U plane. For large values of $f_c + \delta f$ the ratio tends to 1, as expected (in that limit the width of the stop-band goes to 0, meaning that no frequencies are filtered out and thus the waveform remains the same).

Using the configuration which gives the best mean performance for the three planes (see bottom right panel of Fig. 6.5), i.e. $f_c = 0.068 \text{ ticks}^{-1}$ and $\delta f = 0.010 \text{ ticks}^{-1}$, we can see how such filter affects the different channels. Fig. 6.6 shows the distribution of the S/N improvement values for all the channels in the raw ADC capture `felix-2020-07-17-21:31:44`, separated by wire plane, after the optimal Chebyshev filter was applied. One can see that there is a clear improvement for both U and V induction wire planes, obtaining a mean change of 1.25 and 1.30 for them respectively. However, in the case of the collection plane X the mean of this distribution is roughly 1, meaning that a good fraction of channels in that plane get a slightly worse S/N after the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.6: Distribution of the relative change of the S/N on the different wire planes from the ProtoDUNE-SP raw data capture *felix-2020-07-17-21:31:44* after the optimal Chebyshev filter was applied. The filter was computed with the Parks-McClellan algorithm using a cutoff of $f_c = 0.068 \text{ ticks}^{-1}$ and a transition width $\delta f = 0.010 \text{ ticks}^{-1}$.

filter is applied. In any case, this is not a big issue as the S/N for collection channels is usually much higher than the one for induction channels.

The results I obtained optimising the low pass filter with the Parks-McClellan method are promising. Nonetheless, the improvement found is rather marginal so I wondered if there could be an alternative approach to the filtering problem which yields better outputs. At this point, I found a possible alternative in matched filters. By construction, this kind of filters offer the best improvement on the S/N.

6.4 Matched filters

In the context of signal processing, a matched filter is the optimal linear filter for maximising the signal-to-noise ratio (S/N) in the presence of additive noise, obtained by convolving a conjugated time-reversed known template with an unknown signal to detect the presence of the template in the signal [54].

Given a known signal sequence $s(t)$ and another (a priori unknown) noise sequence

6.4. Matched filters




Figure 6.7: Left panel: Zoomed match filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture felix-2020-07-17-21:31:44 (blue line). The filter used was directly extracted from the data, being the 32 values around the first peak in the original waveform. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after match filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after match filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$

1278 $n(t)$, the input signal can be written as:

$$x(t) = s(t) + n(t). \quad (6.4)$$

1279 Now, considering a linear time-invariant filter, whose impulse-response function I
1280 will refer to as $h(t)$, one can write the output signal as:

$$\begin{aligned} y(t) &= x(t) * h(t) \\ &= (s(t) + n(t)) * h(t) \\ &= y_s(t) + y_n(t), \end{aligned} \quad (6.5)$$

1281 where $y_s(t)$ and $y_n(t)$ are simply the outputs of the filter due to the signal and the noise
1282 components respectively.

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1283 The goal of the matched filter is to detect the presence of the signal $s(t)$ in the input
 1284 sample $x(t)$ at a certain time t_0 , which effectively means we need to maximise the S/N.
 1285 This way, what one wants is to have a filter which gives a much bigger output when the
 1286 known signal is present than when it is not. Putting it in other words, the instantaneous
 1287 power of the signal output $y_s(t)$ should be much larger than the average power of the
 1288 noise output $y_n(t)$ at some time t_0 .

1289 For the case of the filtered signal, one can easily re-write it as an inverse Fourier
 1290 transform:

$$y_s(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega H(\omega)S(\omega)e^{i\omega t}, \quad (6.6)$$

1291 where $H(\omega)$ and $S(\omega)$ are the Fourier transforms of the impulse-response function (i.e.
 1292 the transfer function of the filter) and of the input signal, respectively.

1293 Now focusing on the noise, we can use the Wiener-Khinchin theorem [55] to write
 1294 the mean power of the noise after filtering as:

$$E|y_n(t)|^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega |H(\omega)|^2 S_n(\omega), \quad (6.7)$$

1295 where $S_n(\omega)$ is the power spectral density of the noise.

1296 Having these, one can write the instantaneous S/N at time t_0 as:

$$\begin{aligned} \left(\frac{S}{N} \right)_{t_0} &= \frac{|y_s|^2}{E|y_n(t)|^2} \\ &= \frac{1}{2\pi} \frac{\left| \int_{-\infty}^{\infty} d\omega H(\omega)S(\omega)e^{i\omega t_0} \right|^2}{\int_{-\infty}^{\infty} d\omega |H(\omega)|^2 S_n(\omega)}. \end{aligned} \quad (6.8)$$

1297 Once we have this expression, we need to find the upper limit of it to determine what
 1298 would be the optimal choice for the transfer function. One can use the Cauchy-Schwarz
 1299 inequality, which in the present case takes the form:

$$\left| \int_{-\infty}^{\infty} dx f(x)g(x) \right|^2 \leq \int_{-\infty}^{\infty} dx |f(x)|^2 + \int_{-\infty}^{\infty} dx |g(x)|^2, \quad (6.9)$$

6.4. Matched filters

1300 for any two analytical functions $f(x)$ and $g(x)$. One can prove that making the choice:

$$\begin{aligned} f(x) &= H(\omega) \sqrt{S_n(\omega)} e^{i\omega t_0}, \\ g(x) &= \frac{S(\omega)}{\sqrt{S_n(\omega)}}, \end{aligned} \quad (6.10)$$

1301 leads to the following upper bound for the S/N:

$$\left(\frac{S}{N} \right)_{t_0} \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{|S(\omega)|^2}{S_n(\omega)}. \quad (6.11)$$

1302 From Eqs. (6.8), (6.9) and (6.10) one can also derive the form of the transfer function

1303 such that the upper bound is exactly reached [56]:

$$H(\omega) \propto \frac{S^*(\omega) e^{-i\omega t_0}}{S_n(\omega)}. \quad (6.12)$$

1304 From this last expression we can clearly see the way the matched filter acts. As the
 1305 transfer function is proportional to the Fourier transform of the signal it will try to only
 1306 pick the frequencies present in the signal [57].

1307 The matched filter transfer function can be greatly simplified if the input noise is
 1308 Gaussian. In that case, the power spectral density of the noise is a constant, so it can be
 1309 re-absorbed in the overall normalisation of the transfer function. Moreover, considering
 1310 that the input signal is a real function, one can simply set $S^*(\omega) = S(-\omega)$, which gives:

$$H(\omega) \propto S(-\omega) e^{-i\omega t_0}. \quad (6.13)$$

1311 For a discrete signal, one can think of the input and impulse-response sequences as
 1312 vectors of \mathbb{R}^N . Then, the matched filter tries to maximise the inner product of the signal
 1313 and the filter while minimising the output due to the noise by choosing a filter vector
 1314 orthogonal to the later. In the case of additive noise, that leads to the impulse-response
 1315 vector:

$$h = \frac{1}{\sqrt{s^\dagger R_n^{-1} s}} R_n^{-1} s, \quad (6.14)$$

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.8: Relative improvement in the S/N for the raw data capture `felix-2020-07-17-21:31:44`, using the matched filter following the parametrisation in Eq. (6.17). The black crosses in both panels denote the location of the maximum ratio value.

1316 where s is a reversed signal template sequence of length N equal to the order of the filter
 1317 and R_n is the covariance matrix associated with the noise sequence n . For the Gaussian
 1318 noise case, the covariance matrix is simply the unit matrix, so the above expression
 1319 simplifies again to:

$$h = \frac{s}{|s|}. \quad (6.15)$$

1320 For this first stage of the study, I use a definition of the S/N per channel given by:

$$\text{S/N} = \frac{\max [ADC] - \mu_{noise}}{\sigma_{noise}}, \quad (6.16)$$

1321 where the subscript *noise* refers to a subset of the data obtained by only taking into
 1322 account waveform values within a $\pm 3\sigma$ range around the mean of the data and $\max [ADC]$
 1323 is the maximum of the original waveform. This definition is further discussed in App.
 1324 6.2, where I also show examples of its application to raw data and to a waveform filtered
 1325 with the current low-pass FIR filter.

1326 To test whether this choice of filter is appropriate one needs to choose a signal
 1327 template. As an example of how a matched filter would affect our signal, I simply took
 1328 the filter coefficients to be the 32 ADC values around a signal peak present in the data.

6.4. Matched filters

1329 In Fig. 6.7 (left panel) I plotted a zoomed region for channel 7840 in the raw data capture
1330 `felix-2020-07-17-21:31:44`, after applying the matched filter described before (blue
1331 line). When compared to the raw and FIR filtered case (see App. 6.2), after applying
1332 the match filter the standard deviation of the noise waveform (red line) decreases and at
1333 the same time the signal peaks are enhanced. This leads to an improvement of the S/N
1334 by a factor of 1.92 when compared to the raw waveform.

1335 In order to obtain the matched filter that is more suitable for our data, I explored
1336 different configurations of signal templates. In order to perform this exploration, I
1337 parametrised the signal using the bipolar function:

$$f(x) = -A(x + \delta) e^{-x^2/\sigma^2}, \quad (6.17)$$

1338 where the parameter δ controls the asymmetry between the positive and negative peaks
1339 and σ controls their width. The amplitude parameter A is set such that it keeps the
1340 height of the biggest peak to be less than 200 ADC in absolute value.

1341 As this parametrisation is only adequate for bipolar signals I will focus exclusively
1342 on the induction channels. Also, the optimal configurations I found for the U and V
1343 plane will be kept separate, i.e. I will have two sets of coefficients that will be applied to
1344 either the U and V planes of wires. I do so as I found this was the choice giving the
1345 best performance. Even so, as I will discuss, the differences are not very pronounced. In
1346 case it is not technically possible to separate channels in the firmware according to the
1347 wire plane they come from and use different sets of filter coefficients for them, we can
1348 just find a common unique set of coefficients. In such case, I do not expect our results
1349 to change dramatically.

1350 In Fig. 6.8 I present the results of our parameter scan, for channels in the induction
1351 planes U (left panel) and V (right panel). For each configuration of σ and δ the resulting
1352 matched filter was applied to all channels in the corresponding plane within the data
1353 capture `felix-2020-07-17-21:31:44`, the S/N improvement was computed with respect
1354 to the raw waveforms and then the S/N mean value was kept as a score for such filter.

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1355 One can see that the improvement obtained for the U plane is in general higher than the
1356 one for the V plane. In any case, I got substantially higher ratios than the ones obtained
1357 for the low-pass FIR filters. For the optimal configurations I attained improvements up
1358 to a factor of 1.85 for the U plane and 1.65 for the V plane.

1359 The sets of optimal matched filter coefficients were obtained for the parameters
1360 $\delta = 0.035$, $\sigma = 0.191$ for the U plane and $\delta = 0.018$, $\sigma = 0.191$ for the V plane. I
1361 show these two sets of coefficients in Fig. 6.9 (left panel). Also in Fig. 6.9 (right
1362 panel) I plot the distribution of the S/N improvement after the optimal match filters
1363 for the U and V were applied to the corresponding channels in the raw data capture
1364 `felix-2020-07-17-21:31:44`. As mentioned before, the mean improvement achieved
1365 for the U plane channels is slightly bigger than the one for the V channels. Note, however,
1366 that the spread of the distribution for the V plane is also smaller than the one for the U
1367 plane.

1368 I also performed a similar scan for the case of a low-pass FIR filter using the Parks-
1369 McClellan algorithm. In that case, the parameters to check were the cutoff frequency
1370 and the transition width of the filter. A summary of the results is given in App. 6.3.

1371 Overall, one can see that the improvements on the S/N are much more significant in
1372 the case of the matched filter than it is for the low-pass FIR filters. The analysis of this
1373 and other raw data captures from ProtoDUNE-SP suggest that matched filters increase
1374 the S/N of induction channels by a factor of 1.5 more than the optimal low-pass FIR
1375 filters.

1376 Although these results are by themselves great points in favour of the matched
1377 filter, more studies are needed to completely assess the robustness of this approach. I
1378 proceeded then to test the matched filter with simulated data samples.

1379 6.5 Using simulated samples

1380 In order to further test the matched filter, the next step was to generate and process
1381 data samples using *LArSoft* [58]. In this way, one can control the particle content of

6.5. Using simulated samples




Figure 6.9: Left panel: Optimal matched filter coefficients for the U (blue line) and V (red line) planes. The filters were computed with our parametrisation in Eq. (6.17) for the parameter values $\delta = 0.035$, $\sigma = 0.191$ and $\delta = 0.018$, $\sigma = 0.191$ respectively. Right panel: Distribution of the relative change of the S/N on the two induction wire planes from the ProtoDUNE-SP raw data capture *felix-2020-07-17-21:31:44* after their respective optimal matched filters were applied.

1382 the samples, the orientation of the tracks and their energy, and therefore see how the
1383 matched filter behaves in various situations.

1384 To begin with, I prepared different monoenergetic and isotropic samples containing
1385 a single particle per event. Each sample contains a different particle species, namely
1386 electrons, muons, protons and neutral pions all with a kinetic energy of $E_k = 100$ MeV.
1387 I chose these because of the fairly different topologies they generate in the liquid argon,
1388 ranging from shower-like to track-like. The procedure I followed to generate the samples
1389 and process them is discussed in detail in App. ??.

1390 These were generated with the single particle gun and the Geant4 stage of the
1391 *LArSoft* simulation [58] was performed with the standard configuration for the DUNE
1392 FD 10kt module.

1393 For simplicity, I restricted the particles to start drifting in a single TPC volume
1394 (in this case TPC 0), so I can focus exclusively on the signals coming from one APA.
1395 The chosen kinetic energy for all the particles in my first trial is $E_k = 100$ MeV, so a
1396 necessary check is to see if all our tracks will be typically contained in one TPC volume.
1397 Fig. 6.10 (left panel) shows the distributions of the track lengths in the liquid argon

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.10: Left panel: distributions of the particles track length in the liquid argon for the generated $E_k = 100$ MeV monoenergetic samples, electrons (blue), muons (red), protons (green) and neutral pions (purple). Right panel: distribution of the length of the longest photon in the neutral pion sample after the decay process $\pi^0 \rightarrow \gamma\gamma$.

of all generated particles with $E_k = 100$ MeV. One can see that, in the case of the track-like particles (i.e. muons and protons), their length distributions are quite sharp and centered at relatively low distances (30 and 8 cm, respectively). For electrons, the distribution is quite broad but it does not extend past ~ 30 cm. The case of neutral pions can be misleading, as they decay promptly the track length associated with the true Monte Carlo particle is always < 1 cm. In Fig. 6.10 (right panel) I show the effective length distribution of the longest photon after the pion decays as $\pi^0 \rightarrow \gamma\gamma$, highlighting the number of fully contained photons in the TPC volume per event (either zero, one or both). One can see that the vast majority of events has both photons contained and that just a negligible number of them has none of them contained in the TPC volume. In any case, for the sake of caution, I will only keep the pion events with both photons contained.

Once I have prepared a sample at the Geant4 level, I need to process it through the detector simulation. In order to make adequate estimations of the noise levels and run the filtering and hit finder as I did with the ProtoDUNE data, one needs to turn off the default zero-suppression of the waveforms produced by the simulation. At this first stage I am only concerned with the waveforms with the noise added, so I keep the noise

6.5. Using simulated samples

addition option as true in the configuration. However, for studies related to the hit finder performance one will also need to store the noiseless waveforms in order to retrieve the truth information of the hits. I will discuss this approach next.

After the detector simulation stage, one needs to extract the no zero-suppressed noisy waveforms, along with their offline channel numbers, and store them in a certain format to be analysed later. To reduce the amount of data that will go for processing, I used the information from the Geant4 step of the simulation to select only the active channels, i.e. the channels where some ionisation electrons arrive. Moreover, as said previously, I only extract the waveforms from APA 0 and exclusively the ones coming from induction channels. The resulting ROOT file contains a tree with two branches, one containing the waveforms for each event and channel and the other with the corresponding offline channel numbers.

Finally, to extract the truth values for the orientation of the tracks and the energies of the particles I used a modified analysis module. This gives a ROOT file with a single tree, containing several branches with different information such as the components of the initial momentum of the particles, initial and final xyz location, track length, etc.

For the analysis of the resulting waveforms and truth values I used a custom set of Python libraries (available at [??]). Among other functionalities, these enable the user to read the ROOT files, export the raw data as pandas objects, apply the filters and compute the S/N of both the raw and filtered signals. So far, the default configuration for the filtering uses the set of optimal matched filter coefficients that I found using the ProtoDUNE data samples.

Additionally, for the analysis of the samples it was necessary to use two different reference frames, to study separately the signals coming from the U and V induction wire planes. As I am focussing on a single APA, the U and V wires have a different orientation in the yz plane. In the case of U wires, these are tilted 35.7° clockwise from the vertical (y direction), whereas the V wires are at the same angle but in the counter clockwise direction. Because of this, the best option is to deal with two new coordinate systems rotated by $\pm 35.7^\circ$ along the x axis, so the new y' and y'' directions are aligned with the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.11: Left panel: schematic representation of the two new rotated reference frames used in this analysis (denoted as prime and double prime), viewed from the yz plane. The magenta stack of lines represent the wires in the U plane, whereas the green lines correspond to the wires in the V plane. Right panel: 3D representation of the momentum of one of the generated monoenergetic muons (red arrow) in the original reference frame (black lines), along with the new reference frame used for the U plane waveforms (blue lines). In the yz plane I added the projection of these three.

1444 U and V induction wires. Fig. 6.11 (left panel) shows a schematic representation of
 1445 the original reference frame together with the two rotated ones (denoted by primed and
 1446 double primed). This way, one can easily understand how parallel was a track to the
 1447 wires in the two induction planes. Fig. 6.11 (right panel) shows a 3D representation of
 1448 the momentum of a track (red arrow) in the original reference frame (black lines), along
 1449 with the new reference frame for U wires (blue lines). I added the projection in the yz
 1450 plane of this three, to show the usefulness of the new reference frame to tell whether a
 1451 track is parallel or normal to the wires in the induction plane.

1452 Fig. 6.12 shows the distribution of the average S/N improvement per event when one
 1453 applies the optimal matched filters. I produced separate distributions for the channels
 1454 in the U (red) and V (blue) induction wire planes. Notice that the S/N distributions
 1455 for the track-like particles, i.e. muons (top left panel) and protons (bottom left panel),
 1456 have significantly larger mean values than the distributions of the shower like particles,
 1457 i.e. electrons (top right panel) and neutral pions (bottom right panel). An important

6.5. Using simulated samples




Figure 6.12: Distributions of the mean S/N improvement per event for the corresponding sample after applying the matched filters. Here I separated the change in the U plane (blue) and the V plane (red) channels. From top left to the right: muon, electron, proton and neutral pion. All the events have a fixed kinetic energy of $E_k = 100$ MeV.

difference between these results and the ones seen before for ProtoDUNE data is that, overall, the improvements that I get for simulated data are bigger. This could be due either to the default noise model used in the *LArSoft* simulation or to the simulated hits having higher energy than the ones in the recorded data. Nonetheless, the concluding message is that the previously optimised matched filters give an overall significant improvement of the S/N for the different samples.

About the convention I followed for the plots and results, in the case of the raw and filtered S/N of each event in the sample I simply took the average of the quantities over all the active channels in the event. That is, if a certain event has N_{chan} active channels these two quantities are computed as:

$$(S/N_{fir})_{event} = \frac{\sum_{i=0}^{N_{chan}} (S/N_{fir})_i}{N_{chan}},$$

$$(S/N_{raw})_{event} = \frac{\sum_{i=0}^{N_{chan}} (S/N_{raw})_i}{N_{chan}}.$$
(6.18)

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.13: Angular dependence of the mean S/N and the S/N improvement, for the different monoenergetic samples considered (from top to bottom: electrons, muons, protons and neutral pions). The two columns on the left represent the values for the U plane waveforms. The top subplots show the mean S/N for raw (green) and filtered (red) waveforms whereas the bottom subplots depict the averaged S/N improvement (black).

1468 However, for the ratio of the raw and filtered S/N (what I called the S/N improvement)
 1469 per event I am not just taking the ratio of the previous two quantities but computing
 1470 the average of the individual ratios per channel in the event:

$$\left(\frac{S/N_{fir}}{S/N_{raw}} \right)_{event} = \frac{\sum_{i=0}^{N_{chan}} \left(\frac{S/N_{fir}}{S/N_{raw}} \right)_i}{N_{chan}}, \quad (6.19)$$

1471 and so:

$$\left(\frac{S/N_{fir}}{S/N_{raw}} \right)_{event} \neq \frac{(S/N_{fir})_{event}}{(S/N_{raw})_{event}}. \quad (6.20)$$

1472 6.5.1 Angular dependence

1473 Having these monoenergetic samples, one can also study the angular dependence of the
 1474 performance of the matched filter. This is an important point, as it is a well established
 1475 fact that for certain configurations (an extreme case configuration being signals normal

6.5. Using simulated samples




Figure 6.14: Angular dependence of the mean S/N and the S/N improvement, for the different monoenergetic samples considered (from top to bottom: electrons, muons, protons and neutral pions). The two columns on the left represent the values for the U plane waveforms. The top subplots show the mean S/N for raw (green) and filtered (red) waveforms whereas the bottom subplots depict the averaged S/N improvement (black).

to the wire plane and perpendicular to the induction wires at the same time) the S/N is much lower than average as the corresponding waveforms are severely distorted. In this sense, I am interested to see how the matched filter behaves for these cases and how the S/N improvement on those compare to the average.

Fig. 6.13 shows the angular dependence of the S/N for the monoenergetic $E_k = 100$ MeV isotropic muons, for the different induction wire planes and projections. The angles for each event are given by the components of the initial value of the momentum of the particles, taking the angles of the projections on the xz and yz planes with respect to the z axis (more accurately, one needs to compute these angles twice for each event, a pair for the $xy'z'$ coordinate system and the other for the $xy''z''$). The top row shows the dependence on the angles corresponding to the U plane, i.e. $\theta_{xz'}$ and $\theta_{y'z'}$, whereas the bottom row shows the angular dependence viewed from the V plane, $\theta_{xz''}$ and $\theta_{y''z''}$. In each plot, the top subplot represents the mean values of the S/N for the raw (blue) and

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1489 matched filtered (red) signals, and the bottom subplot the averaged S/N improvement
1490 (black). The solid lines represent the mean value obtained for the corresponding angular
1491 value, whereas the semitransparent bands represent one standard deviation around the
1492 mean at each point.

1493 As expected, the S/N is in general higher when tracks are parallel to the APA (i.e.
1494 $\theta_{xz} \sim 0$) and lower when it is normal to the plane ($\theta_{xz} \sim \pm 90^\circ$). In the same way, tracks
1495 parallel to the wires ($\theta_{yz} \sim \pm 90^\circ$) tend to have higher S/N than those perpendicular to
1496 these ($\theta_{yz} \sim \pm 0$).

1497 Fig. 6.14 shows the corresponding angular dependence information for the $E_k =$
1498 100 MeV electrons sample. Notice that, in this case, the S/N behaviour discussed above
1499 does not hold. A possible explanation can be that, because most hits in these events
1500 are produced by the secondary particles generated in the EM shower, the signal peaks
1501 whose S/N ratios were computed do not correspond to the directional information of
1502 the primary electron.

1503 6.5.2 Distortion and peak asymmetry

1504 As a little case of study, I selected two of the simulated $E_k = 100$ MeV monoenergetic
1505 muon events. With respect to the U induction plane, one is parallel to the APA (low
1506 $\theta_{xz'}$) and to the wires (high $\theta_{y'z'}$) and the other is normal to the APA plane (high $\theta_{xz'}$)
1507 and perpendicular to the wires (low $\theta_{y'z'}$). As expected from the results on the angular
1508 dependence discussed above, the former has a higher S/N (before and after the filtering)
1509 when compared to the latter. An interesting thing to notice about these two samples
1510 is that, even though one has a much bigger S/N than the other, it is the one with the
1511 smallest S/N the one that got the biggest averaged S/N improvement. In Table 6.1
1512 I included all the relevant parameters of these two $E_k = 100$ MeV muon events I am
1513 considering, namely, the angles with respect to the $xy'z'$ reference frame, the values of
1514 the S/N, the S/N improvement and also the so-called peak asymmetry Δ_{peak} that I will
1515 discuss next.

1516 One can try to understand better what is going on with these two events by looking

6.5. Using simulated samples




Figure 6.15: Selected consecutive waveforms corresponding to two monoenergetic $E_k = 100$ MeV muon events, one is parallel to the APA and to the wires in the U plane (left panel) and the other is normal to the APA plane and perpendicular to the U plane wires (right panel). The solid lines represent the raw waveforms whereas the dashed lines correspond to the waveforms after the matched filter was applied. The waveforms on the left panel have been scaled by a factor of 0.15 to have similar amplitude to the ones on the right panel.

Table 6.1: Characteristic parameters of the two monoenergetic muon events selected, relative to the U plane: projected angles in the xz' and $y'z'$ planes, S/N values for the raw and filtered waveforms, mean improvement of the S/N and peak asymmetry.

	$\theta_{xz'} (\circ)$	$\theta_{y'z'} (\circ)$	S/N _{raw}	S/N _{MF}	$\frac{S/N_{MF}}{S/N_{raw}}$	Δ_{peak} (ADC)
High ("parallel")	-1.58	-77.76	41.65	112.44	2.83	-35.73
Low ("normal")	76.92	-2.56	8.07	25.46	3.12	-10.38

at the raw and filtered data from some of their active channels. Fig. 6.15 shows a selection of consecutive raw and filtered U plane waveforms from the event with high S/N (left panel) and the one with low S/N (right panel). Notice that to show both collections of waveforms at a similar scale I had to apply a factor of 0.15 to the waveforms with high S/N. Additionally, next to each waveform I included the values of the raw and

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.16: Left panel: peak asymmetry distribution for the case of the monoenergetic $E_k = 100$ MeV muon sample. Each value corresponds to a single bipolar signal peak from a channel in any event. The blue distribution represents the peaks on U plane channels, whereas the red corresponds to signal peaks in V wires. Right panel: relation between the mean peak asymmetry per event with the S/N for U channel waveforms from the $E_k = 100$ MeV muon sample. The top subplot shows the decimal logarithm of the mean S/N for the raw (red) and the matched filtered (blue) waveforms. The bottom subplot contains the mean S/N improvement ratio after the matched filter was applied.

1522 matched filtered S/N for the corresponding channel. The first thing to notice in this plot
 1523 is that the amplitude of the signal peaks from the normal track have a much smaller
 1524 amplitude, and also appear quite distorted when compared to the others. On the other
 1525 hand, although the matched filtered S/N is still smaller, the relative improvement is
 1526 bigger than in the parallel case.

1527 A way I found to quantify the difference between the shapes within these two events
 1528 is their different peak asymmetry. One can define the peak asymmetry as the (signed)
 1529 difference between the positive and the negative peaks of the bipolar shape, i.e.:

$$\Delta_{peak} \equiv h_+ - h_-, \quad (6.21)$$

1530 where both heights h_+ and h_- are positive defined. Fig. 6.16 (left panel) shows the
 1531 distribution of this peak asymmetry for all the waveforms corresponding to channels
 1532 in the U (blue) and V (red) planes for the monoenergetic muon sample. One can
 1533 see that these distributions are clearly shifted to negative values (with mean values

6.5. Using simulated samples

1534 $\mu_{\Delta}^U = -10.57$ ADC and $\mu_{\Delta}^V = -5.72$ ADC respectively). It is interesting to notice
1535 that the peak asymmetry value of the sample with high S/N sits at the left tail of the
1536 distribution whereas the corresponding value of the sample with low S/N lies around
1537 the mean.

1538 Now, one can try to correlate the peak asymmetry with the S/N and the S/N change
1539 per event. Fig. 6.16 (right panel) shows the result of comparing (minus) the mean
1540 peak asymmetry per event to the averaged raw (red) and matched filtered (blue) S/N
1541 per event (top subplot). The horizontal lines sit at the mean value obtained in the fit
1542 and represent the width of the $-\Delta_{peak}$ bins used, while the vertical lines indicate one
1543 standard deviation around that mean value. Notice that, when taking decimal logarithm
1544 on both, there is an approximate linear relation between these quantities, except for
1545 peak asymmetry values bigger than -5 ADC where the S/N remains constant.

1546 Also, in the bottom subplot of Fig. 6.16 (right panel) I show the relation between
1547 the peak asymmetry and the mean S/N improvement. In this case, one see that there is
1548 a maximum at $\Delta_{peak} \sim -10$ ADC. As mentioned previously, this is also the value of the
1549 mean of the peak asymmetry distribution. In fact, it is expected that our filter favours
1550 the signal peaks with the most common values of the peak asymmetry, as this was one
1551 of the features I target in our filter coefficient optimisation through the parameter δ .

1552 These results suggest that events with poorer values of the mean S/N, usually
1553 associated to non-favourable track orientations, tend to have smaller values of the mean
1554 peak asymmetry (in absolute value). Nonetheless, because our matched filters have
1555 been optimised to account for these asymmetries, the improvement on the S/N for these
1556 events is sizeable if not better than the one for events which already had a high S/N.

1557 6.5.3 Hit sensitivity

1558 One of the advantages of the matched filter, directly related to increasing the S/N, is
1559 the capability of picking hits that before fell below the threshold. For instance, Fig. 6.17
1560 shows the raw ADC data from an example event (electron, $E_k = 100$ MeV) with the
1561 produced true hits superimposed (black boxes), together with the hits produced by the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.17: Raw data display in the plane time (in firmware ticks) vs. offline channel number for an $E_k = 100$ MeV electron event. The produced true hits are superimposed (black boxes) as well as the hits coming from the standard hit finder chain (blue circles) and the hit finder using the matched filter (green triangles).

standard hit finder chain (blue circles), i.e. using the current FIR filter, and the hits obtained using the matched filters (green triangles). Both the standard and the matched filter hit finders were run with a threshold of 10 ADC. Notice that the standard hits match well the true ones at the initial part of the event (where we have a track-like object), but they miss most of the hits produced by the EM shower at later times. On the other hand, the hits produced with the matched filter have a better agreement with the true hits even for the more diffuse shower activity.

Notwithstanding that now I get more hits with this combination of matched filter and low threshold as a results of the enhancement of the signal peaks relative to the noise level, it is also true that I pick some spurious hits not related to any real activity if one lowers the thresholds too much. Therefore, some optimisation of the threshold is needed. Basically one will need to make a trade-off between precision and sensitivity.

Having this in mind, I tried to compare the produced hits one gets from the standard hit finder and the ones resulting from applying the matched filter with the true hits.

6.5. Using simulated samples

1576 By running the hit finders on our samples with different values of the threshold one
1577 can understand, for instance, how low one can set the threshold without getting mostly
1578 spurious hits and then evaluate the gains obtained from this.

1579 Because now I am also interested in seeing how the hit sensitivity changes with the
1580 energy, I prepared new isotropic samples with the same types of particles as before
1581 (muons, electrons, protons and neutral pions) but with a flat kinetic energy distribution
1582 ranging from 5 to 100 MeV.

1583 In order to estimate the hit sensitivity, given a certain sample, one needs to recover
1584 the set of true hits to be able to compare these with the ones produced. To do so,
1585 a modification in the procedure I was using to extract the raw waveforms is needed.
1586 For this kind of study I run the detector simulation in two steps, first I produce the
1587 waveforms without noise and extract them in the same format I used for the raw data,
1588 then the noise is added and the noisy waveforms are then written to a file as well.

1589 To have a better comparison between the true hits and the ones produced from
1590 the raw waveforms after applying the two filters, I applied also the FIR filter and the
1591 matched filters to the noiseless waveforms and then I run the hit finder with a minimal
1592 threshold (in this case I used 1 ADC) on these noiseless filtered waveforms. In this way
1593 I generated two sets of true hits, I will refer to them as standard true hits (with the
1594 current/default FIR filter) and matched filter true hits respectively. This allows a more
1595 precise matching between the different groups of hits produced, as it will account for
1596 any delays and distortions introduced by the FIR and the matched filters.

1597 In the case of the raw waveforms (with noise), I run the hit finder on them, with
1598 different values of the threshold, after applying either the FIR or the matched filters. I
1599 will name them simply standard hits and matched filter hits respectively. Then, I match
1600 the generated hits to the true hits (the standard hits with the standard true hits and
1601 the matched filter hits with the matched filter true hits). The matching is performed by
1602 comparing the channel number and the timestamp of the hits. To count as a match,
1603 I require that all hits with the same channel number and timestamp have overlapping
1604 hit windows, i.e. the time windows between their hit end and hit start times need to

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.18: Dependence of the precision (blue), sensitivity (red) and F_1 (green) scores on the threshold values used in the hit finder, for the FIR (left panel) and matched filter (right panel) cases. The results were obtained after matching the hits to the true hits in the case of the isotropic muon sample with kinetic energy in the range 5 to 100 MeV, taking only into account the induction plane channels. The points represent the mean value while the error bars indicate one standard deviation around that mean value.

overlap. If more than one hit in one of the groups have hit overlap with the same hit in the other group I only count the hit with closer hit peak time value.

The generation of the samples, the procedure to produce the standard hits (with the default FIR filter) and matched filter hits and the matching of these with the true hits is described in detail in App. ??.

To quantify the performance of the two hit finder approaches, I use a classical method from statistical classification known as confusion matrix [59]. This is basically a way of sorting the outputs of a binary classifier, considering the true values of the classification and the predicted values. It divides the outputs in four categories: true positive (TP, both true and predicted values are 1), false negative (FN, true value is 1 but predicted is 0), false positive (FP, true value is 0 but predicted is 1) and true negative (TN, both true and predicted values are 0)).

The contents of the confusion matrix allow us to compute other derived scores to judge the performance of our classifiers. In this study, I will make use of three of these

6.5. Using simulated samples

1619 metrics, namely the precision or positive predictive value:

$$\text{PPV} = \frac{\text{TP}}{\text{TP} + \text{FP}}, \quad (6.22)$$

1620 the sensitivity or true positive rate:

$$\text{TPR} = \frac{\text{TP}}{\text{TP} + \text{FN}}, \quad (6.23)$$

1621 and the F_1 score [60]:

$$F_1 = \frac{2\text{TP}}{2\text{TP} + \text{FP} + \text{FN}}, \quad (6.24)$$

1622 which is the harmonic mean of the precision and the sensitivity.

1623 In our specific case I am not going to make use of the true negative value, as its
1624 definition in this context can be ambiguous because one does not have clear instances in
1625 the classification process. This way, I will only count the number of true positives as the
1626 total amount of hits I can match between true and raw populations, the number of false
1627 negatives will be the number of missing true hits and the false positive the number of
1628 hits which do not match any true hit.

1629 In Fig. 6.18 I show the precision (blue), sensitivity (red) and F_1 (green) scores I
1630 obtained for different values of the threshold used in the hit finder for the case of the
1631 muon sample. Because the matched filters are only applied to induction channels, I only
1632 consider here hits coming from the U and V planes. The panel on the left corresponds
1633 to the scores I got when I ran the hit finder on the FIR filtered waveforms, whereas the
1634 right panel contains the scores for the matched filter case. The points are centered at
1635 the threshold value used and represent the mean value obtained for each score using all
1636 the generated events, while the error bars indicate one standard deviation around the
1637 mean value.

1638 One can see that the precision for the matched filter case is lower when the thresholds
1639 are very low, as the noise baseline is slightly amplified, but then rises to high values
1640 quicker than for the FIR case. The other difference one can spot is that the sensitivity

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.19: Dependence of the averaged hit sensitivity on the kinetic energy of the events for the matched filter (blue) and standard (red) hits, for the case of the muon (left panel) and electron (right panel) samples, separated between U (top plots) and V (bottom plots) induction wire planes. The top subplots contain the hit sensitivities for the two hit finder alternatives, while the bottom subplots show the ratio between the two. The horizontal lines sit at the mean value and represent the size of the energy bins, while the vertical error bars indicate one standard deviation around that mean value.

in the FIR case starts dropping faster at around the same threshold values where the precision stabilizes around 1, while in contrast for the matched filter this rapid decrease starts at higher threshold values. A similar scan for the same thresholds was performed for the electron sample in the same energy range, yielding similar results.

In Fig. 6.19 I show the averaged hit sensitivity versus the kinetic energy of the events, both for the matched filter hits (blue) and the standard hits (red). The left panel corresponds to the muon sample, whereas the one on the right corresponds to the electron sample, both with kinetic energies between 5 and 100 MeV. In each panel the top plot corresponds to hits in the U plane, while the bottom plot contains the same information for the V plane. Each plot contains two subplots, the one on the top shows the hit sensitivity values for the matched filter and standard hits separate, while the bottom subplot depicts the ratio between the matched filter and standard sensitivities.

6.5. Using simulated samples




Figure 6.20: Distributions of the hit sensitivity in the U (top panels) and V (bottom panels) planes versus the hit sensitivity in the X plane, both for the standard hits (left panels) and the matched filter hits (right panels), in the case of the electron sample and a threshold of 30 ADC.

1653 The horizontal lines are placed at the mean value obtained in the fit and represent the
 1654 width of the E_k bins used, while the vertical error bars indicate one standard deviation
 1655 around that mean value. In both cases the threshold used was 30 ADC, as I required
 1656 the precision to be higher than 0.99 for both matched filter and standard cases.

1657 One can see that, in general, the improvements are better for the U than for the V
 1658 plane. While for the U channels I achieved a mean improvement of 50% and 80% for
 1659 muons and electrons respectively, the improvement in the V plane is stalled at 10% and
 1660 25%. Nevertheless, if I look at the sensitivities for the matched filter hits in both planes
 1661 one can see these have similar mean values for each energy bin, while on the contrary

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.21: Top panels: standard residual plots of the hit sensitivities between the X and U planes. Bottom panels: quantile-quantile plots of the hit sensitivity standard residuals between the X and U planes. In all cases, the left panel corresponds to the standard hits while the right panel represents the matched filter case, all from the electron sample with a 30 ADC threshold.

for the standard hits the sensitivity remains relatively high for the V plane. This way, it looks there was a less significant gain because the hit sensitivity was already high.

Another interesting observation is the different behaviors for muons and electrons. While hit sensitivity for muons grows significantly with energy, in the case of electrons this slightly decreases the higher the kinetic energy of the event is. In any case, when it comes to the improvement on the sensitivities, this remains almost constant in all cases.

Furthermore, we can look at how the concurrence of hits between the different wire planes has changed. For any given event, I expect to have a similar number of hits in the three planes. As the ionisation electrons need to cross the U and V planes prior to reach the collection plane X they will induce current in those wire planes. A way to check the concurrence of hits across planes is looking at the relation between the hit sensitivities for each individual event. One cannot expect the sensitivities to be exactly equal across

6.5. Using simulated samples

1674 planes, but ideally they should be normally distributed around the diagonal.

1675 Fig. 6.20 shows the hit sensitivity in the U (top panels) and V (bottom panels)
1676 planes versus the hit sensitivity in the X plane, for the case of the standard hits (left
1677 panels) and the matched filter hits (right panels). All plots were generated for the
1678 electron sample and a threshold of 30 ADC. From these one can see a clear trend,
1679 when I use the standard hit finder chain the sensitivities in the induction planes are
1680 systematically lower than the hit sensitivity in the X plane, i.e. most of the points sit
1681 below the diagonal (red dashed line). In contrast, when the matched filters are applied,
1682 the majority of the events are distributed around the diagonal. This points out that the
1683 concurrence of hits across planes has improved.

1684 To exemplify the improvement I obtained, one can consider the residuals of the hit
1685 sensitivities for the X and U planes. Assuming the diagonal hypothesis, i.e. given a
1686 dataset of the form (x, y) for any x I take the predicted y value to be equal to the value
1687 of x , I can compute the standard residuals for the hit sensitivities in U given the ones for
1688 X. In Fig. 6.21 (top panels) I show these standard residuals against the corresponding
1689 values of the hit sensitivity in the U plane, for our electron sample with kinetic energy
1690 between 5 and 100 MeV. If I compare the scatter points in the case of the standard
1691 hits (left panel) and the matched filter hits (right panel), I see that the residuals of the
1692 standard hit finder case follow a certain pattern and their mean deviates from 0.

1693 To see clearly if the residuals are normally distributed, in Fig. 6.21 (bottom panels)
1694 I plot the corresponding quantile-quantile plot for both the standard (left panel) and
1695 matched filter (right panel) standard residuals. One can clearly see that the points for
1696 the standard case follow a strongly non-linear pattern, suggesting that the residuals
1697 do not follow a normal distribution. In contrast, for the matched filter hits the points
1698 conform to a roughly linear path, implying that in this case the normality condition is
1699 fulfilled.

1700 All these results hint at the fact that the concurrence of hits across the wire planes
1701 can be strengthened by applying the matched filters.

1702 **Chapter 7**

1703 **DM searches with neutrinos from
1704 the Sun**

1705 **7.1 Motivation**

1706 The idea of detecting neutrino signals coming from the Sun’s core to probe DM is not new.
1707 The main focus of these searches has usually been high-energy neutrinos originated from
1708 DM annihilations into heavy particles [61–64], although recent studies have proposed to
1709 look at the low-energy neutrino flux arising from the decay of light mesons at rest in the
1710 Sun [65–68] previously thought undetectable.

1711 In this chapter I try to demonstrate the capability of DUNE to constrain different
1712 DM scenarios. I used the neutrino fluxes arising from DM annihilations in the core
1713 of the Sun to compute the projected limits that DUNE would be able to set on the
1714 annihilation rates in the Sun and the DM scattering cross sections.

1715 **7.2 Gravitational capture of DM by the Sun**

1716 The Sun and the centre of the Earth are possible sources of DM annihilations, specially
1717 interesting because of their proximity. Their gravitational attraction ensured the capture
1718 of DM from the local halo through repeated scatterings of DM particles crossing them.

Chapter 7. DM searches with neutrinos from the Sun

1719 Only neutrinos produced from DM annihilations can escape the dense interior of these
1720 objects. Therefore, neutrino telescopes are the most useful experimental layouts to
1721 pursue DM searches from their cores.

1722 The neutrino flux from DM annihilations inside the Sun depends on the DM capture
1723 rate, which is proportional to the DM scattering cross section, and the annihilation rate,
1724 which is proportional to the velocity-averaged DM annihilation cross-section. The total
1725 number of DM particles inside the Sun follows the Boltzmann equation [65]:

$$\frac{dN_{DM}}{dt} = C_{\odot} - A_{\odot} N_{DM}^2, \quad (7.1)$$

1726 where C_{\odot} and A_{\odot} are the total Sun DM capture and annihilation rates respectively.
1727 In this expression I neglected the evaporation term, proportional to N_{DM} , which only
1728 contribute for $m_{DM} \lesssim 4$ GeV [69]. As the current threshold of neutrino telescopes is
1729 a few GeV, this region falls below the probed range but can be important in future
1730 low-energy projects.

1731 This equation has an equilibrium solution:

$$N_{DM}^{eq} = \sqrt{\frac{C_{\odot}}{A_{\odot}}}, \quad (7.2)$$

1732 which represents the amount of DM inside the Sun if the capture and annihilation have
1733 reached equilibrium. As the Sun is approximately 4.6 Gyr old, it is usually assumed that
1734 equilibrium has been achieved. Therefore, the anomalous neutrino flux from the Sun
1735 would only depend on the DM scattering cross section, enabling us to set limits on this
1736 quantity. If one does not assume equilibrium, some assumptions on the DM annihilation
1737 cross section are necessary to extract predictions from neutrino signals.

1738 Here, I am going to consider three possible scenarios for the DM interactions: DM
1739 scattering off electrons, spin-dependent (SD) and spin-independent interactions off nuclei.
1740 For the case of these last two, the cross sections will be given in terms of the SD and
1741 SI elastic scattering DM cross section off protons (assuming that DM interactions off

7.2. Gravitational capture of DM by the Sun

1742 protons and neutrons are identical), σ_p^{SD} and σ_p^{SI} , as [4, 65]:

$$\sigma_i^{\text{SD}} = \left(\frac{\tilde{\mu}_{A_i}}{\tilde{\mu}_p} \right)^2 \frac{4(J_i + 1)}{3J_i} |\langle S_{p,i} \rangle + \langle S_{n,i} \rangle|^2 \sigma_p^{\text{SD}}, \quad (7.3)$$

$$\sigma_i^{\text{SI}} = \left(\frac{\tilde{\mu}_{A_i}}{\tilde{\mu}_p} \right)^2 A_i^2 \sigma_p^{\text{SI}}, \quad (7.4)$$

1743 where $\tilde{\mu}_{A_i}$ is the reduced mass of the DM-nucleus i system, $\tilde{\mu}_p$ is the reduced mass of
1744 the DM-proton system, A_i and J_i the mass number and total angular momentum of
1745 nucleus i and $\langle S_{p,i} \rangle$ and $\langle S_{n,i} \rangle$ the expectation value of the spins of protons and neutrons
1746 averaged over all nucleons, respectively (see Ref. [70] for a review on spin expectation
1747 values).

1748 Since the Sun is mainly composed of Hydrogen, the capture of DM from the halo
1749 is expected to occur mainly through spin-dependent scattering. However, since the
1750 spin-independent cross section is proportional to the square of the atomic mass, heavy
1751 elements can contribute to the capture rate (even though they constitute less than 2%
1752 of the mass of the Sun). Heavy elements can also contribute to the spin-dependent cross
1753 section if the DM has also momentum-dependent interactions.

1754 DM particles can get captured by the Sun if after repeated scatterings off solar
1755 targets their final velocity is lower than the escape velocity of the Sun. In the limit of
1756 weak cross sections, this capture rate can be approximately written as [4]:

$$C_{\odot}^{\text{weak}} = \sum_i \int_0^{R_{\odot}} dr 4\pi r^2 \int_0^{\infty} du_{\chi} \frac{\rho_{\chi}}{m_{\chi}} \frac{f_{v_{\odot}}(u_{\chi})}{u_{\chi}} \omega(r) \int_0^{v_e(r)} dv R_i^-(\omega \rightarrow v) |F_i(q)|^2, \quad (7.5)$$

1757 where the summation extends over all possible nuclear targets. In this expression, R_{\odot}
1758 is the radius of the Sun, ρ_{χ} is the local DM density, m_{χ} the mass of the DM particle,
1759 $f_{v_{\odot}}(u_{\chi})$ the DM velocity distribution seen from the Sun's reference frame, $R_i^-(\omega \rightarrow v)$
1760 is the differential rate at which a DM particle with velocity v scatters a solar target of
1761 mass m_i to end up with a velocity ω and $|F_i(q)|$ is the nuclear form factor of target i .

1762 The differential scattering rate takes a rather simple form when considering velocity-

Chapter 7. DM searches with neutrinos from the Sun

¹⁷⁶³ independent and isotropic cross sections. In that case, this quantity is given by [4, 71]:

$$R_i^-(\omega \rightarrow v) = \frac{2}{\sqrt{\pi}} \frac{\mu_{i,+}^2}{\mu_i} \frac{v}{\omega} n_i(r) \sigma_i \left[\chi(-\alpha_-, \alpha_+) + \chi(-\beta_-, \beta_+) e^{\mu_i(\omega^2 - v^2)/u_i^2(r)} \right], \quad (7.6)$$

¹⁷⁶⁴ where μ_i is the ratio between the DM mass and the mass of target i , $\mu_{i,\pm}$ is defined as:

$$\mu_{i,\pm} \equiv \frac{\mu_i \pm 1}{2}, \quad (7.7)$$

¹⁷⁶⁵ $n_i(r)$ is the density profile of target i in the solar medium, $u_i(r)$ is the most probable

¹⁷⁶⁶ velocity of target i given by:

$$u_i(r) = \sqrt{\frac{2T_\odot(r)}{m_i}}, \quad (7.8)$$

¹⁷⁶⁷ where $T_\odot(r)$ is the temperature of the Sun, the quantities α_\pm and β_\pm are defined as:

$$\alpha_\pm \equiv \frac{\mu_{i,+}v \pm \mu_{i,-}\omega}{u_i(r)}, \quad (7.9)$$

$$\beta_\pm \equiv \frac{\mu_{i,-}v \pm \mu_{i,+}\omega}{u_i(r)}, \quad (7.10)$$

¹⁷⁶⁸ and the function $\chi(a, b)$ is a Gaussian integral of the form:

$$\chi(a, b) \equiv \int_a^b dx e^{-x^2}. \quad (7.11)$$

¹⁷⁶⁹ Finally, if one assumes the DM halo velocity distribution in the galactic rest frame

¹⁷⁷⁰ to be a Maxwell-Boltzmann distribution, one can write the halo velocity distribution for

¹⁷⁷¹ an observer moving at the speed of the Sun with respect to the DM rest frame as:

$$f_{v_\odot}(u_\chi) = \sqrt{\frac{3}{2\pi}} \frac{u_\chi}{v_\odot v_d} \left(e^{-\frac{3(u_\chi - v_\odot)^2}{2v_d^2}} - e^{-\frac{3(u_\chi + v_\odot)^2}{2v_d^2}} \right), \quad (7.12)$$

¹⁷⁷² where:

$$\omega^2 = u_\chi + v_e(r)^2, \quad (7.13)$$

¹⁷⁷³ is the DM velocity squared, v_\odot the relative velocity of the Sun from the DM rest frame

7.2. Gravitational capture of DM by the Sun




Figure 7.1: Input solar parameters used in our capture rate computation as functions of the Sun's radius, from left to right: temperature (with respect to the temperature at the core), mass (in solar masses) and electron number density (with respect to the electron density at the core). All quantities shown correspond to the standard solar model BS2005-OP [3].

1774 and $v_d \simeq \sqrt{3/2}v_\odot$ the velocity dispersion.

1775 For the case of strong scattering cross section, Eq. (7.5) ceases to be valid, as it
 1776 escalates indefinitely with the cross section. In that limit, the capture rate saturates to
 1777 the case where the probability of interaction is equal to one, which can be written as:

$$C_\odot^{\text{geom}} = \pi R_\odot^2 \left(\frac{\rho_\chi}{m_\chi} \right) \langle v \rangle \left(1 + \frac{3}{2} \frac{v_e^2(R_\odot)}{v_d^2} \right) \xi(v_\odot, v_d), \quad (7.14)$$

1778 where $v_d = \sqrt{8/3\pi}v_d$ is the mean velocity in the DM rest frame and the factor $\xi(v_\odot, v_d)$
 1779 accounts for the suppression due to the motion of the Sun:

$$\xi(v_\odot, v_d) = \frac{v_d^2 e^{-\frac{3v_\odot^2}{2v_d^2}} + \sqrt{\frac{\pi}{6}} \frac{v_d}{v_\odot} (v_d^2 + 3v_e^2(R_\odot) + 3v_\odot^2) \text{Erf} \left(\sqrt{\frac{3}{2}} \frac{v_\odot}{v_d} \right)}{2v_d^2 + 3v_e^2(R_\odot)}. \quad (7.15)$$

1780 Having these into account, one can write the total capture rate as a combination of
 1781 both contributions, allowing a smooth transition between the two, as:

$$C_\odot = C_\odot^{\text{weak}} \left(1 - e^{C_\odot^{\text{geom}}/C_\odot^{\text{weak}}} \right). \quad (7.16)$$

1782 I computed the capture rate from Eq. (7.16) in the case of interactions with
 1783 electrons. To do so, I used the standard solar model BS2005-OP [3]. Fig. 7.1 shows the

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.2: Capture rates as a function of the DM mass for the DM-electron interactions (red lines), SD DM-nucleons interactions (green lines) and SI DM-nucleons interactions (blue lines). Solid lines represent the values computed in this work while the dashed lines are the one given in Ref. [4]. All the rates are shown for a choice of scattering cross section of $\sigma_i = 10^{-40} \text{ cm}^2$.

1784 three parameters from the solar model that are needed for the computation, the solar

1785 temperature (left panel), mass (central panel) and electron density (right panel) profiles.

1786 For the case of the interactions off nuclei, the computations are more convoluted

1787 as one needs to add up the contributions of the different most abundant nuclei in

1788 the Sun. Also, in contrast to the electron scenario where the form factor is trivially

1789 $|F_e(q)|^2 = 1$, for any nucleus i one would need to consider some appropriate nuclear

1790 density distribution (either a Gaussian approximation, a Woods-Saxon distribution, etc)

1791 which would complicate the calculations even further.

1792 That is the reason why, at this stage of our study, I decided to take an alternative

1793 approach to the computation of the DM-nucleus capture rates. I used the **DarkSUSY**

1794 software, that allows us to compute these quantities performing a full numerical

1795 integration over the momentum transfer of the form factors. The default standard

1796 solar model used by **DarkSUSY** is BP2000¹ [72].

¹This is what they say in their manual, but I fear it is somewhat outdated. It appears to me this

7.2. Gravitational capture of DM by the Sun

1797 In Fig. 7.2 I show the results I obtained for the capture rates, for the case of
 1798 interactions off electrons (red solid line), SD (green solid line) and SI (blue solid line)
 1799 interactions of nucleons. In all cases I used a value of the scattering cross sections of
 1800 $\sigma_i = 10^{-40} \text{ cm}^2$. Note here one of the limitations of the **DarkSUSY** approach, one can
 1801 not extend the computation below $m_{\text{DM}} = 1 \text{ GeV}$. Nevertheless, this is not something
 1802 to worry about in this case, as I will discuss next. As a comparison, I added also the
 1803 values computed in Ref. [4] (same color scheme, dashed lines). One can see there is good
 1804 agreement between these and the **DarkSUSY** computation of the SD and SI interactions
 1805 for $m_{\text{DM}} \geq 1 \text{ GeV}$. In this regime their computations also matches quite well our
 1806 result for the electron capture rate. However, these start to differ significantly below
 1807 $m_{\text{DM}} = 1 \text{ GeV}$, being their estimate up to a factor of 5 bigger than ours for low masses.
 1808

1808 Let us comment briefly about the assumption I made before about not including
 1809 an evaporation term in the Boltzmann equation. If I include this term in the equation
 1810 (which will be proportional to the number of DM particles) the equilibrium solution
 1811 takes the form:

$$N_{\text{DM}}^{eq} = \sqrt{\frac{C_\odot}{A_\odot}} \frac{1}{\kappa + \frac{1}{2} E_\odot \tau_{eq}}, \quad (7.17)$$

1812 where E_\odot is the total evaporation rate, τ_{eq} is the equilibrium time in the absence of
 1813 evaporation:

$$\tau_{eq} = \frac{1}{\sqrt{C_\odot A_\odot}}, \quad (7.18)$$

1814 and κ is defined as:

$$\kappa \equiv \sqrt{1 + \left(\frac{E_\odot \tau_{eq}}{2} \right)^2}. \quad (7.19)$$

1815 Now, it is easy to proof that in case evaporation dominates $\kappa \gg 1$ and therefore:

$$N_{\text{DM}}^{eq} \simeq \frac{C_\odot}{E_\odot}. \quad (7.20)$$

1816 In contrast, if evaporation is irrelevant $\kappa \simeq 1$ and one recovers Eq. (7.2).

model is relatively old and do not see why they are not using others like [3]. Maybe one can double-check in the code to make sure.

Chapter 7. DM searches with neutrinos from the Sun

1817 In this way, one can define the evaporation mass as the mass for which the number
1818 of DM particles in equilibrium approaches Eq. (7.20) at 10% level:

$$\left| N_{DM}^{eq}(m_{\text{evap}}) - \frac{C_{\odot}(m_{\text{evap}})}{E_{\odot}(m_{\text{evap}})} \right| = 0.1 N_{DM}^{eq}(m_{\text{evap}}). \quad (7.21)$$

1819 This can be regarded as the minimum testable mass one can reach using the annihilation
1820 products of the DM in the Sun.

1821 It was reported in Ref. [4] that, in the case of both SD and SI DM interactions
1822 off nuclei, this value ranges from 2 to 4 GeV depending on the specific scattering
1823 cross section value, compatible with the usual assumptions in the literature. What is
1824 interesting is the case of the electron capture. It was found that, when one applies a
1825 cutoff in the velocity distribution of the DM trapped in the Sun slightly below the escape
1826 velocity, the evaporation mass for the DM-electron interaction decreases remarkably. For
1827 a moderate choice of $v_c(r) = 0.9v_e(r)$ one gets an evaporation mass of around 200 to
1828 600 MeV. This possibility opens a region of the parameter space that could be tested
1829 with neutrino detectors.

1830 7.3 Neutrino flux from DM annihilations

1831 When WIMPs annihilate inside the Sun a flux of high-energy neutrinos is expected from
1832 heavy quarks, gauge bosons and $\tau^+\tau^-$ final states, which decay before losing energy
1833 in the dense solar medium, as they will produce a continuum spectra up to $E_{\nu} \sim m_{\chi}$
1834 (in the case of direct annihilation to neutrinos one would have a line at $E_{\nu} = m_{\chi}$) [66].
1835 This kind of signal has been extensively studied in the literature, allowing to put strong
1836 limits on the SD WIMP-proton cross section for large m_{χ} . However, the number of
1837 high-energy neutrinos per WIMP annihilation is small and the spectrum depends on the
1838 unknown final state. Moreover, background rejection is easier for large m_{χ} but neutrinos
1839 with $E_{\nu} \gtrsim 100$ GeV are significantly attenuated by interactions in the Sun.

1840 Nevertheless, most WIMP annihilation final states eventually produce a low-energy
1841 neutrino spectrum. In this case one does not just consider the more massive final

7.4. Computing limits from solar neutrino fluxes

states but also annihilations into e^+e^- , $\mu^+\mu^-$ and light quarks [65]. In particular, light mesons would be produced and stopped in the dense medium, thus decaying at rest and producing a monoenergetic neutrino signal. The decay-at-rest of kaons will produce a $E_\nu = 236$ MeV ν_μ while in the case of pions one would have a $E_\nu = 29.8$ MeV ν_μ . In practice only K^+ and π^+ contribute to these signals, as K^- and π^- are usually Coulomb-captured in an atomic orbit and get absorbed by the nucleus. There is also a low-energy neutrino signal coming from muon decays, which are produced in kaon or pion decays, leptonic decays of other hadrons and heavy leptons or even directly from WIMP annihilations, which can decay at rest and contribute to the previous low-energy neutrino flux with a well known spectrum below 52.8 MeV.

These monoenergetic MeV neutrinos were previously considered undetectable but, due to the large yield, the known spectra and the modern advances in the detector technology, these low-energy neutrino flux can be a good probe of the SD WIMP-proton cross-section in standard solar WIMP capture scenario, as it is sensitive to low WIMP masses and insensitive to the particular final state. A good place to look for these signals are next-generation neutrino experiments such as DUNE.

7.4 Computing limits from solar neutrino fluxes

In order to use the neutrino fluxes from DM annihilations in the Sun, the first thing I need to do is to determine the expected number of atmospheric background events, for a given exposure, after directionality selection has been applied. I can write this number as:

$$N_B = \eta_B \int d\Omega \int_{E_{min}}^{E_{max}} dE_\nu \frac{d^2\Phi_{atm}^\mu}{dE_\nu d\Omega} \times \left(A_{eff}^{(\mu)}(E_\nu) T \right), \quad (7.22)$$

where η_B is the background efficiency, E_{min} and E_{max} the minimum and maximum energies to integrate over, $d^2\Phi_{atm}^\mu/dE_\nu d\Omega$ the differential flux of atmospheric muon neutrinos, $A_{eff}^{(\mu)}$ is the effective area of DUNE to muon neutrinos and T is the exposure time. The effective area can be expressed as the product of the neutrino-nucleus scattering cross section and the number of nuclei in the fiducial volume of the detector. This way

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.3: *NuWro* computed $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section as a function of the neutrino energy E_μ . The black line shows to the total cross section, whereas the others correspond to the different contributions (in red quasi-elastic scattering, in green resonant pion exchange, in blue deep inelastic scattering and in purple meson exchange current).

1868 for DUNE I can write:

$$A_{eff}^{(\mu)}(E_\nu) = (6.0 \times 10^{-10} \text{ m}^2) \left(\frac{\sigma_{\nu - \text{Ar}}^{(\mu)}(E_\nu)}{10^{-38} \text{ cm}^2} \right) \left(\frac{M_{target}}{40 \text{ kT}} \right), \quad (7.23)$$

1869 where $\sigma_{\nu - \text{Ar}}^{(\mu)}$ is the $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section. In Fig. 7.3 I
 1870 show the computed value of this cross section as a function of the neutrino energy E_ν ,
 1871 in the range of interest both for the atmospheric background and signal events. It was
 1872 computed using the **NuWro** Monte Carlo neutrino event generator [73], including the
 1873 charged-current contributions of the quasi-elastic scattering (red line), resonant pion
 1874 exchange (green line), deep inelastic scattering (blue line) and meson exchange current
 1875 (purple line).

1876 The background rejection will depend on the resolution of the detector and the
 1877 selection one applies on the events. A geometry argument can be used to estimate
 1878 the maximum background rejection one can achieve in this case, considering one can

7.4. Computing limits from solar neutrino fluxes

1879 efficiently discriminate all events coming from a direction different from that of the
 1880 Sun. In that case, the optimal background efficiency will simply be the relative angular
 1881 coverage of the Sun. Taking the angular diameter of the Sun as seen from the Earth to
 1882 be 0.5° , I have:

$$\eta_B^{(opt)} \approx \frac{\pi \left(\frac{0.5}{2}\right)^2}{360 \times 180} \simeq 3.03 \times 10^{-6}. \quad (7.24)$$

1883 This value will give a very optimistic estimate of the number of background events.
 1884 However, it can be regarded as an lower limit, as it represents the best case scenario.

1885 In Fig. 7.4 I show the fluxes of atmospheric neutrinos at the Homestake mine during
 1886 solar minimum, taken from Ref. [5]. The values are averaged over the two angular
 1887 directions. In blue I have the flux of muon neutrinos while in red I indicate the flux
 1888 of electron neutrinos. Additionally, the dashed lines correspond to both antineutrino
 1889 species.

1890 Using these values for the muon neutrino and the corresponding total CC cross
 1891 section, one can compute the number of expected background events by integrating over
 1892 the given energy range (as in this case the angular integral is trivial). As for the energy
 1893 range to integrate over, I choose the range for DUNE specified in [42], $E_{min} = 10^{-1}$ GeV
 1894 and $E_{max} = 10$ GeV. Taking all these into account, I found the number of background
 1895 events to be:

$$N_B \simeq \eta_B \times (3.827 \times 10^4) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right). \quad (7.25)$$

1896 In order to estimate the sensitivity of DUNE to this kind signal, one can consider a
 1897 hypothetical data set where the number of observed neutrinos is taken to be the expected
 1898 number of background events rounded to the nearest integer, $N_{obs} = \text{round}(N_B)$ [74].
 1899 Now, if I assume that the number of signal and background events seen by DUNE are
 1900 given by Poisson distributions with means equal to the expected number of signal and
 1901 background events, N_S and N_B , one can denote by N_S^{90} to the number of expected
 1902 signal events such that the probability of having an experimental run with a number of
 1903 events greater than N_{obs} is 90%. This number can be obtained as the numerical solution

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.4: Expected atmospheric neutrino flux as a function of the neutrino energy E_ν at Homestake at solar minimum, taken from Ref. [5]. The blue solid (dashed) line correspond to muon neutrinos (antineutrinos) and the red solid (dashed) line correspond to electron neutrinos (antineutrinos).

1904 to the equation:

$$1 - \frac{\Gamma(N_{obs} + 1, N_S^{90} + N_B)}{N_{obs}!} = 0.9, \quad (7.26)$$

1905 where $\Gamma(x, y)$ is the upper incomplete gamma function.

1906 The number of signal events is related to the neutrino flux from DM annihilations in
 1907 a similar way as the background events to the atmospheric neutrino flux. In this case I
 1908 have:

$$N_S = \eta_S \Gamma_A^{eq} \int_{z_{min}}^{z_{max}} dz \frac{dN_\nu}{dAdN_A dz} \times (A_{eff}^\mu(z)T), \quad (7.27)$$

1909 where η_S is the signal efficiency, Γ_A^{eq} is the total annihilation rate of DM particles at
 1910 equilibrium, $\Gamma_A^{eq} = A_\odot (N_{DM}^{eq})^2$, z_{min} and z_{max} the minimum and maximum relative
 1911 energies to integrate over (in such a way that $z_{min,max} \leq E_{min,max}/m_{DM}$ for each m_{DM})
 1912 and $dN_\nu/dAdN_A dz$ the muon neutrino flux per DM annihilation in the Sun.

1913 Knowing N_S^{90} one can use the relation in Eq. (7.27) to obtain $\Gamma_A^{eq,90}$ for different
 1914 values of the DM mass. From there I can directly translate those values into the

7.5. Example: Kaluza-Klein Dark Matter

1915 upper limits for DUNE on the DM scattering cross sections, for a given exposure. The
1916 relation between the annihilation rate and the DM-nucleon cross section comes from the
1917 equilibrium condition through the solar DM capture rate. The details of the evolution
1918 of the number of DM particles inside the Sun and the computation of the capture rates
1919 are discussed in App. 7.2.

1920 7.5 Example: Kaluza-Klein Dark Matter

1921 Even though there are plenty of BSM theories which provide viable dark matter
1922 candidates, Kaluza-Klein type of models [75, 76] within the universal extra dimensions
1923 (UED) paradigm naturally predict the existence of a massive, stable particle that can
1924 play the role of the dark matter. In the UED scenario all the SM fields can propagate
1925 in one or more compact extra dimensions [77], as opposed to the idea of brane worlds
1926 [78, 79], where just gravity can propagate in the bulk while SM particles live at fixed
1927 points.

1928 Furthermore, in UED there is no violation of the translational invariance along the
1929 extra dimensions, thus leading to degenerate KK modes masses and also the conservation
1930 of the KK number in the effective four dimensional theory. At loop level, radiative
1931 corrections and boundary terms shift the masses of the KK modes and break KK
1932 number conservation into a KK parity. As a result, this theory only contains interactions
1933 between an even number of odd KK modes and therefore the lightest among the first KK
1934 excitations will be stable. This particle is usually denoted as the lightest Kaluza-Klein
1935 particle (LKP) and its mass is proportional to $1/R$, being R the size of the extra
1936 dimension.

1937 A viable DM candidate needs to be electrically neutral and non-baryonic, therefore
1938 good candidates among the first Kaluza-Klein excitations would be the KK neutral
1939 gauge bosons and the KK neutrinos [80]. Another possible candidate is the first KK
1940 excitation of the graviton, which receives negligible radiate contributions and therefore
1941 has a mass almost equal to $1/R$, but it has been shown that the lightest eigenstate from

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.5: Feynman diagrams for B^1B^1 annihilation into SM fermions.




Figure 7.6: Feynman diagrams for B^1B^1 annihilation into a Higgs boson pair.

7.5. Example: Kaluza-Klein Dark Matter




Figure 7.7: Computed spectra of muon neutrinos at the DUNE FD site from B^1 annihilations in the Sun for three different values of M_{LKP} , plotted in relative energy units for legibility.

1942 the mixing of the gauge mass states (B^1, W_3^1) would be lighter, as B^1 and W_3^1 receive
 1943 negative radiative corrections [81]. It is also understood that, when these corrections
 1944 become sizeable, the eigenstates become approximately pure B^1 and W_3^1 states as the
 1945 Weinberg mixing angle grows small with the KK number [81]. In that case, the LKP
 1946 can be well-approximated as being entirely B^1 .

1947 I need to compute the neutrino flux produced by the annihilations of the LKP in
 1948 the core of the Sun, taking into account their propagation in the solar medium, as well
 1949 as neutrino oscillations. To this end I used `WimpSim` [82, 83] to generate one million
 1950 annihilation events in the Sun over a time span of four years and propagate them to
 1951 the DUNE FD location ($44^\circ 20' \text{ N}, 103^\circ 45' \text{ W}$), for different values of M_{LKP} . In Fig.
 1952 7.7 I show the obtained muon neutrino spectra arriving to the detector from LKP
 1953 annihilations in the Sun, per unit area and per annihilation, plotted in relative energy
 1954 units for different values of the mass. As one could expect the spectra get steeper the
 1955 higher is the mass, due to the absorption of high-energy neutrinos in the solar medium.
 1956 Also, one can see the peak at $z = 1$ due to the direct annihilation into neutrinos $\chi\chi \rightarrow \nu\bar{\nu}$.

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.8: Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin-dependent B^1 -proton scattering cross section as a function of M_{LKP} (green dots). I also show the previous limits from IceCube [6] (blue line) and Antares [7] (red line) on the LKP cross section. The shaded area represents the disfavoured region (at 95% confidence level) on the mass of the LKP from LHC data [8].

Now, one can estimate the sensitivity of DUNE to this particular model by using the methods I previously discussed. To begin with, I will use the optimistic estimation of the background efficiency in Eq. (7.24) to get our upper bound. Using it, one can directly compute the number of expected background events to be $N_B = 0.1101$ for an exposure of 400 kT yr. Then, Eq. (7.26) give us a value of $N_S^{90} = 2.20$ for the 90% exclusion number of expected signal events. By using the NuWro generated cross sections and the computed neutrino fluxes from B^1 annihilations in the Sun I can estimate the limits on the SD and SI DM-nucleus cross section using the relation in Eq. (7.2) and the capture rates I computed with DarkSUSY.

In Fig. 7.8 I show the projected sensitive for DUNE on the spin-dependent B^1 -proton scattering cross section versus the mass of the DM particle, for a exposure of 400 kT yr (green dots). I also include the previous results from IceCube [6] (blue line) and Antares [7] (red line). The shaded area represents the disfavoured region from combined searches for UED by ATLAS and CMS [8].

7.6. High energy DM neutrino signals




Figure 7.9: Computed spectra of muon neutrinos at the DUNE FD site from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilations in the Sun for the DM masses $m_{\text{DM}} = 10 \text{ GeV}$ (red line), 50 GeV (green line) and 100 GeV (blue line), plotted in relative energy units.

From the experimental point of view, this estimation lacked a detailed simulation of the detector response and thus this must be consider as a mere optimistic sensitivity computation. However, it shows the potential of DUNE to constrain this kind of exotic scenarios, showing the region where it will be in a position to compete with other neutrino telescopes. A more detailed analysis is needed if I am to make a realistic estimation. Even though the region of the parameter space where DUNE would be sensitive to this particular model is quite constrained by collider searches [8] and other rare decay measurements [84, 85], it still constitutes an alternative indirect probe.

7.6 High energy DM neutrino signals

To have better estimates on the capability of the DUNE FD to constrain the parameter space of DM using solar neutrino fluxes, I need to start accounting for the detector resolution effects and the topologies of the different signatures. As a starting point, I will focus on specific annihilation channels. For the case of DUNE, the relevant ones are mainly the hard channels $\tau^+\tau^-$ and $\nu\bar{\nu}$ and the soft channel $b\bar{b}$. These are the open annihilation channels for relatively low mass WIMPs that will actually give neutrino fluxes. Other channels, like W^+W^- and ZZ , are open for more massive WIMPs, but

Chapter 7. DM searches with neutrinos from the Sun

1987 those will produce usually a higher energy neutrino flux that will be out of reach for
1988 DUNE (usually the maximum neutrino energy is taken to be $E_{max} = 10$ GeV).

1989 In Fig. 7.9 I show the `WimpSim` [82, 83] generated muon neutrino spectra at the
1990 DUNE FD location ($44^\circ 20' N, 103^\circ 45' W$) from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel)
1991 annihilations in the core of the Sun, for different DM masses. Here, one can clearly see
1992 the meaning of the previous distinction between hard and soft channels. For the same
1993 DM mass value, the muon neutrino spectrum from the $\tau^+\tau^-$ channel is more flat and
1994 reaches higher energies than the one from the $b\bar{b}$ channel, which drops faster.

1995 In this case, I prepared two sets of files, one for $\tau^+\tau^-$ and the other for $b\bar{b}$, for DM
1996 masses in the range from 5 to 100 GeV (actually for $b\bar{b}$ the first mass point I took is
1997 7.5 GeV, as a WIMP with $m_{DM} = 5$ GeV can not kinematically self annihilate into $b\bar{b}$).
1998 Then, I prepared the `WimpSim` output fluxes in a specific way to use them as inputs to
1999 `NuWro`, which simulates the neutrino interaction with the argon.

2000 Because `WimpSim` outputs an event list together with the fluxes, I can use the former
2001 to generate the events. The direction of these is given in terms of the azimuth and
2002 altitude angles viewed from the specified location, so first I need to convert these into the
2003 DUNE FD coordinates. Once I have done it, each event can be processed with `NuWro`.
2004 To increase the number of samples and optimise the computation time, I generate 100
2005 interactions (i.e. `NuWro` events) for each `WimpSim` event². I restrict the event generation
2006 to charged current interactions, but I allow all the different contributions to the CC
2007 cross section, i.e. quasielastic scattering (QEL), meson exchange current process (MEC),
2008 resonant pion production (RES) and deep inelastic scattering (DIS). I just take into
2009 account the CC contribution because I am only interested in final states with charged
2010 leptons, as we have better chances of reconstructing the kinematics of CC events.

2011 For the atmospheric fluxes I follow a similar procedure, only that this time I do not
2012 have a set of events but the fluxes binned in azimuth and altitude angles. This way, I
2013 transform these to DUNE coordinates and process the fluxes for each bin separated with

²This also solves a problem related with the generation of the neutrino interactions in `NuWro`, as if you only produce one event each time you launch `NuWro` it will always produce an interaction of the dominant interaction type for that particular energy.

7.6. High energy DM neutrino signals




Figure 7.10: Distribution of the muon neutrino energies from the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilation channels, for $m_{\text{DM}} = 10 \text{ GeV}$, separated by CC interaction type: QEL (blue), MEC (orange), RES (green) and DIS (red).

2014 NuWro.

2015 At this point, I have two sets of events with different energies and final states.
 2016 In Fig. 7.10 one can see the distribution of the muon neutrino energies for the case
 2017 $m_{\text{DM}} = 10 \text{ GeV}$, both for the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) channels, separated
 2018 by interaction. One can clearly see that there are different energy regimes where the
 2019 primary interaction type is different. This leads to a plurality of event topologies,
 2020 therefore making it difficult to implement a general approach to the selection of events
 2021 in detriment of the background. As a way to proceed, I decided to split our samples,
 2022 based on the different interaction modes and contents of the final state, into a CC DIS
 2023 sample and a single proton CC QEL sample.

2024 **7.6.1 DIS events**

2025 To begin with, I consider the high energy part of the spectrum. In this region DIS events
 2026 dominate, i.e. interactions of the form $\nu_\mu + q_d(\bar{q}_u) \rightarrow \mu^- + q_u(\bar{q}_d)$. Therefore, our final
 2027 estates will contain a muon and a hadronic jet from the fragmentation of the outgoing
 2028 quark. As all these events have $E_\nu \gtrsim 1 \text{ GeV}$ the momentum transfer to the remnant
 2029 nucleus is negligible, for this reason the neutrino energy can be effectively reconstructed

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.11: Distributions of θ_μ (left panel), θ_j (central panel) and θ_{plane} (right panel) for the $b\bar{b}$ sample with $m_{\text{DM}} = 10 \text{ GeV}$ (blue) and the atmospheric background (red).

2030 just taking into account the momenta of the muon and the jet. This technique was
 2031 successfully used in Ref. [86] to select monoenergetic DM solar neutrino events from $\nu\bar{\nu}$
 2032 annihilation channels.

2033 Using momentum conservation one sees that the plane generated by the momenta
 2034 of the muon and the jet needs to also contain the momentum of the neutrino. As we
 2035 are interested in neutrinos coming from the Sun, the momentum of the neutrino can be
 2036 regarded as known beforehand. This will allow us to define the angle of the outgoing
 2037 muon and jet with respect to the incoming neutrino. Moreover, one can also use that
 2038 information to reject poorly reconstructed jets, checking for deviations of these from the
 2039 momentum conservation plane.

2040 To account for the limited angular resolution of the detector, I smeared the momenta
 2041 of the muons and hadrons. In a liquid argon TPC muons are expected to be tracked with
 2042 high precision, therefore I take the associated angular resolution to be 1° . In the case of
 2043 jets, it is expected that for the hadrons dominating the cascade a detector like DUNE
 2044 has an angular resolution between 1° to 5° [42], so I take the latter, more conservative,
 2045 estimate.

2046 As a first selection step, I will just take into account particles with kinetic energies
 2047 above the detection threshold of DUNE. For muons and photons the specified threshold
 2048 energy is 30 MeV, for charged pions 100 MeV and for other hadrons 50 MeV [42]. This
 2049 way, if the outgoing muon in a certain event has an energy lower than the required

7.6. High energy DM neutrino signals




Figure 7.12: Left panel: signal efficiencies (blue lines) and background rejections (red lines) for events passing the cuts $\theta < \theta_{\text{cut}}$ for the jet (solid lines) and muon (dashed lines) angles. Right panel: signal efficiency (blue line) and background rejection (red line) for events passing the cut $\theta_{\text{plane}} < \theta_{\text{cut}}$ for the momentum conservation plane deviation.

threshold I will drop such event. For the case of hadrons and photons, I will only require to have at least one particle above the energy threshold, so then one can compute the jet momentum using the (smeared) momenta of the N particles above threshold as:

$$\vec{p}_j = \sum_{i=1}^N \vec{p}_i. \quad (7.28)$$

Additionally, I will also define an estimation of the deposited hadronic energy as:

$$E_j^{\text{dep}} = m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}} + \sum_{i=1}^N \sqrt{|\vec{p}_i|^2 + m_i^2}. \quad (7.29)$$

This quantity is useful to select events with enough hadronic visible energy in the detector. For events where most of the hadronic energy is scattered across plenty of hadrons with individual energies below the detection threshold, this estimation will give $E_j^{\text{dep}} \leq 0$. In these cases it could be expected that the jet momentum is poorly reconstructed, and therefore I require events to pass the cut $E_j^{\text{dep}} > 0$.

For the events I can compute the angles for the muon and jet with respect to the

Chapter 7. DM searches with neutrinos from the Sun

2060 incoming neutrino as:

$$\cos \theta_\mu = \hat{p}_\nu \cdot \hat{p}_\mu, \quad (7.30)$$

$$\cos \theta_j = \hat{p}_\nu \cdot \hat{p}_j, \quad (7.31)$$

2061 and the deviation from the momentum conservation plane as:

$$\sin \theta_{plane} = \left| \frac{\hat{p}_\mu \times \hat{p}_\nu}{|\hat{p}_\mu \times \hat{p}_\nu|} \cdot \hat{p}_j \right|. \quad (7.32)$$

2062 In Fig. 7.11 I show some distributions of these quantities for the case of the $b\bar{b}$ sample
 2063 with $m_{DM} = 10$ GeV (blue histograms) and for the atmospheric backgrounds (red).
 2064 In order to select the atmospheric events I followed the same criteria as for the signal
 2065 events. However, because in the signal case I used the true direction of the neutrino
 2066 as input, as it should be that of the Sun at that time and therefore known, in the
 2067 atmospheric case I used a set of solar positions as our ansatz for the neutrino direction.
 2068 From the distributions, one can see that the muon and the jet for the signal events are
 2069 predominantly forward and also that the deviations from the momentum conservation
 2070 plane are peaked at zero, as one should expect.

2071 Now, I can start applying cuts to maximise our signal selection efficiency while at
 2072 the same time I try to minimise the amount of atmospheric background events passing
 2073 the selection. To this end, I will need to find some lower and upper cuts for θ_j and
 2074 θ_μ and an upper bound for θ_{plane} . In Fig. 7.12 I show how upper bound cuts in the
 2075 different angular variables affect the signal efficiency (blue lines) and the background
 2076 rejection (red lines). Notice that the signal efficiency behaves in a quite different way
 2077 when I apply cuts in the jet and the muon angles. On the contrary, the cuts on both
 2078 variables have a similar effect on the background rejection.

2079 In order to obtain the optimal set of cuts, I perform a multidimensional scan. I
 2080 do this separately for the $\tau^+\tau^-$ and the $b\bar{b}$ samples. For each case, I scan the possible
 2081 cuts for each mass point and then I take the mean value of the signal efficiency for

7.6. High energy DM neutrino signals




Figure 7.13: Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) DIS samples as functions of the DM mass, m_{DM} , obtained by applying the optimal angular cuts $\theta_\mu < 27^\circ$, $4^\circ < \theta_j < 26^\circ$ and $\theta_{\text{plane}} < 3.5^\circ$.

2082 each configuration, to get the mean efficiency for each set of cuts. I do a similar scan
 2083 for the atmospheric sample independently. Then, I take the sets of cuts such that
 2084 the background rejection achieved is greater than 99.8% and search for the one which
 2085 maximises the $\tau^+\tau^-$ and $b\bar{b}$ sample mean efficiencies. I found that with the cuts $\theta_\mu < 27^\circ$,
 2086 $4^\circ < \theta_j < 26^\circ$ and $\theta_{\text{plane}} < 3.5^\circ$ I get a background rejection of 99.80% while achieving
 2087 a 49.40% and 44.92% mean signal efficiencies for the $\tau^+\tau^-$ and $b\bar{b}$ signals respectively.

2088 In Fig. 7.13 I show the signal efficiencies as a function of the DM mass for the $\tau^+\tau^-$
 2089 (blue line) and the $b\bar{b}$ (red line) DIS events, after applying the cuts discussed above, as
 2090 well as the energy threshold and hadronic visible energy selections. One can see that
 2091 the efficiency grows with the mass, as annihilations of more massive DM particles will
 2092 produce a neutrino spectrum centered at higher energies, where DIS events dominate.
 2093 Notice also that the efficiency is higher for the $\tau^+\tau^-$ case at every mass point, as in
 2094 general this channel produces neutrinos at higher energies than the corresponding $b\bar{b}$
 2095 channel.

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.14: Distributions of $\cos \theta_\mu$ (left panel), $\cos \theta_p$ (central panel) and $\cos \theta_N$ (right panel) for the $\tau^+\tau^-$ QEL sample with $m_{\text{DM}} = 5$ GeV (blue) and the atmospheric background (red).

2096 7.6.2 Single proton QEL events

2097 Now, one can try to explore the low energy tail of the neutrino energy distributions. This
 2098 regime is dominated by the QEL interactions, i.e. events of the type $\nu_\mu + n \rightarrow \mu^- + p$.
 2099 In this case, as the typical energies are $E_\nu \lesssim 1$ GeV, the momentum transfer to the
 2100 remnant nucleus is sizeable. Therefore, I can not make the approximation I did before
 2101 and assume that the momentum of the muon and the proton will give an adequate
 2102 estimation of the reconstructed neutrino energy.

2103 In any case, as before, I can take the direction of the incoming neutrino as known.
 2104 That way, one can estimate the energy of the neutrino as:

$$E_\nu^{reco} = E_\mu + E_p + m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}}, \quad (7.33)$$

2105 and using momentum conservation I can write the momentum of the remnant nucleus
 2106 as:

$$\vec{p}_N = \hat{p}_\nu (E_\mu + E_p + m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}}) - \vec{p}_\mu - \vec{p}_p. \quad (7.34)$$

2107 As in the previous case, I need to drop the events where the muon or the proton fall
 2108 below the kinetic energy detection threshold [42]. Also, I again apply a smearing to the
 2109 momenta of the particles, a 1% for muons and 5% for protons.

2110 Having done that, one can compute the following angular variables for our selected

7.6. High energy DM neutrino signals




Figure 7.15: Left panel: value of the loss function for the training sample (blue line) and accuracy for the validation sample (red line) versus the number of iterations for the MLP classifier training. Right panel: distributions of the predicted probabilities assigned by the MLP classifier to the test sample for the $\tau^+\tau^-$ QEL signal with $m_{\text{DM}} = 5 \text{ GeV}$ (blue) and the atmospheric background (red).

2111 events:

$$\cos \theta_\mu = \hat{p}_\nu \cdot \hat{p}_\mu, \quad (7.35)$$

$$\cos \theta_p = \hat{p}_\nu \cdot \hat{p}_p, \quad (7.36)$$

$$\cos \theta_N = \hat{p}_\nu \cdot \hat{p}_N. \quad (7.37)$$

2112 Fig. 7.14 shows the distributions of these angular variables for the $\tau^+\tau^-$ QEL
 2113 sample with $m_{\text{DM}} = 5 \text{ GeV}$ (blue) and the atmospheric background (red). Again, for
 2114 the atmospheric events I used a random solar position as the ansatz for the incoming
 2115 neutrino direction. Notice that now, opposed to the DIS case where the signal had very
 2116 sharp distributions for the variables considered, the shapes of the angular distributions
 2117 for signal and background are not that much different.

2118 This effectively means that the usual approach of applying simple angular cuts would
 2119 not work as well as in the previous situation. Therefore, as a possible solution, I tried to
 2120 use a multilayer perceptron (MLP) classifier to separate between signal and background
 2121 events. Thus, the power of the hypothesis test will serve as an estimate of the signal

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.16: Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) single proton QEL samples as functions of the DM mass, m_{DM} , obtained by requiring a minimum predicted probability from the MLP classifier of 0.97 in order to achieve a background rejection greater than 99.8%.

efficiency, and in the same way one can take the size of the test to be our background rejection.

For each DM mass value and channel, as well as for the background sample, I divide our events into training, validation and test samples. The input variables for the classifier were the reconstructed neutrino energy from Eq. (7.33) and the angular variables defined in Eqs. (7.35 - 7.37). I used the MLP classifier implemented in `scikit-learn` [87], with a total of five hidden layers, the rectified linear unit activation function and adaptive learning rate. In order to account for fluctuations due to artifacts in the training process I repeated the training a thousand times for each sample, redefining each time the training, validation and test subsets, so one can take as our signal efficiency and background rejection the mean values of the powers and sizes of the tests.

The results of one of these training processes for the $\tau^+\tau^-$ QEL signal with $m_{\text{DM}} = 5$ GeV is shown in Fig. 7.15. On the left panel I show the loss function values (blue) and accuracy (red) at each iteration for the training and the validation samples respectively.

7.6. High energy DM neutrino signals

2136 The training stops either when the maximum number of iterations is reached (1000 in
2137 this case) or when the accuracy for the validation sample reaches a certain tolerance
2138 (I chose 10^{-4} as our tolerance). On the right panel I have the distributions for the
2139 predicted probability by the model, separated in true signal (blue) and background
2140 (red) events, for the test sample. One can see that both populations are well separated,
2141 obtaining a power of 44.97% and a size of 0.17% when I require a predicted probability
2142 greater than 0.97.

2143 Applying this criteria for each sample, I obtain the mean signal efficiencies shown in
2144 Fig. 7.16. Notice that the efficiencies for the channel $\tau^+\tau^-$ (blue line) are consistently
2145 lower than the ones for the $b\bar{b}$ channel (red line). This can be due to the fact that, for
2146 each DM mass point, the neutrino spectrum coming from the $b\bar{b}$ annihilation channel is
2147 centered at lower energies when compared to the $\tau^+\tau^-$ spectrum. This directly translates
2148 into more low energy neutrinos undergoing QEL interactions, which give signals that
2149 can be easily separated from the atmospheric background. This explanation also help us
2150 understand why in both cases the signal acceptance drops when the DM mass increases.
2151 In all cases, the background rejection took values between 99.8% to 99.9%. I will assume
2152 a 99.8% background rejection value in all cases to keep our estimation conservative.

2153 7.6.3 Results

2154 In order to estimate the DM-nucleon cross section sensitivities in the present case I need
2155 again to compute the expected number of background events. As I am now separating
2156 events by interaction type Eq. (7.25) does not hold anymore, as in that case I integrated
2157 over the total neutrino-argon cross section. In this instance, the expected background
2158 events for DIS events is approximately given by:

$$N_B^{DIS} \simeq \eta_B^{DIS} \times (4.655 \times 10^3) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right), \quad (7.38)$$

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.17: Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin-dependent DM-nucleon scattering cross section as a function of m_{DM} , for the annihilation channels $\tau^+\tau^-$ (blue) and $b\bar{b}$ (red) separated by interaction type (up triangles denote DIS interactions whereas down triangles represent QEL interactions). I also show the previous limits from IceCube [9] (solid lines) and the projected sensitivities for Pingu [10] (dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines), as well as the direct detection limits from PICASSO [12] (solid green line) and PICO-60 C₃F₈ [13] (dashed green line).

2159 whereas for QEL events we have:

$$N_B^{QEL} \simeq \eta_B^{QEL} \times (2.248 \times 10^4) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right). \quad (7.39)$$

2160 Now, using these together with Eqs. (7.26) and (7.27) one can obtain the 90% C.L.
 2161 upper limit on the total annihilation rate at equilibrium for both kind of events. Then,
 2162 applying the computed DM-nucleons capture rates I can translate these into limits on
 2163 the DM-nucleon cross section by means of Eqs. (7.2), (7.5) and (7.6).

2164 Fig. 7.17 shows the obtained limits on the SD DM-nucleon cross section for DUNE,
 2165 using the DIS (up triangles) and QEL (down triangles) events both for the $\tau^+\tau^-$ (blue)
 2166 and the $b\bar{b}$ (red) samples, for an exposure of 400 kT yr. I also include the corresponding

7.7. Example: Leptophilic Dark Matter

2167 current limits from IceCube [9] (solid lines), as well as the projected sensitivities of Pingu
2168 [10] (dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines). For comparison, I
2169 also show the reported direct detection limits from PICASSO [12] (solid green line) and
2170 PICO-60 C₃F₈ [13] (dashed green line).

2171 Notice that, for most of the mass range, the limits one can set by using the DIS
2172 events are stronger than those of the QEL interactions, except for the low mass part
2173 of both the $\tau^+\tau^-$ and the $b\bar{b}$ curves where the QEL events dominate. In general, the
2174 expected sensitivity of DUNE for DM masses $\lesssim 25$ GeV surpasses the stronger current
2175 indirect limits. However, experiments like Hyper-Kamiokande are foreseen to have an
2176 overall better sensitivity in this kind of searches, as they have a bigger active volume
2177 and accept a broader energy range.

2178 A pending question is what happens when we add the RES and MEC charged-current
2179 interaction contributions. In that case it would probably be more convenient to split
2180 the samples by final state interaction topologies. Also, another necessary improvement
2181 would be adding a full detector simulation and reconstructions. This will also require
2182 considering the effect of poorly reconstructed events or final states containing neutral
2183 particles such that they mimic the desired topology at the reconstruction level.

2184 7.7 Example: Leptophilic Dark Matter

2185 In general, the capture rate of DM particles by the Sun via interactions with electrons is
2186 several orders of magnitude smaller than the capture via DM-nucleus scattering. Thus,
2187 it would be sub-leading even when nucleon capture is loop suppressed. As I showed in
2188 Fig. 7.2, the capture rate via scattering off electrons only surpasses the capture rates
2189 via DM-nucleons interactions for DM masses $\lesssim 100 - 500$ MeV.

2190 However, if one considers a model where DM-nucleon interactions are forbidden even
2191 at loop level, then electron interactions will be the sole contributor to DM capture in
2192 the Sun. One can describe such scenario where the DM particles couple to leptons but
2193 not to the quark sector using effective operators.

Chapter 7. DM searches with neutrinos from the Sun

2194 In general, assuming that the DM particle is a Dirac fermion, the dimension six
 2195 operators describing the interaction between two DM particles and two leptons can be
 2196 written as:

$$\mathcal{L}_{eff} = G \sum_i (\bar{\chi} \Gamma_\chi^i \chi) (\bar{\ell} \Gamma_\ell^i \ell), \quad (7.40)$$

2197 where $G = 1/\Lambda^2$ is the effective coupling strength, Λ the cut-off of the effective field
 2198 theory and ℓ denotes any lepton. In principle, one should consider all the possible
 2199 Lorentz structures Γ_f^i in order to have a complete set of effective operators.

2200 However, some combinations will induce interactions with nucleons at loop level. As
 2201 we are specifically interested in interactions which forbid any communication with the
 2202 quark sector, I will not consider those [88]. In addition, some of the effective operators
 2203 give rise to velocity-suppressed scattering cross sections between DM particles and
 2204 leptons. I will also neglect those, as the suppression goes with the square of the DM
 2205 halo velocity which in units of the speed of light is $\sim 10^{-6}$.

2206 This way, the only Lorentz tensor structure that do not induce interactions with
 2207 quarks at loop level and gives a contribution to the scattering cross section that is not
 2208 velocity suppress is the axial-axial interaction. The effective Lagrangian is then given
 2209 by:

$$\mathcal{L}_{eff} = \frac{c_A^\chi c_A^\ell}{\Lambda^2} (\bar{\chi} \gamma^\mu \gamma^5 \chi) (\bar{\ell} \gamma_\mu \gamma^5 \ell), \quad (7.41)$$

2210 where c_A^χ and c_A^ℓ are the couplings for the different species. As the DM coupling appears
 2211 as a common factor for any lepton choice, I will redefine the corresponding coupling c_A^ℓ
 2212 to absorb c_A^χ . Also, for simplicity, I will assume that the couplings between the DM
 2213 particles and the leptons are flavour independent, i.e. I have just two couplings, c_A^e for
 2214 charged leptons and c_A^ν for neutrinos.

2215 In the case of a scalar DM particle, the lowest order effective interaction with
 2216 leptons happens through a dimension five operator, generating scalar and pseudoscalar
 2217 interactions. However, the former induces interactions with quarks at two loop level
 2218 whereas the latter gives a velocity suppressed scattering cross section.

2219 From the effective Lagrangian in Eq. (7.41) it can be shown that the axial-axial

7.7. Example: Leptophilic Dark Matter

contribution to the scattering cross section for the fermionic DM and a charged lepton
is given by:

$$\sigma_{\text{DM}-e}^{AA} = 3(c_A^e)^2 \frac{m_e^2}{\pi \Lambda^4}. \quad (7.42)$$

If the DM interacts exclusively with fermions, then the only annihilation channels that will give us a measurable neutrino flux coming out of the Sun are $\tau^+\tau^-$ and $\nu\bar{\nu}$. The former channel, already explored previously in the more mainstream scenario of the DM capture via scattering off nucleons, is open only for $m_{\text{DM}} > m_\tau \simeq 1776.86 \pm 0.12$ MeV [89], a mass region where the solar DM capture by electrons is at least one order of magnitude smaller than the capture via interactions with nucleons. On the contrary, the latter allows us to explore a region where the capture rate via scattering off electrons dominates over the rest.

One downside of focusing in such low mass range is that it falls below the usual limit of $m_{\text{evap}} \sim 4$ GeV usually explored in the literature. The pretext to explore this region is the result discussed previously reported in Ref. [4], where DM evaporation in the Sun for the case of capture via electron scattering could be negligible for masses as low as $m_{\text{evap}} \sim 200$ MeV. This result is quite sensitive to the high velocity tail of the DM velocity distribution in equilibrium inside the Sun, and therefore full numerical simulations would be needed to assess the impact of this effect. However, this falls out of the scope of our work.

In this case, as I have a specific realisation of the interaction between the DM and leptons, one can estimate the relic density of our DM for different values of the couplings and the effective field theory scale Λ . The first step to do so is compute the self-annihilation cross section. Because I consider cold relics, at the freeze-out time our DM particles were non-relativistic and so one can expand the annihilation cross section in terms of the relative velocity v between two annihilating DM particles as [90]:

$$\sigma_{\text{ann}}^{AA}|v| \approx \frac{1}{2\pi\Lambda^4} \sum_\ell \left(c_A^\ell\right)^2 m_\chi^2 \sqrt{1 - \frac{m_\ell^2}{m_\chi^2} \left[\frac{m_\ell^2}{m_\chi^2} + \frac{1}{12} \left(2 - \frac{m_\ell^2}{m_\chi^2}\right) v^2 \right]}, \quad (7.43)$$

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.18: Left panel: Projected 90% confidence level sensitivity of DUNE (400 kT yr) to the scale Λ of an EFT containing only leptophilic DM axial-axial interactions (blue line). Right panel: In both cases the corresponding limits from DarkSide-50 [14] (dotted green line) and XENON1T [15] (dashed red line) are also shown, together with the configurations for which the correct relic density is achieved (black line), all for the coupling values $c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$.

2244 where the sum includes all the possible lepton final states with mass m_ℓ .
 2245 Solving the Boltzmann equation for the evolution of the DM density gives as a
 2246 solution a relic density of:

$$\Omega_\chi h^2 \approx \frac{(1.04 \times 10^9) x_F}{M_{Pl} \sqrt{g_*} (a + 3b/x_F)}, \quad (7.44)$$

2247 where $x_F = m_\chi/T_F$ being T_F the freeze-out temperature, g_* the number of relativistic
 2248 degrees of freedom at freeze-out and a and b the terms in the annihilation cross section
 2249 expansion $\sigma_{ann}|v| \approx a + bv^2 + \mathcal{O}(v^4)$. Using the current best fit for the relic DM density
 2250 $\Omega_\chi h^2 = 0.1198 \pm 0.0012$ [91] one can use these relations to compute the required effective
 2251 theory scale Λ at which the correct density is achieved for any combinations of m_χ and
 2252 c_A^ℓ .

2253 As discussed before, in the low DM mass region QEL interactions dominate. Moreover,
 2254 if I focus on direct annihilation to neutrinos, the energy of the muon neutrino flux is
 2255 known as it must be equal to the mass of the DM particle, $E_\nu = m_\chi$. That way, now
 2256 I do not need to use Eq. (7.33) in order to estimate the momentum transfer to the

7.7. Example: Leptophilic Dark Matter

2257 remnant nucleus, I can simply take:

$$\vec{p}_N = \hat{p}_\nu m_\chi - \vec{p}_\mu - \vec{p}_p. \quad (7.45)$$

2258 To estimate the signal efficiency and background rejection for this case I used again
2259 the MLP classifier from `scikit-learn`, using the same specifications as before. The
2260 only difference now is that I add also the reconstructed neutrino energy as one of the
2261 features to train the classifier with, because the characteristic monoenergetic flux for
2262 each m_χ value will help to distinguish between signal and background events.

2263 In this case, for masses below ~ 500 MeV I obtain a signal efficiency close to unity
2264 while keeping a background rejection of 99.9%. For bigger values of the mass, the signal
2265 efficiency drops significantly if I require to keep the background acceptance under 0.01%.
2266 However, because this kind of search is dominated by the background, sacrificing the
2267 signal acceptance to keep the background rejection to a minimum enhances the reach
2268 of the analysis. This way, for DM masses of the order of $m_\chi \sim 1$ GeV I end up with
2269 efficiencies as low as 1%.

2270 Now, estimating the number of background events using Eq. (7.39) one can go on
2271 and apply Eqs. (7.26) and (7.27) together with Eq. (7.42) to derive the sensitivity of
2272 DUNE to this kind of model. Fig. 7.18 (left panel) shows the potential reach of DUNE
2273 to constrain the EFT scale Λ this model containing only leptophilic DM axial-axial
2274 interactions (blue line), for a choice of couplings $c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$. I also included
2275 the current limits on the DM-electron scattering cross section from DarkSide-50 [14]
2276 (dotted green line) and XENON1T [15] (dashed red line), reworked with Eq. (7.42) to
2277 show their implications for the EFT scale. The values of Λ for which the correct DM relic
2278 density value is achieved for each mass are also shown (black line). This tells us that,
2279 for that specific choice of couplings, DUNE would be sensitive to DM configurations
2280 allowed by the relic density constraint up to a mass of $m_\chi \sim 400$ MeV.

2281 In Fig. 7.18 (right panel) I show the same upper limits but for the DM-electron
2282 scattering cross section. From this view one can see that DUNE would be able to

Chapter 7. DM searches with neutrinos from the Sun

offer complementary information to the low energy DM-electron interaction searches performed by direct detection experiments, in a slightly higher mass range.

With the present example, although it focuses on a very specific realisation of the DM interactions, I show the potential of DUNE to constrain exotic DM scenarios. Thanks to its low backgrounds and superb angular resolution DUNE will be able to help with the systematic searches for dark sectors physics.

2289 Chapter 8

2290 Particle ID in GArSoft

2291 ND-GAr is a magnetised, high-pressure gaseous argon TPC (HPgTPC), surrounded by
2292 an electromagnetic calorimeter (ECal) and a muon detector (commonly refer to as μ ID).
2293 A detailed discussion on the requirements, design, performance and physics of ND-GAr
2294 can be found in the DUNE ND CDR [48] and the ND-GAr whitepaper (cite).

2295 In DUNE Phase II ND-GAr will fulfill the role of TMS, measuring the momentum
2296 and sign of the charged particles exiting ND-LAr. Additionally, it will be able to measure
2297 neutrino interactions inside the HPgTPC, achieving lower energy thresholds than those
2298 of the ND and FD LArTPCs. By doing so ND-GAr will allow to constrain the relevant
2299 systematic uncertainties for the LBL analysis even further.

2300 The goal of the present chapter is to review the requirements that the physics program
2301 of DUNE impose on ND-GAr, present the current status of its design and describe the
2302 GArSoft package, its simulation and reconstruction software.

2303 As decided during the DUNE Phase II workshop in June 2023 [reference], we want
2304 to build ND-GAr physics case by showing:

- 2305 • That ND-GAr can constrain systematic uncertainties that ND-LAr might miss.
- 2306 • The impact on the neutrino oscillation results if such systematic uncertainties are
2307 missed.
- 2308 • That ND-GAr is necessary to reach DUNE's main physics goals.

Chapter 8. Particle ID in GArSoft

2309 This way, the design of ND-GAr will be physics driven.

2310 In order to study the effects of final state interactions (FSI) in CC interactions,
2311 ND-GAr should be able to measure the spectrum of protons and charged pions at low
2312 energies. ND-GAr also needs to be able to measure the pion multiplicity, specially for
2313 energies above 100 MeV as at these energies the pions shower in the LAr, to inform the
2314 pion mass correction in the ND and FD LArTPCs.

2315 In order to correctly identify electrons, muons, pions, kaons and protons ND-GAr
2316 can use a combination of: dE/dx measurements in the HPgTPC, E_{ECAL}/p using the
2317 ECAL total energy and the momentum obtained from magnetic spectroscopy in the
2318 HPgTPC and penetration information through the ECAL and muon tagger.

2319 8.1 dE/dx measurement in the TPC

2320 Among the parameters extracted from the track fitting, ionisation is particularly useful
2321 for particle identification, as it is a function of the particle velocity. Although for the
2322 case of relativistic particles this dependence is not very strong, measuring the track on
2323 a large number of points may allow us to estimate the amount of ionisation accuratel.
2324 This, paired with a measurement of the momentum, may allow us to identify the particle
2325 type.

2326 The first calculation of the energy loss per unit length of relativistic particles using
2327 a quantum-mechanical treatment is due to Bethe [?]. Using this approach, the mean
2328 ionisation rate of a charged particle traveling through a material medium is (using
2329 natural units $G = \hbar = c = 1$):

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{4\pi Ne^4}{m_e \beta^2} z^2 \left(\log \frac{2m_e \beta^2 \gamma^2}{I} - \beta^2 \right), \quad (8.1)$$

2330 where N is the number density of electrons in the medium, e the elementary charge, m_e
2331 is the electron mass, z the charge of the particle in units of e , β is the velocity of the
2332 particle, $\gamma = (1 - \beta^2)^{-1}$ and I denotes the effective ionisation potential averaged over
2333 all electrons. This relation is known as the Bethe-Bloch formula.

8.1. dE/dx measurement in the TPC

From Eq. (8.1) one can see that the ionisation loss does not depend explicitly on the mass of the charged particle, that for non-relativistic velocities it falls as β^{-2} , then goes through a minimum and increases as the logarithm of γ . This behaviour at high velocities is commonly known as the relativistic rise. The physical origin of this effect is partly due to the fact that the transverse electromagnetic field of the particle is proportional to γ , therefore as it increases so does the cross section.

It was later understood that the relativistic rise could not grow indefinitely with γ . A way to add this feature in the Bethe-Bloch formula is by introducing the so-called density effect term. It accounts for the polarisation effect of the atoms in the medium, which effectively shield the electromagnetic field of the charged particle halting any further increase of the energy loss [?]. Denoting the correction as $\delta(\beta)$, one can rewrite Eq. (8.1) as:

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{4\pi Ne^4}{m_e \beta^2} z^2 \left(\log \frac{2m_e \beta^2 \gamma^2}{I} - \beta^2 - \frac{\delta(\beta)}{2} \right). \quad (8.2)$$

In general, the form of $\delta(\beta)$ depends on the medium and its state of aggregation, involving the usage of tabulated parameters and implicit relations [?].

Another standard method to compute the amount of ionisation a charged particle produces is the so-called photo-absorption ionisation (PAI) model proposed by Allison and Cobb [?]. Within their approach, the mean ionisation is evaluated using a semiclassical calculation in which one characterises the continuum material medium by means of a complex dielectric constant $\epsilon(k, \omega)$. However, in order to model the dielectric constant they rely on the quantum-mechanical picture of photon absorption and collision. Therefore, in the PAI model the computation of the ionisation loss involves a numerical integration of the measured photo-absorption cross-section for the relevant material.

In a particle physics experiment, the typical way of determining the energy loss per unit length as a function of the particle velocity is studying identified particles over a range of momenta. Once we have established this relation we can use it for other, unknown particles. In this sense, it makes sense to have a regular mathematical expression for this relation that one can use.

Chapter 8. Particle ID in GArSoft

2361 It happens that neither the Bethe-Bloch theory nor the PAI model from Allison and
2362 Cobb offer a close mathematical form for the ionisation curve. This is the reason why a
2363 full parametrisation of the ionisation curves can be useful. A parametrisation originally
2364 proposed for the ALEPH TPC [?] and later used by the ALICE TPC [?] group that
2365 manages to capture the features of the ionisation energy loss is:

$$f(\beta\gamma) = \frac{P_1}{\beta^{P_4}} \left(P_2 - \beta^{P_4} - \log \left[P_3 + \frac{1}{(\beta\gamma)^{P_5}} \right] \right), \quad (8.3)$$

2366 where P_i are five free parameters. Hereafter, we will refer to Eq. (8.3) as the ALEPH
2367 dE/dx parametrisation.

2368 8.1.1 Energy calibration

2369 In order to obtain the amount of energy loss by a charged particle due to ionisation
2370 in our TPC we need to determine the conversion between the charge deposited in our
2371 readout planes and the actual energy depositions. This procedure is known as energy
2372 calibration.

2373 In a general, the first step of the calibration involves a non-uniformity correction,
2374 to make sure that the detector response is uniform throughout the TPC. These are
2375 typically divided into three categories, non-uniformities in the transverse YZ plane,
2376 non-uniformities along the drift direction X and variations of the detector response
2377 over time (would not apply to us as the detector is not built yet). These would correct
2378 for effects such as electron diffusion and attenuation, space charge effects or channel
2379 misconfiguration. However, because at the moment I am only interested in making sure
2380 we recover a sensible result from our simulation, I will not apply uniformity corrections
2381 to our charge deposits.

2382 Other effects, like electron-ion recombination or ADC saturation, lead to a non-linear
2383 relation between the observed charge and the deposited energy in the detector, with the
2384 observed readout charge saturating at high ionisation energies. In this case, because we
2385 are dealing with gaseous argon and therefore recombination is not as important as in

8.1. dE/dx measurement in the TPC




Figure 8.1: Left panel: distribution of the fraction of Geant4-level energy deposits per track with residual range less than 20% of the total track length, for the isotropic proton sample. Right panel: distribution of the ionisation per unit length of the energy deposits in the proton sample after removing the tracks with less than 30% of their energy deposits in the last 20% of the track.

2386 liquid, we do not simulate recombination effects in the TPC. Even so, the simulation of
 2387 the electronic response will still introduce charge saturation, and one needs to correct
 2388 for it in order to obtain the exact amount of energy loss due to ionisation.

2389 By default, the track fitting algorithm in GArSoft provides a `TrackIonization`
 2390 object associated to each reconstructed track. It contains two collections of charge
 2391 deposits, one for each fitting direction, consisting on pairs of charge values (dQ , in ADC)
 2392 and step sizes (dx , in cm).

2393 In order to estimate the ionisation loss in the ND-GAr TPC, I have used an MC
 2394 sample consisting of single, isotropic protons propagating in the TPC. The starting points
 2395 of the protons were sampled inside a $50 \times 50 \times 25$ cm box centered at $(100, -150, 1250)$,
 2396 and their momenta are uniformly distributed in the range $0.25 - 1.75$ GeV. I ran the
 2397 simulated sample through GArSoft's default detector simulation and reconstruction, and
 2398 then a custom analyser module that extracts the ionisation data together with other
 2399 reconstructed track information from the Kalman fit.

2400 For studying the energy loss of the protons I select the reconstructed tracks that
 2401 range out (i.e. slow down to rest) inside the TPC. A characteristic feature of the energy
 2402 loss profile of any stopping ionising particle is the so-called Bragg peak, a pronounced

Chapter 8. Particle ID in GArSoft




Figure 8.2: Left panel: distribution of the reconstructed ionisation charge per unit length for our MC stopping proton sample. The different colors indicate how many consecutive dQ/dx pairs were grouped together. Right panel: distribution of the median change in dQ/dx per track after $N_{group} = 4$ clusters were reclustered together.

peak that occurs immediately before the particle comes to rest. From Eq. (8.1) we can see that this behaviour is expected, as the energy loss for non-relativistic particles is inversely proportional to β^2 . In data, a way of identifying the Bragg peak, and thus select the stopping particles, is checking the number of energy deposits towards the end of the track. In this case, I count the fraction of the Geant4 simulated energy deposits with a residual range value (the distance from a given energy deposit to the last deposit in the track trajectory) less than a 20% of the corresponding track length¹. The distribution of this fraction of energy deposits for our proton sample is shown in Fig. 8.1 (left panel). We can clearly see two well separated peaks in this distribution, one centered at 0.2 and another, narrower, one centered at a higher value. The first one corresponds to non-stopping protons, as in that case the number of energy deposits towards the end of the track is uniformly distributed due to the absence of the Bragg peak. In that way, I apply a cut in this distribution, requiring that at least 30% of the simulated energy deposits sit in the last 20% of the tracks, to ensure that the Bragg peak is present.

Fig. 8.1 (right panel) shows the distribution of the energy loss per unit length for the Geant4 simulated energy deposits of the selected stopping protons. We can see that it

¹As we are applying this selection at the Geant4 level we could have simply selected the stopping protons using the `EndProcess` labels from the simulation. However, the Bragg peak identification method displayed here could serve as a starting point for a selection of stopping protons in real data.

8.1. dE/dx measurement in the TPC

follows the expected shape of a Landau distribution, which describes the fluctuations of the ionisation energy losses [?]. This distribution has a characteristic asymmetric PDF, with a long right tail that translates into a high probability for high-energy ionisation losses. The origin of these fluctuations is mainly the possibility of transferring a high enough energy to an electron, so it becomes a ionising particle itself.

Now, from the point of view of the reconstruction, the objects that we have available to extract the ionisation information for the different reconstructed tracks are the collections of dQ and dx pairs, as stated before. The dQ values come from adding up the amplitude of all the reconstructed hits in a cluster, which is the input object to the Kalman fit.

Fig. 8.2 (left panel) shows the distribution of the ionisation charge deposits per unit length for the track in the stopping proton sample (blue line). As one can notice, this distribution does not resemble the expected shape of the Landau PDF. This distribution peaks sharply at 0 and has a heavy tailed behaviour. Notice, however, how the distribution changes its shape as we group together N_{group} consecutive charge deposit pairs (red, purple and green lines). The distribution in the $N_{group} = 4$ case already has a shape which resembles that of the Geant4-level ionisation per unit length, so I will proceed using this amount of reclustering for the reconstruction-level depositions.

An extra factor I need to account for, when reclustering is applied, is how the overall dQ/dx per track changes. To do so, we can look at the ratio between the median dQ/dx after and before the reclustering. Fig. 8.2 (right panel) shows the median enhancement in dQ/dx per track for the stopping proton sample in the case $N_{group} = 4$. Fitting a Landau distribution convolved with a Gaussian², I estimate the most probable value of this ratio to be $G_{group} = 2.24 \pm 0.05$.

At this point, I am left with determining the conversion between the charge deposits per unit length dQ/dx and the energy deposits per unit length dE/dx . To this end, we need a way of comparing the two. I can use the residual range z to get a prediction of

²In the literature, this distribution is often referred to as Landau+Gaussian or langau. In the following, I will use LanGauss to refer to such PDF.

Chapter 8. Particle ID in GArSoft




Figure 8.3: Distribution of the Geant4-simulated energy losses per unit length versus residual range for the stopping proton sample. The overlaid points represent the fitted most probable value of the dE/dx distribution in each residual range bin, whereas the curve is their best fit to the Bragg-Kleeman formula from Eq. (8.4).

2447 the most probable dE/dx by using the following empirical parametrisation [?]:

$$\frac{dE}{dx}(z) = \frac{z^{\frac{1}{p}-1}}{p\Lambda^{\frac{1}{p}}}, \quad (8.4)$$

2448 which is quoted in the literature as the Bragg-Kleeman formula. In order to obtain the
 2449 p and Λ parameters I perform a fit using the energy losses and the residual ranges given
 2450 by the Geant4 stage of our proton sample.

2451 Within our simulation, the residual range is sampled with a maximum size of
 2452 5 mm. Therefore, to perform the fit to the Bragg-Kleeman formula, we can use a
 2453 fine-grained residual range binning. For each of the residual range bins I extract the
 2454 dE/dx distribution and fit it to a LanGauss distribution, to obtain the value of the
 2455 most probable dE/dx in the bin together with a statistical uncertainty. I then fit Eq.
 2456 (8.4) to these most probable values and the centres of the residual range bins. This

8.1. dE/dx measurement in the TPC




Figure 8.4: Fitted most probable dQ/dx values for each dE/dx bin (red points), obtained from the stopping proton sample. The overlaid curve (black line) represents the best fit to the logarithmic calibration function from Eq. (8.5).

procedure is depicted in Fig. 8.3, where I show the distribution of the energy loss per unit length versus the residual range, together with the most probable dE/dx values and their uncertainty in each bin (red points) and the curve with the best fit of the Bragg-Kleeman relation to those values (black line). The best fit is obtained for the parameter values $p = 1.8192 \pm 0.0005$ and $\Lambda = 0.3497 \pm 0.0008$ cm/MeV³.

Having an analytical expression that relates the residual range to dE/dx , I can take our reconstruction-level residual ranges from the stopping proton sample and compute the most probable energy loss associated.

In order to parametrise the charge saturation, we can use the following logarithmic

³These strange units for Λ come from dimensional analysis, just to keep the Bragg-Kleeman formula (8.4) consistent.

Chapter 8. Particle ID in GArSoft

2466 function inspired by the modified box model for recombination:

$$\frac{dE}{dx} = \frac{e^{\frac{dQ}{dx}B\frac{W_{ion}}{G_{group}C}} - A}{B}, \quad (8.5)$$

2467 where A and B are the calibration parameters we need to determine, W_{ion} is the average
 2468 energy to produce an electron-ion pair, G_{group} is the gain from the reclustering discussed
 2469 above and C is the calibration constant to convert number of electrons to ADC counts,
 2470 commonly refer to as gain (also to be obtained in the fit). In this case, I use a value for the
 2471 electron-ion production energy of $W_{ion} = 26.4$ eV [?]. This value, used in our simulation
 2472 as well, was measured for gaseous argon in normal conditions, and therefore should
 2473 be checked in the future to describe correctly the high-pressure argon-CH₄ mixture of
 2474 ND-GAr.

2475 For the calibration fit I follow a procedure similar to the previous one for Eq. (8.4).
 2476 Binning the dE/dx range, I fit a LanGauss distribution to the corresponding dQ/dx
 2477 distribution to obtain the most probable value. The resulting data points (red bars) are
 2478 shown in Fig. 8.4 (top panel), the horizontal error bars depict the width of the dE/dx
 2479 bin whereas the vertical bars represent the error associated to the most probable value
 2480 estimation. A fit to the logarithmic function in Eq. (8.5) is also shown (black line).
 2481 For this I weighted the data points using the inverse of their relative error, obtaining a
 2482 reduced chi-square value of $\chi^2/ndf = 2.22 \times 10^{-6}$. The best fit parameters I found from
 2483 this fit are $A = 0.883 \pm 0.064$, $B = 5.6 \pm 1.3$ cm/MeV and $C = 4.94 \pm 0.49$ ADC/e. In
 2484 Fig. 8.4 (bottom panel) I show the residuals between the data points and the fit.

2485 The value for the gain I obtained from the fit is in reasonable agreement with our
 2486 expectation. This value is set in GArSoft to 5 ADC/e by default.

2487 One interesting thing to check is what induces this non-linear relation between charge
 2488 and energy. The only effects that modify the amount of electrons reaching the readout
 2489 planes in the simulation are the transverse diffusion and the finite electron lifetime.
 2490 Once the electrons reach the readout chambers, the pad response functions are applied,
 2491 together with an electrons-to-ADC conversion and the ADC saturation limit.

8.1. dE/dx measurement in the TPC




Figure 8.5: Fitted most probable dQ/dx values for each dE/dx bin for three different ADC bit limits, 10 (blue points), 12 (default, yellow points) and 16-bit (red points).

Table 8.1: Calibration parameters obtained from the fit of the ND-GAr simulated stopping proton sample to the calibration function from Eq. (8.5). The fits were performed for the 10, 12, and 16-bit ADC limits.

	χ^2/ndf	Best fit $\pm 1\sigma$		
		A	B (cm/MeV)	C (ADC/e)
10-bit	$1.83 \times 10^{-6}/12$	-9.3 ± 3.9	270 ± 69	27.1 ± 5.4
12-bit	$2.67 \times 10^{-5}/12$	0.883 ± 0.064	5.6 ± 1.3	4.94 ± 0.49
16-bit	$1.44 \times 10^{-5}/12$	0.949 ± 0.024	3.53 ± 0.58	4.52 ± 0.29

By default, GArSot applies a 12-bit ADC limit, which can be changed in the simulation configuration. However, it can only be increased up to 16-bit, as we represent the ADC collection as a `std::vector<short>`. This way, I tried to change the saturation parameter to see how it affects the relation between reconstructed charge and energy. Fig. 8.5 shows a comparison between the most probable dQ/dx for 10, 12 and 16-bit ADC limits. As expected, the lower the limit is the sooner the charge saturates. For higher ADC limits the relation between energy and charge remains linear up to higher dE/dx values, but even for the 16-bit limit the saturation is noticeable for values $\gtrsim 0.5$ MeV/cm.

In Tab. 8.1 I also show the results of fitting the samples with 10 and 16-bits ADC

Chapter 8. Particle ID in GArSoft




Figure 8.6: Top panel: area normalised dE/dx distributions for the true (solid grey) and the reconstructed energy deposits in the stopping proton sample, both after applying the calibration (blue) and the calibration and the normalisation correction (yellow). Also shown is the distribution obtained by applying a correction factor to the dQ/dx values but not the calibration (red). Bottom panel: fractional residuals for the uncorrected (blue), corrected (yellow) and uncalibrated (red) samples.

limits to the calibration function from Eq. (8.5), using the weights based on their relative error as described previously. One interesting feature to notice is how different the best fit points look for the 10-bit ADC saturation when compared to the other two, which are consistent with each other.

At this point we can compare the dE/dx distribution one gets from Geant4, i.e. the true energy loss distribution, and the distribution I found by applying the calibration function to our collection of reconstructed dQ/dx values. Fig. 8.6 (top panel) shows the true (solid grey) and reconstructed (blue, labeled as uncorrected) distributions together. The dashed vertical lines indicate the region of validity of the calibration fit, i.e. the left

8.1. dE/dx measurement in the TPC

and right edges of the first and last dE/dx bin respectively. Notice that these histograms are area-normalised, as the total number of true energy deposits is much higher than the number of reconstructed charge deposits. This is due to a combination of effects, like the finite spatial resolution of the detector, the hit clustering used in the track fitting and the reclustering we have applied here.

The two distributions are significantly different. That can be seen clearly when looking at the fractional residuals, shown in Fig. 8.6 (bottom panel). In particular, the position of the peak is off, which could bias the mean energy loss predictions. It seems like the difference between these may be due to an overall scaling factor. One possibility is to scale the most probable value of the reconstructed distribution to the most probable value predicted by Geant4. I do this by fitting both distributions using a LanGauss function, obtaining $dE/dx_{MPV, true} = 0.1145 \pm 0.0005$ MeV/cm and $dE/dx_{MPV, reco} = 0.0928 \pm 0.0005$ MeV/cm for the true and reconstructed most probable values respectively. These can be translated into a scaling factor $S = 0.579 \pm 0.006$.

The result of applying the scaling correction can be seen in Fig. 8.6 (top panel). The corrected dE/dx distribution (yellow, labeled as corrected) peaks around the same value the true distribution does, as expected. Moreover, the high energy region is also slightly better described. For low ionisations, below the lower limit of the calibration fit, the differences between true and reconstructed are still significant. This low energy excess may be migration of some events from the peak region. The overall effect of the correction can be seen in the fractional residual plot in Fig. 8.6 (bottom panel).

One can also check what happens if instead of applying the logarithmic calibration we simply scale the dQ/dx distribution (post reclustering) to have the same most probable value as the true dE/dx distribution. In this case, following an analogous procedure to the one described earlier, I found the scaling factor $S_{uncalibrated} = 0.414 \pm 0.002$ MeV/ADC⁴. The resulting distribution (red, labeled as uncalibrated) is also shown in in Fig. 8.6 (top panel). The behaviour of the new distribution is similar to the corrected case at low energy losses, around the peak of the true distribution, but it is worse at describing the

⁴Notice that now the scaling factor is not dimensionless, as it acts more like a conversion factor here.

Chapter 8. Particle ID in GArSoft




Figure 8.7: Left panel: fractional residuals between the true and the corrected dE/dx means (blue) and the 60% truncated means (yellow), for each event in the stopping proton sample. Right panel: fractional residuals between the true and the uncorrected (blue), corrected (yellow) and uncalibrated (red) dE/dx 60% truncated means, for each event in the stopping proton sample.

2538 high energy tail. This is expected, it is in the high ionisation regime where saturation
 2539 effects apply and therefore calibration is needed.

2540 **8.1.2 Truncated dE/dx mean**

2541 Once we have a collection of dE/dx values for each reconstructed track, we can compute
 2542 the corresponding most probable ionisation loss per unit length of the particle. This
 2543 is the value predicted by the Bethe-Bloch or the PAI models, and together with a
 2544 measurement of the momentum it allows for particle identification.

2545 However, estimating the most probable dE/dx value for each reconstructed track
 2546 is not a trivial task. As mentioned before, the dE/dx distributions follow Landau-like
 2547 distributions. Therefore, one should perform e.g. a LanGauss fit to correctly estimate
 2548 the most probable values. Automating this kind of fits is often problematic, as they
 2549 usually incur in convergence problems. Moreover, the reconstructed dE/dx distributions
 2550 we obtain tend to have relatively small statistics, which may also produce poor fits. In
 2551 practice, doing these unsupervised fits may degrade our performance, and a more robust
 2552 method is preferred.

2553 A possibility could be taking the mean of the reconstructed dE/dx distribution for

8.1. dE/dx measurement in the TPC




Figure 8.8: Estimated values of the mean dE/dx bias (left panel) and resolution (right panel) obtained using the corrected data from the stopping proton sample, for different values of the truncation factor.

each particle. The problem with this approach is that the high energy Landau tail, combined with our limited statistics, can induce large fluctuations in the computation of the mean. Imagine you have two protons with the same kinetic energy, but due to reconstruction problems in one case you did not get as many charge deposits reconstructed in its high ionisation loss region. If you do not remove the tails the computed dE/dx means will be significantly different.

In order to avoid those fluctuations, one can compute the mean of a truncated dE/dx distribution instead. By keeping only a given fraction of the lowest energy deposits we obtain an estimate of the mean energy loss that is more resilient to reconstruction inefficiencies and statistical effects. In Fig. 8.7 (left panel) I show a comparison between the $\langle dE/dx \rangle$ computed by taking the mean of the full distribution (blue line) and the 60% lowest energy clusters (yellow line), for the stopping proton sample. The fractional residuals are computed for each proton, taking the corresponding means using their collections of true and reconstructed energy deposits. One can see that using the simple mean translates into a high bias and uncertainty in the $\langle dE/dx \rangle$ estimation, whereas applying the truncation reduces both significantly.

Additionally, I performed a comparison between the 60% truncated mean dE/dx obtained using the different calibration methods discussed earlier, namely the uncorrected

Chapter 8. Particle ID in GArSoft

2572 (blue), corrected (yellow) and uncalibrated (red) distributions. The results are shown
 2573 in Fig. 8.7 (right panel). While the widths of these distributions are similar, the bias
 2574 obtained for the corrected sample, i.e. calibration function and correction factor applied,
 2575 is a factor of ~ 2 lower than in the uncalibrated case and almost three times smaller
 2576 than for the uncorrected sample.

2577 The next step is to optimise the level of truncation we are going to apply to our
 2578 data. To do so, I used different truncation factors, i.e. the percentage of energy-ordered
 2579 reconstructed energy deposits we keep to compute the mean, on the corrected dE/dx
 2580 sample of the stopping protons. Then, following the same procedure of computing the
 2581 fractional residuals as before, I fitted the resulting histograms using a double Gaussian
 2582 function. This is simply the sum of two Gaussian functions of the type:

$$g(x; \mu, \sigma, A) = A e^{-\frac{(x-\mu)^2}{2\sigma^2}}. \quad (8.6)$$

2583 I do not add the classical normalisation factor of the Gaussian, $1/\sqrt{2\pi}\sigma$, therefore
 2584 the amplitude A simply represents the maximum of the function. One of the two
 2585 Gaussian functions describes the core part of the distribution, while the other captures
 2586 the behaviour of the tails.

2587 For each truncation factor, I look at the bias and the resolution I obtain. I define
 2588 these as the weighted means of the corresponding parameters in the fits:

$$\bar{x} = \frac{A_{core} x_{core} + A_{tail} x_{tail}}{A_{core} + A_{tail}}, \quad (8.7)$$

2589 where A_{core} and A_{tail} are the amplitudes of the core and tail distributions respectively
 2590 and x is either the mean μ or the width σ of said distributions.

2591 Fig. 8.8 shows the bias (left panel) and the resolution (right panel) I obtained for
 2592 the stopping proton sample, using different values of the truncation. From these, it
 2593 can be seen that a truncation factor of 50% minimises the bias in the estimation, while
 2594 70% gives the best resolution. That way, I settled on the intermediate value of 60%

8.1. dE/dx measurement in the TPC




Figure 8.9: Examples of the truncated mean dE/dx LanGauss fits for various $\beta\gamma$ bins, from a simulated FHC neutrino sample.

truncation, which yields a $\langle dE/dx \rangle$ resolution of 5.00 ± 0.08 % for stopping protons.

8.1.3 Mean dE/dx parametrisation

Now that we have a way to estimate the mean energy loss of a particle in the HPgTPC, we can determine the value of the free parameters in the ALEPH formula, Eq. (8.3). For this, I used a sample of 10^5 reconstructed FHC neutrino events inside ND-GAr. In this case I cannot use the stopping proton sample, as we need to cover the full kinematic range of interest for the neutrino interactions in our detector.

The original data does not contain an estimation of the velocity of the tracks, instead the tracks have a value for the reconstructed momentum and the associated PDG code of the Geant4-level particle that created the track. Therefore, one can select some of the

Chapter 8. Particle ID in GArSoft




Figure 8.10: Resulting one and two dimensional projections of the posterior probability distributions of the ALEPH $\langle dE/dx \rangle$ parameters obtained by fitting the 60% truncated mean dE/dx values from a FHC neutrino sample in ND-GAr. The vertical dashed lines in the 1D distributions represent the 16th, 50th and 84th percentiles.

2605 particles in the data, in this case I selected electrons, muons, pions and protons, and
 2606 compute β and γ using the reconstructed momentum and their mass. In terms of $\beta\gamma$
 2607 the mean dE/dx does not depend on the particle species, so one can consider all the
 2608 dataset as a whole. For this fit, I will express β in terms of the $\beta\gamma$ product as:

$$\beta = \frac{\beta\gamma}{\sqrt{1 + (\beta\gamma)^2}}, \quad (8.8)$$

2609 which can be easily proven from the definition of γ .

2610 Next, I bin the data in $\beta\gamma$. I chose a fine binning so as to capture the different

8.1. dE/dx measurement in the TPC




Figure 8.11: Truncated mean dE/dx obtained for the FHC neutrino sample as a function of the $\beta\gamma$ product (upper panel). Also shown are the fitted most probable values for each $\beta\gamma$ bin (red points) and the best fit obtained using the ALEPH parametrisation (black line). The residuals resulting from the fit are shown in the lower panel.

2611 features of the ionisation curve. Instead of fixing the bin width, I select them so each one
 2612 has approximately the same statistics. Then, for each $\beta\gamma$ slice, I compute the median
 2613 and the interquartile range (IQR) of the $\langle dE/dx \rangle$ distribution. Using these, I make a
 2614 histogram in the range [median – IQR, median + 5 IQR], which I fit to a LanGauss
 2615 function in order to extract the MPV. Using this range accounts for the asymmetric
 2616 nature of the distributions, while also helps avoiding a second, lower maximum present
 2617 at low $\beta\gamma$, probably a result of reconstruction failures.

2618 A few examples of these fits are shown in Fig. 8.9. The chosen values of $\beta\gamma$ sit in
 2619 very distinct points along the $\langle dE/dx \rangle$ curve, going from the high ionisation region at
 2620 low velocities (top left panel), to the minimum point (top right panel), the beginning of
 2621 the relativistic rise (bottom left panel), and the plateau produced by the density effect

Chapter 8. Particle ID in GArSoft




Figure 8.12: Distribution of the 60% truncated mean $\langle dE/dx \rangle$ versus reconstructed momentum for the FHC neutrino sample. The black lines indicate the predictions of the ALEPH parametrisation for electrons, muons, charged pions and protons.

2622 (bottom right panel).

2623 I used the resulting most probable $\langle dE/dx \rangle$ values and the centres of the $\beta\gamma$ bins as
 2624 the points to fit to the ALEPH formula. For this particular fit I used the least-squares
 2625 method to get a first estimation of the ALEPH parameters. Applying some uniform
 2626 priors, I then used these values as the starting point of a 100000 steps MCMC. In
 2627 Fig. 8.10 I show the posterior probability distributions I obtain for each parameter.
 2628 The reported best fit points are based on the 16th, 50th, and 84th percentiles in the
 2629 marginalised distributions.

2630 The resulting fit (black line), compared to the data points (red points) and the
 2631 underlying distribution is shown in Fig. 8.11 (top panel). The overall fit is good, with a
 2632 reduced chi-squared of $\chi^2/ndf = 1.02$. However, there are some regions where the fit
 2633 does not describe the data correctly, like the very low $\beta\gamma$ regime, where the fit severely
 2634 underestimates for energy losses $\gtrsim 50$ keV/cm, and the start of the relativistic raise,
 2635 where we have a slight overestimation. This is a result of those points having a larger

8.2. Muon and pion separation in the ECal and MuID




Figure 8.13: Estimated values of the mean dE/dx bias (left panel) and resolution (right panel) obtained for the true protons in a FHC neutrino sample.

uncertainty when compared to the ones around the dip or the plateau areas. These differences can be better seen in the residual plot, Fig. 8.11 (bottom panel).

8.1.4 Particle identification

8.2 Muon and pion separation in the ECal and MuID

As it could be seen from Fig. 8.12, it is not possible to separate muons and charged pions in the HPgTPC using dE/dx for momenta $\gtrsim 300$ MeV/c. In ND-GAr, approximately 70% of the interactions in FHC mode will be ν_μ CC (compared to the 47% of $\bar{\nu}_\mu$ CC interactions when operating in RHC mode), while 24% are neutral currents. Out of these, around 53% and 47% of them will produce at least one charged pion in the final state, respectively. Fig. 8.14 shows a comparison between the spectra of the primary muons and the charged pions for ν_μ CC interactions in ND-GAr producing one or more charged pions. From this, one can see that (i) the majority of muons and charged pions are not going to be distinguishable with a $\langle dE/dx \rangle$ measurement, and that (ii) particle identification is necessary both to classify correctly the ν_μ CC events and identify the primary muon within them.

ND-GAr counts with two other subdetectors which can provide additional information for this task, namely the ECal and MuID. The current ECal design, described in (ref

Chapter 8. Particle ID in GArSoft




Figure 8.14: True momentum distribution for the primary muon in ν_μ CC $N\pi^\pm$ interactions inside the fiducial volume of ND-GAr (blue line), compared to the post FSI charged pion spectrum (red line).

section), consists of 42 layers, made of 5 mm of Pb, 7 mm of plastic scintillator and a 1 mm PCB board. The total thickness of this calorimeter is 1.66 nuclear interaction lengths or 1.39 pion interaction lengths. The MuID design is in a more conceptual stage, however it is envisioned to feature layers with 10 cm of Fe and 2 cm of plastic scintillator⁵. With its three layers, it will have a thickness of 1.87 or 1.53 nuclear or pion interaction lengths, respectively.

Because pion showers are dominated by inelastic nuclear interactions, the signatures of these particles in the calorimeter will look significantly different from those of muons. Although our ECal is not thick enough to fully contain the hadronic showers of the charged pions at their typical energies in FHC neutrino interactions, they can still be used to understand whether the original particle was more hadron-like or MIP-like. In Fig. 8.15 I have two examples of energy distributions created by a muon (left panel) and a charged pion (right panel) of similar momenta interacting in the ECal. These figures represent the transverse development of the interactions. For each of them, I

⁵It is not mentioned anywhere, but I assume that there should also be another layer of PCB board of 1 mm. However, in this case its contribution to the total thickness of the sampling calorimeter would be negligible.

8.2. Muon and pion separation in the ECal and MuID




Figure 8.15: Distributions of energy deposits in the ECal for a muon (left panel) and a charged pion (right panel) with similar momenta. The energy is projected onto the plane perpendicular to the principal component of the hits, and the positions are relative to the center of the interaction.

2667 computed the principal component and centre of mass of the interaction, projecting
 2668 the position of the hits onto the plane perpendicular to that direction, and taking the
 2669 distances relative to the centre. It can be seen that the muon follows an almost MIP-like
 2670 behaviour, being the central bin in the histogram the one with the highest deposited
 2671 energy. On the other hand, the pion not only deposits more energy overall, but also this
 2672 energy is more spread-out among the different hits. It is this kind of information that
 2673 would allow us to tell apart muons from pions.

2674 This way, I identify three main action points that need to be addressed if one wants
 2675 to use these detectors to distinguish between muons and charged pions. These are:

- 2676 1. the way we make the associations between tracks in the HPgTPC to the activities
 2677 (what in GArSoft we call clusters) in the ECal and the MuID,
- 2678 2. what variables or features one can extract from the calorimeters that encapsulate
 2679 the information we are interested about,
- 2680 3. and how to carry out the classification problem.

Chapter 8. Particle ID in GArSoft

2681 8.2.1 Track-ECal matching

2682 One of the main players in the muon and pion separation is the way we associate clusters
2683 in the ECal to reconstructed tracks in the TPC. Missing some associations or making
2684 wrong ones can bias the ECal quantities that we can use for classifying particles. The
2685 current algorithm in GArSoft provides precise associations, i.e. most of the associations
2686 that it produces are correct, but it appears to miss an important number of associations
2687 (at least when using the default configuration).

2688 The current TPC track-ECal cluster association algorithm is divided in four parts.
2689 It first checks whether the track end point fulfils certain conditions to be extrapolated.
2690 There are two cut values in this step, one for the drift direction and other radial.

2691 If the point can be extrapolated, the code computes the coordinates of the centre
2692 of curvature using the Kalman fit estimates at the track end ($y, z, 1/R, \phi, \tan\lambda$). It
2693 then compares the distance between this and the cluster in the (z, y) plane with R . This
2694 introduces another cut in the perpendicular direction.

2695 The next step is different for clusters in the barrel or in one of the end caps. If it
2696 is a barrel cluster the algorithm extrapolates the track up to the radial distance of the
2697 cluster. There are three possible outcomes, the extrapolated helix can cut the cylinder
2698 of radius r_{clus} two, one or zero times. I get the cut point that is closer to the cluster and
2699 check that it is either in the barrel or the end caps. Computing the difference between
2700 the x coordinates of the cluster and the extrapolated point, the module checks that this
2701 is not greater than a certain cut. If the cluster is in an end cap, I propagate the track
2702 up to the x position of the cluster. Then, the algorithm computes the angle in the (z, y)
2703 plane between the centre of curvature and the cluster, α , and the centre of curvature
2704 and the propagated point, α' . A cut is applied to the quantity $(\alpha - \alpha')R$.

2705 If the cluster contains more than a certain number N of hits, I apply an extra cut to
2706 the dot product of the direction of the track at the propagated x value and the cluster
2707 direction.

2708 The code makes sure to only associate one end of the track (if any) to a cluster.

8.2. Muon and pion separation in the ECal and MuID




Figure 8.16: Left panel: comparison between the precision (blue), sensitivity (yellow) and F_1 score (red) obtained for the default (horizontal lines) and new algorithms, both with the χ^2 -based direction estimator (squares) and cheating the directions (circles), for different values of the χ^2 cut. Right panel: comparison of the performance of the new algorithm when applying the cluster t_0 correction (squares) and when (circles).

2709 However, it can associate more than one track to the same cluster. This makes sense,
2710 as different particles can contribute to the same cluster in the ECal, but it makes it
2711 difficult to quantify the relative contributions of the tracks to a certain cluster.

2712 As a way of comparing the performance of this algorithm, a new, simpler association
2713 module was written. The goal was to have a simple and robust algorithm, which depends
2714 on as few parameters as possible and that can produce a one-to-one matching between
2715 tracks and ECal clusters.

2716 For each reconstructed track, the new algorithms applies the same procedure to the
2717 forward and the backward fits irrespective of their end point positions. It first gets the
2718 Kalman fit parameters at the corresponding end point together with the X position, x_0 ,
2719 (y_0 , z_0 , $1/R$, ϕ_0 , $\tan\lambda$).

2720 For each ECal cluster, I compute the radial distance to the centre of the TPC and
2721 find the ϕ value in the range $[\phi_0, \phi_0 + \text{sign}(R)\phi_{max}]$ that makes the propagated helix
2722 intersect with the circle defined with such radius. The (x, y, z) position of the helix for
2723 the ϕ value found (if any) is then computed. In case there are two intersections, I keep
2724 the one that minimises the distance between (y, z) and (y_c, z_c) .

2725 I then calculate χ^2 value based on the Euclidean distance between the propagated

Chapter 8. Particle ID in GArSoft

2726 point and the cluster:

$$\chi^2/ndf = \frac{\sum_{n=0}^2 (x^{(n)} - x_c^{(n)})^2}{3}. \quad (8.9)$$

2727 If there was no intersection I store a -1 instead. In the end, for each reconstructed track
2728 in the event one ends up with two collections of χ^2 values, one for each ECal cluster
2729 and fit directions.

2730 The current code only supports having ECal clusters associated to one end of each
2731 track. We have two options to decide what track end to keep. The first one tries to
2732 cheat the selection, looking at the distance between the two track ends and the true
2733 start position of the associated MC particle. The second one keeps the track end with
2734 more χ^2 entries below the cut.

2735 This feature of only considering one track end limits the algorithm, making it not
2736 suitable for reconstructing events with particles originating outside the TPC. However,
2737 as for the moment the main concern of the group is the study of neutrino interactions
2738 off the gaseous argon, this is an acceptable assumption.

2739 In order to associate a cluster to a track, I take all clusters with a χ^2 value in the
2740 range $[0, \chi_{cut}^2]$. If a cluster has been assigned to more than one track we leave it with
2741 the one with the lowest χ^2 .

2742 This default behaviour of the algorithm can be modified to associate more than one
2743 track to each cluster. Not only that, but the χ^2 values can be used to assign relative
2744 weights to the different contributions.

2745 To evaluate the performance of the association method, I use a binary classification
2746 approach. In this case, I check the leading MC Track IDs associated to the reconstructed
2747 tracks and ECal clusters. I count an association as true positive (TP) if both Track
2748 IDs coincide. An association is considered false positive (FP) when the Track IDs are
2749 different. If a cluster has not been associated to any track but it shares the Track ID
2750 with a reconstructed track it is counted as a false negative (FN).

2751 For the testing, I used a sample of 10000 FHC neutrino events inside the HPgTPC.
2752 Fig. 8.16 (left panel) shows the precision (blue line), sensitivity (yellow line), and F_1

8.2. Muon and pion separation in the ECal and MuID




Figure 8.17: Schematics of a possible option to deal with track-ECal associations in non-zero t_0 neutrino interaction events, trying to correct for the drift direction uncertainty in a cluster-by-cluster basis using the cluster time, $t_{cluster}$.

2753 score (red line) I obtained for different values of χ^2_{cut} . For comparison, the same metrics
 2754 computed for the default algorithm with the current configuration are also shown (dashed
 2755 lines). In the case of the new algorithm, I used both the χ^2 -based method to estimate
 2756 the track direction described earlier (square markers) and the cheated direction from the
 2757 Geant-level information (circle markers). For either of these we achieve similar values of
 2758 the precision compared to the old code, while having a considerably higher sensitivity.
 2759 It can be seen that cheating the direction of the tracks only makes a difference at high
 2760 χ^2_{cut} , past the optimal value of the cut around the F_1 score maximum. Therefore, I set
 2761 the χ^2 method as the default.

2762 One of the possible weak points of this approach is that it relies on the position along
 2763 the drift direction to make the decisions. Within the current ND-GAr design implemented
 2764 in GArSoft, the timing information is provided by the ECal. That effectively means
 2765 that prior to make the track-ECal associations the reconstructed x positions of the track
 2766 trajectories differ from the simulated ones by an amount:

$$x_{reco}^{(n)} - x_{sim}^{(n)} = v_{drift} t_0, \quad (8.10)$$

2767 where v_{drift} is the mean drift velocity in our medium and the initial time is in the range
 2768 $t_0 \in [0, t_{spill}]$ where t_{spill} is the spill length. For a 10 μs spill this translates into a
 2769 maximum 30 cm uncertainty on the drift direction position.

2770 The current default in GArSoft sets $t_0 = 0$, but the functionality to randomly sample

Chapter 8. Particle ID in GArSoft

2771 this within the spill time is in place. Therefore, we need to understand what is the impact
2772 of a non-zero t_0 on the associations algorithm and foresee possible ways of minimising a
2773 loss in performance.

2774 Fig. 8.17 represents a possible option to tackle the association problem when having
2775 events with a non-zero initial time t_0 . The black and white circles represent the original
2776 points, whereas the squares indicate the corrected positions. The end points of the track
2777 and the propagated points up to the cluster radius are indicated using filled and unfilled
2778 markers respectively. The red square represents the position of the cluster.

2779 Here I try to correct for the drift coordinate position using the time associated to the
2780 cluster. Assuming that the drift time is much larger than the propagation time, $t_{cluster}$
2781 could be used as a good estimation of the t_0 . An alternative can be using the earliest
2782 time associated to a hit in said cluster. Doing this for each cluster before computing
2783 the χ^2 value could be used as an alternative to knowing the specific value of the t_0 , as
2784 when the association is correct this will provide the right correction but its impact is
2785 small enough to not change the position significantly in the case the cluster does not
2786 correspond to a given track.

2787 I tested the effect of this correction again using a sample of 10000 FHC neutrino
2788 events. Fig. 8.16 (right panel) shows the precision (blue line), sensitivity (yellow line),
2789 and F_1 score (red line) for the case the cluster t_0 correction is applied (square markers)
2790 and for the no correction case (circle markers), as a function of χ^2_{cut} . In this case, the
2791 differences are particularly notorious at low values of the cut. It makes sense, as the t_0
2792 effect becomes subdominant when the distance we consider grows large. Overall, the
2793 correction increases the sensitivity while keeping the precision almost unchanged. As a
2794 result, I apply the t_0 correction to the generated samples as the default.

2795 8.2.2 Classification strategy

2796 The problem of the muon and charged pion separation has to be viewed in the broader
2797 context of the particle identification in our detector. Focusing on the beam neutrino
2798 interactions, it is clear that we are going to have muons and pions spanning a broad

8.2. Muon and pion separation in the ECal and MuID




Figure 8.18: Momentum distribution for the reconstructed muons (top panel) and charged pions (bottom panel) in a FHC neutrino sample, together with the fraction of them reaching the ECal (red) and MuID (blue). Each entry corresponds to a reconstructed track, backtracked to a true muon or pion which has not produced any other reconstructed track.

2799 momentum range. Not only that, but we will also have other particles with similar
 2800 characteristics that will make the classification even more challenging. Therefore, we are
 2801 presented with a task that will depend heavily on the kinematic range we are looking at
 2802 each time, as both the available information and the possible impurities of other particle
 2803 species vary.

2804 For instance, distinguishing muons from pions could be difficult at low momenta, as
 2805 a great number of them do not reach the ECal. Therefore, we could think of tailoring a
 2806 version of the classification for that particular case, which could be complemented with
 2807 a dE/dx measurement. Likewise, for momenta $\gtrsim 1$ GeV muons and pions reach the
 2808 calorimeters efficiently, but so do protons. Because of this, one can try to train another
 2809 classifier for this energy range, and rely on other methods to remove as many of the
 2810 protons as possible.

Chapter 8. Particle ID in GArSoft




Figure 8.19: Predicted truncated mean dE/dx versus momentum, for electrons, muons, charged pions and protons, obtained using the ALEPH parametrisation. The vertical dashed lines represent the boundaries of the six regions used for the muon and pion classification training.

Fig. 8.18 shows the momentum distribution of the reconstructed muons (top) and pions (bottom) in a FHC sample. It also contains the fraction of particles reaching the ECal (red) and MuID (blue), for the different momentum bins. In Fig. 8.19 I show the mean dE/dx of different particles as a function of the momentum, computed using the ALEPH parametrisation with the best fit parameters found in Subsec. 8.1.2.

Using these two sources as references, I decided to approach the classification by dividing the problem into six different momentum regions. A summary of these can be found in Tab. 8.2. The basic idea is to exploit all the information that is available in each region and . For the problem at hand, I prepared separated samples of isotropic single muons and pions, with momenta uniformly distributed along the corresponding momentum range. I did not generate samples for the first region, as it is assumed that the separation can be achieved using dE/dx only. For the last region, I generated particles up to a momentum of 10 GeV/c, as that is well above the typical energies of muons and pions from FHC neutrino interactions in ND-GAr.

To tackle this classification problem, I make use of Boosted Decision Trees (BDT). A

8.2. Muon and pion separation in the ECal and MuID

Table 8.2: Momentum ranges and description of the PID approach assumed for the muon and pion classification task.

Momentum range	Description
$< 0.1 \text{ GeV}/c$	All tracks can be separated with dE/dx
$[0.1, 0.3) \text{ GeV}/c$	Use ECal for reaching muons and pions, dE/dx for the rest
$[0.3, 0.8) \text{ GeV}/c$	Use ECal for muons and pions, dE/dx for protons
$[0.8, 1.5) \text{ GeV}/c$	Use ECal and MuID for muons and pions, dE/dx for protons
$[1.5, 3.0) \text{ GeV}/c$	Use ECal and MuID for muons and pions, ToF for protons
$\geq 3.0 \text{ GeV}/c$	Use ECal and MuID for muons and pions, dE/dx and ToF for protons

2826 decision tree uses a flowchart-like structure to make decisions based on some input data.
2827 It starts from a root node, which represents the complete dataset, and then it splits
2828 this based on the variable or feature which gives the best separation between classes,
2829 creating two new nodes. The process repeats for each node until it reaches a certain
2830 limit, like a maximum number of splits or some tolerance criteria. The last set of nodes
2831 are often called leave nodes, and represent the final prediction of the classifier.

2832 Boosting refers to a family of methods to combine the predictions from multiple
2833 classifiers, following a sequential approach where each new model learns from the errors
2834 of the previous one. The process starts with a simple decision tree, which is used to
2835 make predictions on the training data. Then, the data points which the first model got
2836 wrong are assigned higher weights, and another decision tree is trained on the data with
2837 adjusted weights. The predictions of the two trees are then combined, and the cycle
2838 repeats for a predefined number of iterations. Gradient boosting uses the direction of
2839 the steepest error descent to guide the learning process and improve the accuracy with
2840 each iteration.

2841 8.2.3 Feature selection and importance

2842 I am using samples with 100k FHC neutrino interactions, no truth cuts whatsoever.
2843 From the Geant4 I am only keeping the true momentum and PDG. From the track reco
2844 I keep the momentum (end momentum, both forward and backward fits).

Chapter 8. Particle ID in GArSoft

2845 For the ECal, I compute a series of variables per each reconstructed track. As there
2846 can be more than one ECal cluster associated to each track, what I do is collect all
2847 associated clusters and compute these variables from the complete collection of hits
2848 associated to each track. I can roughly divide the variables in three types: energy-related,
2849 geometry-related and statistical.

2850 **• Energy-related ECAL**

- 2851 – ECAL total energy (ClusterTotalEnergy): sum of the energy of all the ECAL
2852 hits.
- 2853 – Hit mean energy (HitMeanEnergy): mean of the hits energy distribution.
- 2854 – Hit std energy (HitStdEnergy): standard deviation of the hits energy distribution.
- 2855 – Hit max energy (HitMaxEnergy): maximum of the hits energy distribution.

2856 **• Geometry-related ECAL**

- 2857 – Mean distance hit-to-cluster (DistHitClusterMean): mean of the distance
2858 distribution between the hits and the corresponding cluster's main axis.
- 2859 – RMS distance hit-to-cluster (DistHitClusterRMS): root mean square of the
2860 distance distribution between the hits and the corresponding cluster's main
2861 axis.
- 2862 – Maximum distance hit-to-centre (DistHitCenterMax): maximum of the
2863 distance distribution between the hits and the centre of the TPC.
- 2864 – Time-of-Flight velocity (TOFVelocity): slope obtained when fitting a straight
2865 line to the hit time versus hit distance to the centre (i.e. $d = v \times t$).

2866 **• Energy and geometry ECAL**

- 2867 – Radius 90% energy (Radius90E): distance in the hit-to-cluster distribution
2868 for which 90% of the total energy is contained in the hits that are closer to
2869 the axis (i.e. radius that contains 90% of the energy).

8.2. Muon and pion separation in the ECal and MuID

• Statistical ECAL

- 2871 – Number of hits (NHits): total number of hits associated to the track.
- 2872 – Number of layers with hits (NLayers): not really a count of all layers with
- 2873 hits but the difference between the last and the first layer with hits.

2874 For the MuID, now that the bug affecting the position of the reco hits has been
2875 solved (07/07/23) I also included some related quantities in the analysis ntuples. They
2876 also fall in the same classification I use for the ECAL variables and are also defined per
2877 reco track as there is also the possibility of having more than one MuID cluster for each
2878 track.

• Energy-related MuID

- 2880 – MuID total energy (ClusterMuIDTotalEnergy): sum of the energy of all the
- 2881 MuID hits.
- 2882 – Hit MuID mean energy (HitMuIDMeanEnergy): mean of the MuID hits
- 2883 energy distribution.
- 2884 – Hit MuID std energy (HitMuIDStdEnergy): standard deviation of the MuID
- 2885 hits energy distribution.
- 2886 – Hit MuID max energy (HitMuIDMaxEnergy): maximum of the MuID hits
- 2887 energy distribution.

• Geometry-related MuID

- 2889 – Max distance MuID hits (DistHitMuIDMax): maximum distance between
- 2890 pairs of MuID hits (not sure this is a good variable, distribution looks nuts).
- 2891 – Maximum distance MuID hit-to-centre (DistHitCenterMuIDMax): maximum
- 2892 of the distance distribution between the MuID hits and the centre of the
- 2893 TPC.

• Statistical MuID

Chapter 8. Particle ID in GArSoft

- 2895 – Number of hits (NHitsMuID): total number of MuID hits associated to the
- 2896 track.
- 2897 – Number of layers with hits (NLayersMuID): not really a count of all layers
- 2898 with MuID hits but the difference between the last and the first layer with
- 2899 MuIDhits.

2900 8.2.4 Hyperparameter optimisation

2901 8.2.5 Probability calibration

2902 8.2.6 Performance

2903 8.3 ECal time-of-flight

2904 8.3.1 Arrival time estimations

2905 8.3.2 Proton and pion separation

2906 8.4 Charged pion decay in flight

2907 As discussed previously, in GArSoft the TPC tracks are formed after a pattern recognition
2908 algorithm and a Kalman filter are applied to the TPC clusters. These two steps can
2909 find discontinuities in the track candidates (e.g. due to a particle decay) when these
2910 so-called breakpoints are large enough. However, for some, more subtle, cases they may
2911 miss them and form a single reconstructed track. It has been noted in the literature
2912 that Kalman filters offer, as a by-product, additional information to form test statistics
2913 to identify these breakpoints [?, ?].

2914 Considering the mean life of the charged pion, $\tau = (2.6033 \pm 0.0005) \times 10^{-8}$ s, one
2915 can estimate that about 12% of the pions with momentum $p \sim \mathcal{O}(500 \text{ MeV})$ (roughly
2916 the peak of the pion momentum distribution in ν_μ CC interactions off argon) decay
2917 inside the TPC. Fig. 8.20 (left panel) shows the amount of charged pions decaying in the
2918 full TPC and fiducial volumes from an isotropic, monoenergetic sample of 10^5 negatively

8.4. Charged pion decay in flight




Figure 8.20: Left panel: number of non-decaying, decaying and decaying in the fiducial volume pions for a MC sample of 10^5 , $p = 500$ MeV isotropic positively charged pions inside the TPC. Right panel: event display for a positive pion decaying inside the fiducial volume, with a single reconstructed track for the pion and muon system.

2919 charged pions with $p = 500$ MeV. We see that about 10% of those decayed, with more
2920 than half of them decaying inside the TPC fiducial volume.

2921 Fig. 8.20 (right panel) shows an example event display of a charged pion (magenta
2922 line) decays in flight inside the TPC, but because the angle of the muon (blue line) is
2923 small both were reconstructed as one single track (black line). In this case, the composite
2924 track reaches the ECal, where it undergoes a muon-like interaction, thus being classified
2925 as a muon.

2926 A way to understand what decaying pion tracks were totally or partially reconstructed
2927 together with the daughter muon is looking at the relative energy contributions to the
2928 reconstructed track. In order to select a sample of such events, I require that a minimum
2929 50% of the total energy comes from the pion and at least 20% from the muon.

2930 To identify potential decays we can use the information we obtain from the Kalman
2931 filter at each step of the fitted track. The simplest test we can think about is computing
2932 the χ^2 of the mismatch between all the parameters in the forward and the backward fits:

$$\chi_k^{2 (FB)} = (\hat{x}_k^B - \hat{x}_k^F)^T [V^{(\hat{x}_k, B)} + V^{(\hat{x}_k, F)}]^{-1} (\hat{x}_k^B - \hat{x}_k^F), \quad (8.11)$$

2933 where \hat{x}_k^F , \hat{x}_k^B are the Kalman filter state vector estimates at step k in the forward and
2934 backward fits and $V^{(\hat{x}_k, F)}$, $V^{(\hat{x}_k, B)}$ the covariance matrices of \hat{x}_k^F and \hat{x}_k^B respectively.

Chapter 8. Particle ID in GArSoft




Figure 8.21: Values of $\chi_k^2(FB)$ (top left panel), F_k (top right panel), $D_k^{1/R}$ (bottom left panel) and D_k^ϕ (bottom right panel) versus position along the drift direction for a reconstructed track in a positive pion decay event. The vertical red dashed line indicates the true location of the decay point.

- 2935 Using the values of the χ^2 at measurement k for the forward and backward fits we can
 2936 compute another χ^2 value that characterises the overall track fit:

$$\chi_{track}^2 = \chi_k^2(F) + \chi_k^2(B) + \chi_k^2(FB), \quad (8.12)$$

- 2937 which remains approximately constant for all k .

2938 An alternative approach proposed in the context of the NOMAD experiment was
 2939 using a fit with a more elaborate breakpoint hypothesis, so we can perform a comparison
 2940 of the χ^2 with and without breakpoints. This can be achieved by using some alternative
 2941 parametrisation with extra parameters, which allows some of the track parameters to
 2942 be discontinuous at certain points. A decay changes the momentum magnitude and
 2943 direction, so we can use the new state vector:

$$\alpha = \left(y, z, 1/R_F, 1/R_B, \phi_F, \phi_B, \tan\lambda_F, \tan\lambda_B \right)^T. \quad (8.13)$$

8.4. Charged pion decay in flight

2944 As we already have the estimates from the standard Kalman filter and their
2945 covariance matrices at each point, we do not need to repeat the Kalman fit for the new
2946 parametrisation. Instead, I can compute the values of α at each point k that minimise
2947 the χ^2 resulting from comparing them to $\{\hat{x}_k^B, \hat{x}_k^F\}$. Introducing the two 5×8 matrices:

$$H^F = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad H^B = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (8.14)$$

2948 we can write this as:

$$\begin{aligned} \chi_k^{2(FB)}(\alpha) &= (\hat{x}_k^F - H^F \alpha)^T \left[V^{(\hat{x}_k, F)} \right]^{-1} (\hat{x}_k^F - H^F \alpha) \\ &\quad + (\hat{x}_k^B - H^B \alpha)^T \left[V^{(\hat{x}_k, B)} \right]^{-1} (\hat{x}_k^B - H^B \alpha). \end{aligned} \quad (8.15)$$

2949 The minimum of $\chi_k^{2(FB)}(\alpha)$ is found when the measured new state vector takes the
2950 value:

$$\hat{\alpha}_k = V^{(\hat{\alpha}_k)} H^T (V^{(\hat{x}_k)})^{-1} \hat{X}, \quad (8.16)$$

2951 where $\hat{X} = \{\hat{x}_k^B, \hat{x}_k^F\}$, $V^{(\hat{x}_k)}$ is the block diagonal matrix formed by $V^{(\hat{x}_k, F)}$ and $V^{(\hat{x}_k, B)}$
2952 and $V^{(\hat{\alpha}_k)}$ is the covariance matrix of $\hat{\alpha}_k$, given by:

$$V^{(\hat{\alpha}_k)} = \left(H^T (V^{(\hat{x}_k)})^{-1} H \right)^{-1}. \quad (8.17)$$

2953 From these new fit estimates we can compute the F statistic, which tells us whether
2954 the model with breakpoint provides a statistically significant better fit:

$$F_k = \left(\frac{\chi_{track,k}^2 - \chi_{full,k}^2}{8 - 5} \right) / \left(\frac{\chi_{full,k}^2}{N - 8} \right). \quad (8.18)$$

2955 One can also compute the signed difference of the duplicated variables divided by

Chapter 8. Particle ID in GArSoft




Figure 8.22: Fractional residual distributions of the true and reconstructed decay position along the drift coordinate, using the position of the maximum of $\chi_k^{2 (\text{FB})}$ (left panel) and F_k (right panel) as estimates of the decay position. Also shown are double Gaussian fits to these points (red lines).

2956 their standard deviation at each point. These represent how significant the discontinuity
2957 in each variable is. For any variable η we can write it as:

$$D_k^\eta = \frac{\hat{\eta}_k^B - \hat{\eta}_k^F}{\sqrt{\text{Var}[\hat{\eta}_k^F] + \text{Var}[\hat{\eta}_k^B] - 2\text{Cov}[\hat{\eta}_k^F, \hat{\eta}_k^B]}}, \quad (8.19)$$

2958 In our case, the relevant ones to look at are $D_k^{1/R}$ and D_k^ϕ .

2959 Fig. 8.21 shows the values of $\chi_k^{2 (\text{FB})}$, F_k , $D_k^{1/R}$ and D_k^ϕ as functions of the position
2960 along the drift direction, for an example reconstructed track with 55.5% of the energy
2961 coming from the charged pion and 45.5% from the daughter muon. The true position of
2962 the decay is indicated (dashed red lines). Notice how $\chi_k^{2 (\text{FB})}$ and F_k , $D_k^{1/R}$ reach their
2963 maxima near the decay point. In the former case this indicates a large forward-backward
2964 difference in the track fit. In the later it represents that the extended state vector
2965 improves the fit particularly around that point.

2966 I can estimate the decay position finding resolution by computing the difference
2967 between the X position of the maxima of $\chi_k^{2 (\text{FB})}$ and F_k and the X position of the
2968 true decay. Fig. 8.22 represent the the fractional residual distributions for both cases,
2969 from the sample of tracks containing pion decays. Fitting a double Gaussian to the
2970 distributions (red lines) I find a resolution of 13.62% and 7.45% respectively.

8.4. Charged pion decay in flight




Figure 8.23: Distributions of the extreme values of $\chi_k^2(FB)$ (top left panel), F_k (top right panel), $D_k^{1/R}$ (bottom left panel) and D_k^ϕ (bottom right panel) for non-decaying reconstructed pion tracks (blue) and tracks which include the decay inside the fiducial volume (red).

2971 In principle, the F -statistic should follow a Fisher distribution with $(8 - 5)$ and
 2972 $(N - 8)$ degrees of freedom under the null hypothesis. In most of our cases $N \sim \mathcal{O}(100)$,
 2973 so the probability density functions will look very similar. In this case, it is safe to take
 2974 the limit $N \rightarrow \infty$ in the Fisher PDF:

$$\begin{aligned}\tilde{f}(x; a - b) &= \lim_{N \rightarrow \infty} f(x; a - b, N - a) \\ &= \frac{2^{-\frac{a-b}{2}}}{\Gamma(\frac{a-b}{2})} (a - b)^{\frac{a-b}{2}} x^{\frac{a-b}{2}-1} e^{-\frac{a-b}{2}x}.\end{aligned}\quad (8.20)$$

2975 In our case $a - b = 8 - 5 = 3$, so we would obtain a p-value of 0.05 at $x = 2.60$.

2976 Fig. 8.23 contains the distributions of the maxima of $\chi_k^2(FB)$, F_k and D_k^ϕ and the
 2977 minima of $D_k^{1/R}$ for a sample of non-decaying pion tracks (blue) and another sample of
 2978 reconstructed tracks containing part of the pion and the daughter muon from a decay
 2979 inside the fiducial volume (red). Notice that, even though the values of $F_k^{(max)}$ for the
 2980 decay sample are typically larger than for the non-decaying one, just a small fraction of

Chapter 8. Particle ID in GArSoft




Figure 8.24: Left panel: distributions of the predicted probabilities assigned by the BDT classifier to a test sample of decaying pion+muon tracks (blue) and non-decaying pion tracks (red). Left: signal efficiency versus background acceptance (ROC curve) obtained from the BDT for the test sample.




Figure 8.25: Left panel: dependence of the decay position finding resolution on the true value of the decay angle for the $\chi_k^2(FB)$ (red) and F_k (blue) methods. Right panel: signal efficiency (blue line) and background rejection (red line) from the BDT classifier versus true decay angle.

2981 the events go beyond the aforementioned value of $F = 2.60$. Therefore, from a practical
2982 point of view, it is not the most efficient variable to use for selecting the decay events.

2983 However, looking at the $D_k^{1/R \text{ (min)}}$ distribution we can see there is a big difference
2984 between non-decaying and decaying events in this variable. One can use a combination
2985 of these four variables to distinguish between the pion decay events (signal) and the
2986 non-decaying pions (background).

2987 An approach to this classification could be using a boosted decision tree (BDT). One
2988 of the advantages of BDTs is that they are easy to interpret and identify the relative
2989 importance of the different input variables. Training a BDT with 400 estimators and a

8.5. Neutral particle identification

maximum depth of 4 I can obtain an efficient classification without overtraining. Fig. 8.24 (left panel) shows the distribution of probabilities predicted by the BDT for a test sample. The signal efficiency as a function of background acceptance, the so-called ROC curve, is shown in Fig. 8.24 (right panel). With a relative importance of 0.83, the most important variable turned out to be $D_k^{1/R \text{ (min)}}$.

One thing we can check is how the resolution to the decay and the signal efficiency in the classification changes with the true decay angle. Using an equal-frequency binning for the decay angles, we can repeat the previous steps for each bin.

Fig. 8.25 (left panel) shows the dependence on the decay angle of the decay finding resolution. We can see that for the $\chi_k^2 \text{ (FB)}$ maximum location method the resolution consistently lies between 12 to 16%. However, the $F_k^{(max)}$ approach gives a significantly better resolution for high angle values, reaching the 4 – 6% range for decay angles $\geq 4^\circ$.

For the classification dependence on the angle, I use the same classifier I trained before but evaluating the test sample for each individual angular bin. I compute the signal efficiency in each bin for a fixed value of the background rejection, in this case 90%. Similarly, for the background rejection estimation I use a fixed signal efficiency value of 90%. Fig. 8.25 (right panel) represents the change in signal efficiency (blue) and background rejection (red) with the value of the true decay angles.

8.4.1 Track breakpoints

8.5 Neutral particle identification

8.5.1 ECal clustering

Another important reconstruction item is the clustering algorithm of ECal hits in GArSoft. The default module features a NN algorithm that treats all hits in the same way, independently of the layer each hit comes from. However, the current ECal design of ND-GAr has two very different types of scintillator layers. The inner layers are made out of tiles, which provide excellent angular and timing resolutions. On the other hand, the outer layers are cross scintillator strips. That way, an algorithm that treats hits

Chapter 8. Particle ID in GArSoft

3017 from both kinds of layers differently may be able to improve the current performance.

3018 Inspired by the reconstruction of T2K’s ND280 downstream ECal [?], the idea was
3019 to put together a clustering module that first builds clusters for the different ECal views
3020 (tiles, strips segmented in the X direction and strips segmented in Y direction), and
3021 then tries to match them together to form the final clusters.

3022 Working on a module-by-module basis, the algorithm first separates the hits depending
3023 on the layer type they come from. Then, it performs a NN clustering for the 3 sets of
3024 hits separately. For the tile hits it clusters together all the hits which are in nearest-
3025 neighbouring tiles and nearest-neighbouring layers, for strip hits it looks at nearest-
3026 neighbouring strips and next-to-nearest-neighbouring layers (as the layers with strips
3027 along the two directions are alternated). For strip clusters an additional cut in the
3028 direction along the strip length is needed.

3029 After this first clustering I then apply a recursive re-clustering for each collection
3030 of strip clusters based on a PCA method. In each case, we loop over the clusters with
3031 $N_{hits} \geq 2$, computing the centre of mass and three principal components. Propagating
3032 these axes up to the layers of the rest of the clusters, we check if the propagated point
3033 and the centre of mass of the second cluster are within next-to-nearest-neighbouring
3034 strips. An additional cut in the direction along the strip length is also needed. Moreover,
3035 I require that the two closest hits across the two clusters are at most in next-to-nearest-
3036 neighbouring strips. I merge the clusters if these three conditions are satisfied. The
3037 re-clustering is repeated until no more cluster pairs pass the cuts.

3038 The clusters in each strip view are combined if their centres of mass are close enough
3039 and they point in the same direction. An alternative approach for the strip cluster
3040 merging could be to compute the overlap between the ellipsoids defined by the principal
3041 axes of the clusters, and then merge the pair if the overlap exceeds some threshold.
3042 Further study is needed to understand if this change would have an impact in the overall
3043 clustering performance.

3044 To merge the tile clusters to the combined strip clusters I propagate the principal
3045 axis of the strip cluster towards the inner layers, up to the centre of mass layer of the

8.5. Neutral particle identification




Figure 8.26: Mean values of the F_1 -score marginal distributions for the different free parameters of the new clustering algorithm, with the error bars representing one standard deviation around the mean. The F_1 -score values were computed for the 6561 possible parameter configurations using 1000 ν_μ CC interaction events.

3046 tile cluster. I merge the clusters if the distance between the propagated point and the
 3047 centre of mass is bellow a certain cut.

3048 The last step is to check if clusters in neighbouring modules should be merged
 3049 together, both across two barrel modules, across end cap modules and between barrel
 3050 end cap modules. I check the distance between the two closest hits in the pair of clusters
 3051 and merge them if it passes this and an additional direction cut.

3052 Fig. ?? presents an example of the clustering steps relevant for strip layer hits, from
 3053 the input hits (top left panel) to the NN clustering (top right panel) and re-clustering
 3054 (bottom left panel) for each strip view and the final merging strip clusters (bottom
 3055 right panel). It shows the hits from a single ECal barrel module in a ν_μ CC interaction
 3056 event with a neutral pion and a proton in the final state. The two clusters on the left
 3057 correspond to the photon pair from the π^0 decay and the one on the upper right corner
 3058 is associated to the proton.

3059 This algorithm has a total number of eight free parameters that need to be optimised.
 3060 I used a sample of 1000 ν_μ CC interactions in order to obtain the optimal configuration of
 3061 clustering parameters. This sample was generated up to the default ECal hit clustering

Chapter 8. Particle ID in GArSoft

Table 8.3: Summary of parameters and sampled values used in the optimisation of the clustering algorithm.

Name	Units	Sampled values	Description
PrimMinorCut	strips	2, 3, 4	Distance along strip length in NN clustering
RecMinorCut	strips	3, 5, 8	Distance between propagated point and CM along strip length in re-clustering
RecPerpCut	strips	2, 3, 4	Closest hit pair distance in re-clustering
StripDistCut	strips	3, 5, 8	Distance between CMs in strip cluster merging
StripDirCut	cos	0.5, 0.7, 0.9	Main axes direction cut in strip cluster merging
PointDistCut	tiles	3, 5, 8	Distance between propagated point and CM in strip-tile matching
MergePerpCut	cm	10, 20, 30	Closest hit pair distance in module merging
MergeDirCut	cos	0.5, 0.7, 0.9	Main axes direction cut in module merging

3062 level, so then I could run the new clustering algorithm each time with a different
 3063 configuration of parameters. As the number of parameters is relatively large, I only
 3064 performed a coarse-grained scan of the parameter space. Sampling each of the eight
 3065 parameters at three different points each I obtain 6561 different configurations. These
 3066 parameters, together with the used values, are summarised in Tab. 8.3.

3067 In order to measure the performance of the clustering, I use a binary classification
 3068 approach. For each formed cluster, I identify the Geant4 Track ID of the matching MC
 3069 particle and the energy fraction of each hit. Then, I assign to each cluster the Track ID
 3070 with the highest total energy fraction. For each of the different Track IDs associated to
 3071 the clusters, I select the cluster with the highest energy (only from the hits with the
 3072 same Track ID). I identify such a cluster as the main cluster for that Track ID. I count
 3073 as true positives (TPs) the hits with the correct Track ID in each main cluster. False
 3074 positives (FPs) are the hits with the incorrect Track ID for the cluster they are in, not
 3075 only main clusters. The false negatives (FNs) are the hits with the correct Track ID in
 3076 clusters other than the main.

3077 Fig. 8.26 shows the computed F_1 -score values for the different cuts. In each case, the
 3078 central value represents the mean of the F_1 -score distribution for the specified value of

8.5. Neutral particle identification




Figure 8.27: Left panel: distributions of the number of ECal clusters per photon from π^0 decays for the standard (red) and new (blue) clustering algorithms. Right panel: reconstructed invariant mass distributions for photon pairs from single π^0 events using the standard (red) and new (blue) ECal clustering algorithms.

3079 the corresponding variable and the vertical error bar represents one standard deviation
 3080 around the mean. Also shown are the Pearson correlation coefficients of these central
 3081 values. We can see that five of the variables have a sizeable effect on the F_1 -score, with
 3082 an absolute difference between the last and first values as big as 4%.

3083 The working configuration is obtained as follows. I first select all configurations
 3084 with purity $\geq 90\%$. Among those, I choose the combinations that yield the maximum
 3085 F_1 -score. If more than one configuration remains I select the one with the highest
 3086 sensitivity. Doing so, I end up with a parameter configuration with an efficiency of 88%
 3087 and a 90% purity. Compared with the default algorithm, which gives an efficiency of
 3088 76% and a purity of 91% for the same sample, I have managed to improve the efficiency
 3089 by a factor of 1.16.

3090 8.5.2 π^0 reconstruction

3091 One of the potential applications of the new ECal hit clustering is the reconstruction of
 3092 neutral particles, in particular pions. Neutral pions decay promptly after being produced,
 3093 through the $\pi^0 \rightarrow \gamma\gamma$ channel ($98.823 \pm 0.034\%$) of the time. The photon pair does
 3094 not leave any traces in the HPgTPC (unless one or both of them converts into an
 3095 electron-positron pair), but each of them will produce an electromagnetic shower in

Chapter 8. Particle ID in GArSoft

3096 the ECal.

3097 To test the potential impact of the new algorithm in π^0 reconstruction, I generated
3098 a MC sample of single, isotropic neutral pions inside the HPgTPC. All pions were
3099 generated with a momentum $p = 500$ MeV and their initial positions were uniformly
3100 sampled inside a $2 \times 2 \times 2$ m box aligned with the centre of the TPC. I ran both the
3101 default and the new clustering algorithms, using for the latter the optimised configuration
3102 discussed above.

3103 The first thing to notice is that the number of clusters produced per photon has
3104 decreased. Fig. 8.27 (left panel) shows these distributions for the default (red) and new
3105 (blue) algorithms. Using a simple Gaussian fit, we see that the mean number of ECal
3106 clusters per photon went from 1.82 ± 0.01 to 1.09 ± 0.03 . This effectively means that
3107 with the new algorithm the ECal activity of one true particle is typically reconstructed
3108 as a single object. From the reconstruction point of view this can be an advantage. As
3109 now most of the photon energy ends up in a single ECal cluster, I can simply use cluster
3110 pairs to identify the π^0 decay.

3111 In general, one calculates the invariant mass of the photon pair as:

$$m_{\gamma\gamma} = \sqrt{2E_1E_2(1 - \cos \theta)}, \quad (8.21)$$

3112 where E_i are the energies of the photons and θ the opening angle between them. In this
3113 case I can use the energies deposited in the ECal and their incident directions. This
3114 quantity is computed for all possible pairs of clusters, using their position together with
3115 the true decay point. In a more realistic scenario, e.g. ν_μ CC interaction, one could use
3116 the position of the reconstructed primary vertex instead. I also tried to use the principal
3117 direction of the clusters, but that approach gave considerably worse results. For each
3118 event I only keep the pair with an invariant mass closer to the true π^0 mass value.

3119 Fig. 8.27 (right panel) shows the invariant mass distributions for the photon pairs
3120 we get using the default (red) and the new (blue) ECal clustering algorithms. For the fit
3121 I used a modified version of the Crystal Ball function [?], obtained by taking the limit

8.5. Neutral particle identification

3122 where the parameter controlling the power-law tail goes to infinity:

$$f(x; N, \mu, \sigma, \alpha) = N \cdot \begin{cases} e^{\frac{\alpha(2x-2\mu+\alpha\sigma)}{2\sigma}}; & x \leq \mu - \alpha\sigma, \\ e^{-\frac{(x-\mu)^2}{2\sigma^2}}; & x > \mu - \alpha\sigma. \end{cases} \quad (8.22)$$

3123 Comparing the fitted mean and standard deviation values for the Gaussian cores, we
3124 see that the distribution for the new algorithm is a 67% narrower and also peaks much
3125 closer to the true m_{π^0} value, going from 101.3 ± 0.4 MeV to 130.8 ± 0.6 MeV.

3126 Chapter 9

3127 Event selection in ND-GAr

3128 9.1 CAFs and CAFAna

3129 9.2 Event selection

3130 9.2.1 ν_μ CC selection

3131 9.2.2 Charged pion multiplicity

³¹³² Chapter 10

³¹³³ Conclusions

³¹³⁴ Appendix A

³¹³⁵ An appendix

₃₁₃₆ Bibliography

- ₃₁₃₇ [1] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
₃₁₃₈ *Detector Technical Design Report, Volume I Introduction to DUNE*, JINST **15**
₃₁₃₉ (2020) T08008 [2002.02967]. 15, 43, 44, 45, 48, 49, 50, 56, 57, 58, 65, 66
- ₃₁₄₀ [2] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
₃₁₄₁ *Detector Technical Design Report, Volume IV: Far Detector Single-phase*
₃₁₄₂ *Technology*, JINST **15** (2020) T08010 [2002.03010]. 15, 53
- ₃₁₄₃ [3] J.N. Bahcall, A.M. Serenelli and S. Basu, *New solar opacities, abundances,*
₃₁₄₄ *helioseismology, and neutrino fluxes*, *Astrophys. J. Lett.* **621** (2005) L85
₃₁₄₅ [[astro-ph/0412440](#)]. 21, 113, 115
- ₃₁₄₆ [4] R. Garani and S. Palomares-Ruiz, *Dark matter in the Sun: scattering off electrons*
₃₁₄₇ *vs nucleons*, JCAP **05** (2017) 007 [[1702.02768](#)]. 21, 111, 112, 114, 115, 116, 139
- ₃₁₄₈ [5] M. Honda, M. Sajjad Athar, T. Kajita, K. Kasahara and S. Midorikawa,
₃₁₄₉ *Atmospheric neutrino flux calculation using the NRLMSISE-00 atmospheric model*,
₃₁₅₀ *Phys. Rev. D* **92** (2015) 023004 [[1502.03916](#)]. 22, 119, 120
- ₃₁₅₁ [6] M. Colom i Bernadich and C. Pérez de los Heros, *Limits on Kaluza-Klein dark*
₃₁₅₂ *matter annihilation in the Sun from recent IceCube results*, Eur. Phys. J. C, **80** 2
₃₁₅₃ (2020) 129 **80** (2019) [[1912.04585](#)]. 22, 124
- ₃₁₅₄ [7] ANTARES collaboration, *Search for Dark Matter in the Sun with the ANTARES*
₃₁₅₅ *Neutrino Telescope in the CMSSM and mUED frameworks*, Nucl. Instrum. Meth. A
₃₁₅₆ **725** (2013) 76 [[1204.5290](#)]. 22, 124

BIBLIOGRAPHY

- 3157 [8] N. Deutschmann, T. Flacke and J.S. Kim, *Current LHC Constraints on Minimal*
3158 *Universal Extra Dimensions*, *Phys. Lett. B* **771** (2017) 515 [[1702.00410](#)]. 22, 124,
3159 125
- 3160 [9] ICECUBE collaboration, *Search for GeV-scale dark matter annihilation in the Sun*
3161 *with IceCube DeepCore*, *Phys. Rev. D* **105** (2022) 062004 [[2111.09970](#)]. 24, 136,
3162 137
- 3163 [10] C.-S. Chen, F.-F. Lee, G.-L. Lin and Y.-H. Lin, *Probing Dark Matter*
3164 *Self-Interaction in the Sun with IceCube-PINGU*, *JCAP* **10** (2014) 049 [[1408.5471](#)].
3165 24, 136, 137
- 3166 [11] N.F. Bell, M.J. Dolan and S. Robles, *Searching for dark matter in the Sun using*
3167 *Hyper-Kamiokande*, *JCAP* **11** (2021) 004 [[2107.04216](#)]. 24, 136, 137
- 3168 [12] E. Behnke et al., *Final Results of the PICASSO Dark Matter Search Experiment*,
3169 *Astropart. Phys.* **90** (2017) 85 [[1611.01499](#)]. 24, 136, 137
- 3170 [13] PICO collaboration, *Dark Matter Search Results from the Complete Exposure of*
3171 *the PICO-60 C₃F₈ Bubble Chamber*, *Phys. Rev. D* **100** (2019) 022001
3172 [[1902.04031](#)]. 24, 136, 137
- 3173 [14] DARKSIDE collaboration, *Constraints on Sub-GeV Dark-Matter–Electron*
3174 *Scattering from the DarkSide-50 Experiment*, *Phys. Rev. Lett.* **121** (2018) 111303
3175 [[1802.06998](#)]. 24, 140, 141
- 3176 [15] XENON collaboration, *Light Dark Matter Search with Ionization Signals in*
3177 *XENON1T*, *Phys. Rev. Lett.* **123** (2019) 251801 [[1907.11485](#)]. 24, 140, 141
- 3178 [16] P.F. de Salas, D.V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C.A. Ternes
3179 et al., *2020 global reassessment of the neutrino oscillation picture*, *JHEP* **02** (2021)
3180 071 [[2006.11237](#)]. 29, 40, 45
- 3181 [17] W. Pauli, *Dear radioactive ladies and gentlemen*, *Phys. Today* **31N9** (1978) 27. 35

BIBLIOGRAPHY

- 3182 [18] F. Reines and C.L. Cowan, *Detection of the free neutrino*, *Phys. Rev.* **92** (1953)
3183 830. 35
- 3184 [19] ALEPH, DELPHI, L3, OPAL, SLD, LEP ELECTROWEAK WORKING GROUP,
3185 SLD ELECTROWEAK GROUP, SLD HEAVY FLAVOUR GROUP collaboration,
3186 *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006)
3187 257 [[hep-ex/0509008](#)]. 36
- 3188 [20] SUPER-KAMIOKANDE collaboration, *Evidence for oscillation of atmospheric*
3189 *neutrinos*, *Phys. Rev. Lett.* **81** (1998) 1562 [[hep-ex/9807003](#)]. 36
- 3190 [21] B. Pontecorvo, *Inverse beta processes and nonconservation of lepton charge*, *Zh.*
3191 *Eksp. Teor. Fiz.* **34** (1957) 247. 36
- 3192 [22] Z. Maki, M. Nakagawa and S. Sakata, *Remarks on the unified model of elementary*
3193 *particles*, *Prog. Theor. Phys.* **28** (1962) 870. 36
- 3194 [23] L. Wolfenstein, *Neutrino Oscillations in Matter*, *Phys. Rev. D* **17** (1978) 2369. 38
- 3195 [24] B.T. Cleveland, T. Daily, R. Davis, Jr., J.R. Distel, K. Lande, C.K. Lee et al.,
3196 *Measurement of the solar electron neutrino flux with the Homestake chlorine*
3197 *detector*, *Astrophys. J.* **496** (1998) 505. 39
- 3198 [25] F. Kaether, W. Hampel, G. Heusser, J. Kiko and T. Kirsten, *Reanalysis of the*
3199 *GALLEX solar neutrino flux and source experiments*, *Phys. Lett. B* **685** (2010) 47
3200 [[1001.2731](#)]. 39
- 3201 [26] SAGE collaboration, *Measurement of the solar neutrino capture rate with gallium*
3202 *metal. III: Results for the 2002–2007 data-taking period*, *Phys. Rev. C* **80** (2009)
3203 015807 [[0901.2200](#)]. 39
- 3204 [27] G. Bellini et al., *Precision measurement of the 7Be solar neutrino interaction rate*
3205 *in Borexino*, *Phys. Rev. Lett.* **107** (2011) 141302 [[1104.1816](#)]. 39

BIBLIOGRAPHY

- 3206 [28] SUPER-KAMIOKANDE collaboration, *Solar neutrino measurements in*
3207 *super-Kamiokande-I*, *Phys. Rev. D* **73** (2006) 112001 [[hep-ex/0508053](#)]. 39
- 3208 [29] SNO collaboration, *Combined Analysis of all Three Phases of Solar Neutrino Data*
3209 *from the Sudbury Neutrino Observatory*, *Phys. Rev. C* **88** (2013) 025501
3210 [[1109.0763](#)]. 39
- 3211 [30] SUPER-KAMIOKANDE collaboration, *Atmospheric neutrino oscillation analysis with*
3212 *external constraints in Super-Kamiokande I-IV*, *Phys. Rev. D* **97** (2018) 072001
3213 [[1710.09126](#)]. 39
- 3214 [31] ICECUBE collaboration, *Measurement of Atmospheric Neutrino Oscillations at*
3215 *6–56 GeV with IceCube DeepCore*, *Phys. Rev. Lett.* **120** (2018) 071801
3216 [[1707.07081](#)]. 39
- 3217 [32] KAMLAND collaboration, *Reactor On-Off Antineutrino Measurement with*
3218 *KamLAND*, *Phys. Rev. D* **88** (2013) 033001 [[1303.4667](#)]. 39
- 3219 [33] RENO collaboration, *Measurement of Reactor Antineutrino Oscillation Amplitude*
3220 *and Frequency at RENO*, *Phys. Rev. Lett.* **121** (2018) 201801 [[1806.00248](#)]. 39
- 3221 [34] DAYA BAY collaboration, *Measurement of the Electron Antineutrino Oscillation*
3222 *with 1958 Days of Operation at Daya Bay*, *Phys. Rev. Lett.* **121** (2018) 241805
3223 [[1809.02261](#)]. 39
- 3224 [35] K.J. Kelly, P.A.N. Machado, S.J. Parke, Y.F. Perez-Gonzalez and R.Z. Funchal,
3225 *Neutrino mass ordering in light of recent data*, *Phys. Rev. D* **103** (2021) 013004. 40
- 3226 [36] P. Dunne, “Latest Neutrino Oscillation Results from T2K.” Jul, 2020. 40
- 3227 [37] MINOS collaboration, *Combined analysis of ν_μ disappearance and $\nu_\mu \rightarrow \nu_e$*
3228 *appearance in MINOS using accelerator and atmospheric neutrinos*, *Phys. Rev. Lett.*
3229 **112** (2014) 191801 [[1403.0867](#)]. 40

BIBLIOGRAPHY

- 3230 [38] K2K collaboration, *Measurement of Neutrino Oscillation by the K2K Experiment*,
3231 *Phys. Rev. D* **74** (2006) 072003 [[hep-ex/0606032](#)]. 40
- 3232 [39] DUNE collaboration, *Long-baseline neutrino oscillation physics potential of the*
3233 *DUNE experiment*, *Eur. Phys. J. C* **80** (2020) 978 [[2006.16043](#)]. 40
- 3234 [40] HYPER-KAMIOKANDE collaboration, *Physics potential of Hyper-Kamiokande for*
3235 *neutrino oscillation measurements*, *PoS NuFact2019* (2019) 040. 40
- 3236 [41] DUNE collaboration, *Snowmass Neutrino Frontier: DUNE Physics Summary*,
3237 [2203.06100](#). 45, 46
- 3238 [42] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
3239 *Detector Technical Design Report, Volume II: DUNE Physics*, [2002.03005](#). 45, 47,
3240 61, 63, 119, 128, 132
- 3241 [43] SUPER-KAMIOKANDE collaboration, *Search for Proton Decay via $p \rightarrow e^+ \pi_0$ and*
3242 *$p \rightarrow \mu^+ \pi_0$ in a Large Water Cherenkov Detector*, *Phys. Rev. Lett.* **102** (2009)
3243 [141801](#) [[0903.0676](#)]. 46
- 3244 [44] S. Raby, *Grand Unified Theories*, in *2nd World Summit: Physics Beyond the*
3245 *Standard Model*, 8, 2006 [[hep-ph/0608183](#)]. 46
- 3246 [45] KAMIOKANDE-II collaboration, *Observation of a Neutrino Burst from the*
3247 *Supernova SN 1987a*, *Phys. Rev. Lett.* **58** (1987) 1490. 47
- 3248 [46] R.M. Bionta et al., *Observation of a Neutrino Burst in Coincidence with Supernova*
3249 *SN 1987a in the Large Magellanic Cloud*, *Phys. Rev. Lett.* **58** (1987) 1494. 47
- 3250 [47] DUNE collaboration, *The DUNE Far Detector Vertical Drift Technology*,
3251 *Technical Design Report*, [2312.03130](#). 51, 52
- 3252 [48] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE) Near*
3253 *Detector Conceptual Design Report, Instruments* **5** (2021) 31 [[2103.13910](#)]. 55, 56,
3254 59, 63, 143

BIBLIOGRAPHY

- 3255 [49] J. Strait, E. McCluskey, T. Lundin, J. Willhite, T. Hamernik, V. Papadimitriou
3256 et al., *Long-baseline neutrino facility (lbnf) and deep underground neutrino*
3257 *experiment (dune) conceptual design report volume 3: Long-baseline neutrino*
3258 *facility for dune june 24, 2015*, 1601.05823. 60
- 3259 [50] DUNE DAQ, “dtp-firmware.”
3260 <https://gitlab.cern.ch/dune-daq/readout/dtp-firmware>, 2020. 75
- 3261 [51] DUNE DAQ, “dtp-simulation.”
3262 <https://gitlab.cern.ch/dune-daq/readout/dtp-simulation>, 2020. 78
- 3263 [52] DUNE DAQ, “dtpemulator.”
3264 https://github.com/DUNE-DAQ/dtpemulator/tree/fmlopez/filter_ana, 2022.
3265 78
- 3266 [53] J. McClellan and T. Parks, *A personal history of the Parks-McClellan algorithm*,
3267 *IEEE Signal Processing Magazine* **22** (2005) 82. 79
- 3268 [54] G. Turin, *An introduction to matched filters*, *IEEE Transactions on Information*
3269 *Theory* **6** (1960) 311. 82
- 3270 [55] J.W. Goodman, *Statistical Optics*, Wiley (1985). 84
- 3271 [56] B. Dwork, *Detection of a pulse superimposed on fluctuation noise*, *Proceedings of*
3272 *the IRE* **38** (1950) 771. 85
- 3273 [57] L. Wainstein and V. Zubakov, *Extraction of Signals from Noise*, Prentice-Hall
3274 (1962). 85
- 3275 [58] E.D. Church, *Larsoft: A software package for liquid argon time projection drift*
3276 *chambers*, 1311.6774. 88, 89
- 3277 [59] S.V. Stehman, *Selecting and interpreting measures of thematic classification*
3278 *accuracy*, *Remote Sensing of Environment* **62** (1997) 77. 102

BIBLIOGRAPHY

- 3279 [60] A.A. Taha and A. Hanbury, *Metrics for evaluating 3d medical image segmentation: analysis, selection, and tool*, *BMC Medical Imaging* **15** (2015) . 103
- 3280
- 3281 [61] J. Silk, K.A. Olive and M. Srednicki, *The Photino, the Sun and High-Energy Neutrinos*, *Phys. Rev. Lett.* **55** (1985) 257. 109
- 3282
- 3283 [62] M. Srednicki, K.A. Olive and J. Silk, *High-Energy Neutrinos from the Sun and Cold Dark Matter*, *Nucl. Phys. B* **279** (1987) 804. 109
- 3284
- 3285 [63] J.S. Hagelin, K.W. Ng and K.A. Olive, *A High-energy Neutrino Signature From Supersymmetric Relics*, *Phys. Lett. B* **180** (1986) 375. 109
- 3286
- 3287 [64] T.K. Gaisser, G. Steigman and S. Tilav, *Limits on Cold Dark Matter Candidates from Deep Underground Detectors*, *Phys. Rev. D* **34** (1986) 2206. 109
- 3288
- 3289 [65] N. Bernal, J. Martín-Albo and S. Palomares-Ruiz, *A novel way of constraining WIMPs annihilations in the Sun: MeV neutrinos*, *JCAP* **08** (2013) 011
- 3290
- 3291 [1208.0834]. 109, 110, 111, 117
- 3292 [66] C. Rott, J. Siegal-Gaskins and J.F. Beacom, *New Sensitivity to Solar WIMP Annihilation using Low-Energy Neutrinos*, *Phys. Rev. D* **88** (2013) 055005
- 3293
- 3294 [1208.0827]. 109, 116
- 3295 [67] C. Rott, S. In, J. Kumar and D. Yaylali, *Dark Matter Searches for Monoenergetic Neutrinos Arising from Stopped Meson Decay in the Sun*, *JCAP* **11** (2015) 039
- 3296
- 3297 [1510.00170]. 109
- 3298 [68] DUNE collaboration, *Searching for solar KDAR with DUNE*, *JCAP* **10** (2021) 065
- 3299 [2107.09109]. 109
- 3300 [69] G. Busoni, A. De Simone and W.-C. Huang, *On the Minimum Dark Matter Mass Testable by Neutrinos from the Sun*, *JCAP* **07** (2013) 010 [1305.1817]. 110
- 3301
- 3302 [70] V.A. Bednyakov and F. Simkovic, *Nuclear spin structure in dark matter search: The Zero momentum transfer limit*, *Phys. Part. Nucl.* **36** (2005) 131
- 3303
- 3304 [hep-ph/0406218]. 111

BIBLIOGRAPHY

- 3305 [71] A. Gould, *WIMP Distribution in and Evaporation From the Sun*, *Astrophys. J.* **321**
3306 (1987) 560. 112
- 3307 [72] J.N. Bahcall, M.H. Pinsonneault and S. Basu, *Solar models: Current epoch and*
3308 *time dependences, neutrinos, and helioseismological properties*, *Astrophys. J.* **555**
3309 (2001) 990 [[astro-ph/0010346](#)]. 114
- 3310 [73] T. Golan, J.T. Sobczyk and J. Zmuda, *NuWro: the Wroclaw Monte Carlo*
3311 *Generator of Neutrino Interactions*, *Nucl. Phys. B Proc. Suppl.* **229-232** (2012)
3312 499. 118
- 3313 [74] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for*
3314 *likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [[1007.1727](#)].
3315 119
- 3316 [75] T. Kaluza, *Zum Unitätsproblem der Physik, Sitzungsber. Preuss. Akad. Wiss.*
3317 *Berlin (Math. Phys.)* **1921** (1921) 966 [[1803.08616](#)]. 121
- 3318 [76] O. Klein, *Quantum Theory and Five-Dimensional Theory of Relativity. (In*
3319 *German and English)*, *Z. Phys.* **37** (1926) 895. 121
- 3320 [77] T. Appelquist, H.-C. Cheng and B.A. Dobrescu, *Bounds on universal extra*
3321 *dimensions*, *Phys. Rev. D* **64** (2001) 035002 [[hep-ph/0012100](#)]. 121
- 3322 [78] N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *The Hierarchy problem and new*
3323 *dimensions at a millimeter*, *Phys. Lett. B* **429** (1998) 263 [[hep-ph/9803315](#)]. 121
- 3324 [79] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*,
3325 *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)]. 121
- 3326 [80] G. Servant and T.M.P. Tait, *Is the lightest Kaluza-Klein particle a viable dark*
3327 *matter candidate?*, *Nucl. Phys. B* **650** (2003) 391 [[hep-ph/0206071](#)]. 121
- 3328 [81] H.-C. Cheng, K.T. Matchev and M. Schmaltz, *Radiative corrections to*
3329 *Kaluza-Klein masses*, *Phys. Rev. D* **66** (2002) 036005 [[hep-ph/0204342](#)]. 123

BIBLIOGRAPHY

- 3330 [82] M. Blennow, J. Edsjö and T. Ohlsson, *Neutrinos from WIMP annihilations using a*
3331 *full three-flavor Monte Carlo*, *JCAP* **01** (2008) 021 [[0709.3898](#)]. 123, 126
- 3332 [83] J. Edsjö, J. Elevant and C. Niblaeus, *WimpSim Neutrino Monte Carlo*. 123, 126
- 3333 [84] U. Haisch and A. Weiler, *Bound on minimal universal extra dimensions from*
3334 *anti-B → X(s)gamma*, *Phys. Rev. D* **76** (2007) 034014 [[hep-ph/0703064](#)]. 125
- 3335 [85] A. Freitas and U. Haisch, *Anti-B → X(s) gamma in two universal extra*
3336 *dimensions*, *Phys. Rev. D* **77** (2008) 093008 [[0801.4346](#)]. 125
- 3337 [86] C. Rott, D. Jeong, J. Kumar and D. Yaylali, *Neutrino Topology Reconstruction at*
3338 *DUNE and Applications to Searches for Dark Matter Annihilation in the Sun*,
3339 *JCAP* **07** (2019) 006 [[1903.04175](#)]. 128
- 3340 [87] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel et al.,
3341 *Scikit-learn: Machine learning in Python*, *Journal of Machine Learning Research*
3342 **12** (2011) 2825. 134
- 3343 [88] J. Kopp, V. Niro, T. Schwetz and J. Zupan, *DAMA/LIBRA and leptонically*
3344 *interacting Dark Matter*, *Phys. Rev. D* **80** (2009) 083502 [[0907.3159](#)]. 138
- 3345 [89] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2020**
3346 (2020) 083C01. 139
- 3347 [90] M. Beltran, D. Hooper, E.W. Kolb and Z.C. Krusberg, *Deducing the nature of dark*
3348 *matter from direct and indirect detection experiments in the absence of collider*
3349 *signatures of new physics*, *Phys. Rev. D* **80** (2009) 043509 [[0808.3384](#)]. 139
- 3350 [91] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron.*
3351 *Astrophys.* **641** (2020) A6 [[1807.06209](#)]. 140