


¹ ADVANCING NEUTRINO
² DETECTION AND TRIGGERING IN
³ DUNE



⁵ Francisco Martínez López

⁶ Submitted in partial fulfillment of the requirements
⁷ of the Degree of Doctor of Philosophy

⁸ School of Physical and Chemical Sciences

⁹ Queen Mary University of London

¹⁰ December 2023

¹¹ Statement of originality

- ¹² I, [insert name as recorded in QM records], confirm that the research included within
¹³ this thesis is my own work or that where it has been carried out in collaboration with, or
¹⁴ supported by others, that this is duly acknowledged below and my contribution indicated.
¹⁵ Previously published material is also acknowledged below.
- ¹⁶ I attest that I have exercised reasonable care to ensure that the work is original, and
¹⁷ does not to the best of my knowledge break any UK law, infringe any third party's
¹⁸ copyright or other Intellectual Property Right, or contain any confidential material.
- ¹⁹ I accept that the College has the right to use plagiarism detection software to check the
²⁰ electronic version of the thesis.
- ²¹ I confirm that this thesis has not been previously submitted for the award of a degree
²² by this or any other university.
- ²³ The copyright of this thesis rests with the author and no quotation from it or information
²⁴ derived from it may be published without the prior written consent of the author.
- ²⁵ Signature: [can be digital signature]
- ²⁶ Date:
- ²⁷ Details of collaboration and publications:
²⁸ [insert details here if applicable]

²⁹ Abstract

³⁰ Hello, here is some text without a meaning. This text should show what a printed text
³¹ will look like at this place. If you read this text, you will get no information. Really? Is
³² there no information? Is there a difference between this text and some nonsense like
³³ “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about
³⁴ the selected font, how the letters are written and an impression of the look. This text
³⁵ should contain all letters of the alphabet and it should be written in of the original
³⁶ language. There is no need for special content, but the length of words should match
³⁷ the language.

O time, thou must untangle this, not I.
It is too hard a knot for me to untie!

Twelfth Night

SHAKESPEARE

³⁸ Acknowledgements

³⁹ In principio fu la Maestra Manuela.

⁴⁰ A heartfelt thank you to . . . for accepting to be the referees of this thesis, and for
⁴¹ the thought-provoking discussions we shared. Their comments brought this manuscript
⁴² to a significant improvement.

⁴³ I feel privileged to have this many eminent researchers in my committee. A very big
⁴⁴ thank you for accepting to be part of it.

⁴⁵ Thank you to . . .

⁴⁶ Un ringraziamento va anche agli amici triestini di nascita, di adozione, o di passaggio,
⁴⁷ matematici, sballerine, o altro, oggi sparpagliati per il mondo a formare una famiglia
⁴⁸ grazie alla quale uno non si sente mai troppo solo e lontano da tutti. Tanto siamo
⁴⁹ dappertutto.

⁵⁰ In particolare grazie a . . .

⁵¹ Prima di giungere ai ringraziamenti più personali, . . .

⁵² Contents

⁵³	Statement of originality	3
⁵⁴	Abstract	5
⁵⁵	Acknowledgements	9
⁵⁶	List of Figures	15
⁵⁷	List of Tables	29
⁵⁸	1 Introduction	31
⁵⁹	2 Neutrino physics	33
⁶⁰	2.1 Neutrinos in the SM	33
⁶¹	2.2 Neutrino oscillations	34
⁶²	2.2.1 Oscillations in vacuum	35
⁶³	2.2.2 Oscillations in matter	36
⁶⁴	2.2.3 Current status of neutrino oscillations	37
⁶⁵	2.3 Open questions in the neutrino sector	38
⁶⁶	3 The Deep Underground Neutrino Experiment	41
⁶⁷	3.1 Overview	41
⁶⁸	3.2 Physics goals of DUNE	43
⁶⁹	3.3 Far Detector	45
⁷⁰	3.3.1 Horizontal Drift	46
⁷¹	3.3.2 Vertical Drift	49

CONTENTS

72	3.3.3 FD Data Acquisition System	51
73	3.4 Near Detector	52
74	3.4.1 ND-LAr	54
75	3.4.2 TMS/ND-GAr	55
76	3.4.3 PRISM	56
77	3.4.4 SAND	58
78	3.5 LBNF beamline	58
79	4 ND-GAr	61
80	4.1 Requirements	61
81	4.2 Reference design	62
82	4.2.1 HPgTPC	62
83	4.2.2 ECal	63
84	4.2.3 Magnet	64
85	4.2.4 Muon system	65
86	4.3 GArSoft	65
87	4.3.1 Event generation	65
88	4.3.2 Detector simulation	66
89	4.3.3 Reconstruction	67
90	5 FWTPG offline software	71
91	6 Matched Filter approach to induction wire Trigger Primitives	73
92	6.1 Motivation	73
93	6.2 Signal-to-noise ratio definition	75
94	6.3 Low-pass FIR filter design	77
95	6.4 Matched filters	80
96	6.5 Using simulated samples	86
97	6.5.1 Angular dependence	92
98	6.5.2 Distortion and peak asymmetry	94

CONTENTS

99	6.5.3 Hit sensitivity	97
100	7 DM searches with neutrinos from the Sun	107
101	7.1 Motivation	107
102	7.2 Gravitational capture of DM by the Sun	107
103	7.3 Neutrino flux from DM annihilations	114
104	7.4 Computing limits from solar neutrino fluxes	115
105	7.5 Example: Kaluza-Klein Dark Matter	119
106	7.6 High energy DM neutrino signals	123
107	7.6.1 DIS events	125
108	7.6.2 Single proton QEL events	130
109	7.6.3 Results	133
110	7.7 Example: Leptophilic Dark Matter	135
111	8 Particle ID in GArSoft	141
112	8.1 dE/dx measurement in the TPC	142
113	8.1.1 Energy calibration	144
114	8.1.2 Truncated dE/dx mean	154
115	8.1.3 Mean dE/dx parametrisation	157
116	8.1.4 Proton identification	160
117	8.2 Muon and pion separation in the ECal and MuID	160
118	8.2.1 Track-ECal matching	160
119	8.2.2 Feature selection and importance	166
120	8.2.3 Hyperparameter optimisation	169
121	8.2.4 Probability calibration	169
122	8.2.5 Performance	169
123	8.3 ECal time-of-flight	169
124	8.3.1 Arrival time estimations	169
125	8.3.2 Proton and pion separation	169
126	8.4 Charged pion decay in flight	169

CONTENTS

127	8.4.1 Track breakpoints	169
128	8.5 Neutral particle identification	169
129	8.5.1 ECal clustering	169
130	8.5.2 π^0 reconstruction	173
131	9 Conclusions	175
132	A An appendix	177
133	Bibliography	179

¹³⁴ List of Figures

¹³⁵	3.1 Schematic diagram of the DUNE experiment and the LBNF beamline [1].	42
¹³⁶	3.2 Schematic diagram showing the operating principle of a LArTPC with wire readout	46
¹³⁸	3.3 Proposed design for the FD-1 and FD-2 modules following the HD principle.	47
¹³⁹	3.4 Schematic representation of an APA frames showing the U, V, X and G wires.	48
¹⁴¹	3.5 A PDS module containing 24 X-ARAPUCAs and the location of the modules on the APAs.	48
¹⁴³	3.6 Proposed design for the FD-3 module following the VD principle.	49
¹⁴⁴	3.7 Schematic representation of the electrode strip configuration for a top and bottom CRU.	50
¹⁴⁶	3.8 Detailed diagram of the DUNE FD DAQ system. Figure taken from Ref. [2].	51
¹⁴⁸	3.9 Representation of the ND hall in Phase II, showing the different subcomponents.	53
¹⁴⁹	3.10 Schematic representation of the external components of ND-LAr, including the cryostat and the PRISM movable system and detailed drawing of one ArgonCube module.	54
¹⁵²	3.11 Schematic view of the TMS detector, highlighting its main parts.	55
¹⁵³	3.12 Cross section of the ND-GAr geometry, showing the HPgTPC, ECal and magnet.	56

LIST OF FIGURES

155	3.13	Predicted beam muon neutrino flux at the ND location for different off-axis positions.	57
156	3.14	Schematic longitudinal section of the LBNF beamline at Fermilab.	58
158	3.15	Predicted neutrino fluxes at the FD in FHC mode and RHC mode.	59
159	4.1	Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers.	63
161	4.2	Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers.	64
163	6.1	<i>Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G.</i>	74
167	6.2	<i>Left panel: Zoomed unfiltered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code> (blue line). The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{noise}}$.</i>	75
174	6.3	<i>Left panel: Zoomed filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code> (blue line). The filter used was the current implementation of the low-pass FIR filter in <code>dtp-firmware</code>. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$</i>	77

LIST OF FIGURES

182	6.4 Power spectrum in decibels for the current implementation of the low-pass	
183	FIR filter in <code>dtp-firmware</code> (blue line), compared to the response of an	
184	optimal filter obtained using the Parks-McClellan algorithm for the same	
185	pass-band (red line). Also for comparison I include the spectrum of the	
186	optimal filter when taking only the integer part of the coefficients (red	
187	dashed line)	78
188	6.5 Relative change in the S/N for the ProtoDUNE-SP raw data capture	
189	<code>felix-2020-07-17-21:31:44</code> , using different values of the cutoff frequency	
190	f_c and the transition width δf . The optimal Chebyshev filters were applied	
191	using just the integer part of the coefficients given by the Parks-McClellan	
192	algorithm.	79
193	6.6 Distribution of the relative change of the S/N on the different wire planes	
194	from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code>	
195	after the optimal Chebyshev filter was applied. The filter was computed	
196	with the Parks-McClellan algorithm using a cutoff of $f_c = 0.068 \text{ ticks}^{-1}$	
197	and a transition width $\delta f = 0.010 \text{ ticks}^{-1}$	80
198	6.7 Left panel: Zoomed match filtered waveform corresponding to channel 7840	
199	from the ProtoDUNE-SP raw data capture <code>felix-2020-07-17-21:31:44</code>	
200	(blue line). The filter used was directly extracted from the data, being	
201	the 32 values around the first peak in the original waveform. The green	
202	dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is	
203	also shown (red line). Top right panel: ADC distribution for channel 7840	
204	after match filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$.	
205	Bottom right panel: noise ADC distribution for channel 7840 after match	
206	filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$	81
207	6.8 Relative improvement in the S/N for the raw data capture <code>felix-2020-07-17-21:31:44</code> ,	
208	using the matched filter following the parametrisation in Eq. (6.17). The	
209	black crosses in both panels denote the location of the maximum ratio value.	84

LIST OF FIGURES

210	6.9 Left panel: Optimal matched filter coefficients for the U (blue line) and V (red line) planes. The filters were computed with our parametrisation in Eq. (6.17) for the parameter values $\delta = 0.035$, $\sigma = 0.191$ and $\delta = 0.018$, $\sigma = 0.191$ respectively. Right panel: Distribution of the relative change of the S/N on the two induction wire planes from the ProtoDUNE-SP raw data capture <i>felix-2020-07-17-21:31:44</i> after their respective optimal matched filters were applied.	87
217	6.10 Left panel: distributions of the particles track length in the liquid argon for the generated $E_k = 100$ MeV monoenergetic samples, electrons (blue), muons (red), protons (green) and neutral pions (purple). Right panel: distribution of the length of the longest photon in the neutral pion sample after the decay process $\pi^0 \rightarrow \gamma\gamma$	88
222	6.11 Left panel: schematic representation of the two new rotated reference frames used in this analysis (denoted as prime and double prime), viewed from the yz plane. The magenta stack of lines represent the wires in the U plane, whereas the green lines correspond to the wires in the V plane. Right panel: 3D representation of the momentum of one of the generated monoenergetic muons (red arrow) in the original reference frame (black lines), along with the new reference frame used for the U plane waveforms (blue lines). In the yz plane I added the projection of these three.	90
230	6.12 Distributions of the mean S/N improvement per event for the corresponding sample after applying the matched filters. Here I separated the change in the U plane (blue) and the V plane (red) channels. From top left to the right: muon, electron, proton and neutral pion. All the events have a fixed kinetic energy of $E_k = 100$ MeV.	91

LIST OF FIGURES

LIST OF FIGURES

255	6.16 Left panel: peak asymmetry distribution for the case of the monoenergetic	
256	$E_k = 100$ MeV muon sample. Each value corresponds to a single bipolar	
257	signal peak from a channel in any event. The blue distribution represents	
258	the peaks on U plane channels, whereas the red corresponds to signal peaks	
259	in V wires. Right panel: relation between the mean peak asymmetry per	
260	event with the S/N for U channel waveforms from the $E_k = 100$ MeV	
261	muon sample. The top subplot shows the decimal logarithm of the mean	
262	S/N for the raw (red) and the matched filtered (blue) waveforms. The	
263	bottom subplot contains the mean S/N improvement ratio after the matched	
264	filter was applied.	96
265	6.17 Raw data display in the plane time (in firmware ticks) vs. offline channel	
266	number for an $E_k = 100$ MeV electron event. The produced true hits are	
267	superimposed (black boxes) as well as the hits comming from the standard	
268	hit finder chain (blue circles) and the hit finder using the matched filter	
269	(green triangles).	98
270	6.18 Dependence of the precision (blue), sensitivity (red) and F_1 (green) scores	
271	on the threshold values used in the hit finder, for the FIR (left panel)	
272	and matched filter (right panel) cases. The results were obtained after	
273	matching the hits to the true hits in the case of the isotropic muon sample	
274	with kinetic energy in the range 5 to 100 MeV, taking only into account	
275	the induction plane channels. The points represent the mean value while	
276	the error bars indicate one standard deviation around that mean value. . .	100

LIST OF FIGURES

- 277 6.19 *Dependence of the averaged hit sensitivity on the kinetic energy of the*
 278 *events for the matched filter (blue) and standard (red) hits, for the case of*
 279 *the muon (left panel) and electron (right panel) samples, separated between*
 280 *U (top plots) and V (bottom plots) induction wire planes. The top subplots*
 281 *contain the hit sensitivities for the two hit finder alternatives, while the*
 282 *bottom subplots show the ratio between the two. The horizontal lines sit at*
 283 *the mean value and represent the size of the energy bins, while the vertical*
 284 *error bars indicate one standard deviation around that mean value.* 102
- 285 6.20 *Distributions of the hit sensitivity in the U (top panels) and V (bottom*
 286 *panels) planes versus the hit sensitivity in the X plane, both for the*
 287 *standard hits (left panels) and the matched filter hits (right panels), in the*
 288 *case of the electron sample and a threshold of 30 ADC.* 103
- 289 6.21 *Top panels: standard residual plots of the hit sensitivities between the X*
 290 *and U planes. Bottom panels: quantile-quantile plots of the hit sensitivity*
 291 *standard residuals between the X and U planes. In all cases, the left*
 292 *panel corresponds to the standard hits while the right panel represents the*
 293 *matched filter case, all from the electron sample with a 30 ADC threshold.* 104
- 294 7.1 *Input solar parameters used in our capture rate computation as functions*
 295 *of the Sun's radius, from left to right: temperature (with respect to the*
 296 *temperature at the core), mass (in solar masses) and electron number*
 297 *density (with respect to the electron density at the core). All quantities*
 298 *shown correspond to the standard solar model BS2005-OP [3].* 111
- 299 7.2 *Capture rates as a function of the DM mass for the DM-electron interactions*
 300 *(red lines), SD DM-nucleons interactions (green lines) and SI DM-nucleons*
 301 *interactions (blue lines). Solid lines represent the values computed in this*
 302 *work while the dashed lines are the one given in Ref. [4]. All the rates*
 303 *are shown for a choice of scattering cross section of $\sigma_i = 10^{-40} \text{ cm}^2$.* . . 112

LIST OF FIGURES

304	7.3 <i>NuWro computed $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section as a function of the neutrino energy E_μ. The black line shows to the total cross section, whereas the others correspond to the different contributions (in red quasi-elastic scattering, in green resonant pion exchange, in blue deep inelastic scattering and in purple meson exchange current).</i>	116
309	7.4 <i>Expected atmospheric neutrino flux as a function of the neutrino energy E_ν at Homestake at solar minimum, taken from Ref. [5]. The blue solid (dashed) line correspond to muon neutrinos (antineutrinos) and the red solid (dashed) line correspond to electron neutrinos (antineutrinos)</i>	118
313	7.5 Feynman diagrams for B^1B^1 annihilation into SM fermions.	120
314	7.6 Feynman diagrams for B^1B^1 annihilation into a Higgs boson pair. . . .	120
315	7.7 <i>Computed spectra of muon neutrinos at the DUNE FD site from B^1 annihilations in the Sun for three different values of M_{LKP}, plotted in relative energy units for legibility.</i>	121
318	7.8 <i>Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin- dependent B^1-proton scattering cross section as a function of M_{LKP} (green dots). I also show the previous limits from IceCube [6] (blue line) and Antares [7] (red line) on the LKP cross section. The shaded area represents the disfavoured region (at 95% confidence level) on the mass of the LKP from LHC data [8]</i>	122
324	7.9 <i>Computed spectra of muon neutrinos at the DUNE FD site from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilations in the Sun for the DM masses $m_{\text{DM}} = 10$ GeV (red line), 50 GeV (green line) and 100 GeV (blue line), plotted in relative energy units.</i>	123
328	7.10 <i>Distribution of the muon neutrino energies from the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilation channels, for $m_{\text{DM}} = 10$ GeV, separated by CC interaction type: QEL (blue), MEC (orange), RES (green) and DIS (red)</i>	125

LIST OF FIGURES

<p>332 7.11 <i>Distributions of θ_μ (left panel), θ_j (central panel) and θ_{plane} (right panel)</i> 333 <i>for the $b\bar{b}$ sample with $m_{DM} = 10$ GeV (blue) and the atmospheric</i> 334 <i>background (red).</i></p> <p>335 7.12 <i>Left panel: signal efficiencies (blue lines) and background rejections (red</i> 336 <i>lines) for events passing the cuts $\theta < \theta_{cut}$ for the jet (solid lines) and</i> 337 <i>muon (dashed lines) angles. Right panel: signal efficiency (blue line) and</i> 338 <i>background rejection (red line) for events passing the cut $\theta_{plane} < \theta_{cut}$ for</i> 339 <i>the momentum conservation plane deviation.</i></p> <p>340 7.13 <i>Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) DIS samples</i> 341 <i>as functions of the DM mass, m_{DM}, obtained by applying the optimal</i> 342 <i>angular cuts $\theta_\mu < 27^\circ$, $4^\circ < \theta_j < 26^\circ$ and $\theta_{plane} < 3.5^\circ$.</i></p> <p>343 7.14 <i>Distributions of $\cos \theta_\mu$ (left panel), $\cos \theta_p$ (central panel) and $\cos \theta_N$</i> 344 <i>(right panel) for the $\tau^+\tau^-$ QEL sample with $m_{DM} = 5$ GeV (blue) and</i> 345 <i>the atmospheric background (red).</i></p> <p>346 7.15 <i>Left panel: value of the loss function for the training sample (blue line)</i> 347 <i>and accuracy for the validation sample (red line) versus the number of</i> 348 <i>iterations for the MLP classifier training. Right panel: distributions of the</i> 349 <i>predicted probabilities assigned by the MLP classifier to the test sample</i> 350 <i>for the $\tau^+\tau^-$ QEL signal with $m_{DM} = 5$ GeV (blue) and the atmospheric</i> 351 <i>background (red).</i></p> <p>352 7.16 <i>Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) single proton</i> 353 <i>QEL samples as functions of the DM mass, m_{DM}, obtained by requiring a</i> 354 <i>minimum predicted probability from the MLP classifier of 0.97 in order to</i> 355 <i>achieve a background rejection greater than 99.8%.</i></p>	<p>126</p> <p>127</p> <p>129</p> <p>130</p> <p>131</p> <p>132</p>
---	---

LIST OF FIGURES

356	7.17 Projected 90% confidence level upper limit for DUNE (400 kT yr) on	
357	the spin-dependent DM-nucleon scattering cross section as a function of	
358	m_{DM} , for the annihilation channels $\tau^+\tau^-$ (blue) and $b\bar{b}$ (red) separated	
359	by interaction type (up triangles denote DIS interactions whereas down	
360	triangles represent QEL interactions). I also show the previous limits	
361	from IceCube [9] (solid lines) and the projected sensitivities for Pingu [10]	
362	(dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines), as well	
363	as the direct detection limits from PICASSO [12] (solid green line) and	
364	PICO-60 C ₃ F ₈ [13] (dashed green line).	134
365	7.18 Left panel: Projected 90% confidence level sensitivity of DUNE (400 kT	
366	yr) to the scale Λ of an EFT containing only leptophilic DM axial-axial	
367	interactions (blue line). Right panel: . In both cases the corresponding	
368	limits from DarkSide-50 [14] (dotted green line) and XENON1T [15]	
369	(dashed red line) are also shown, together with the configurations for which	
370	the correct relic density is achieved (black line), all for the coupling values	
371	$c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$	138
372	8.1 Left panel: distribution of the fraction of Geant4-level energy deposits per	
373	track with residual range less than 20% of the total track length, for the	
374	isotropic proton sample. Right panel: distribution of the ionisation per	
375	unit length of the energy deposits in the proton sample after removing	
376	the tracks with less than 30% of their energy deposits in the last 20% of	
377	the track.	145
378	8.2 Left panel: distribution of the reconstructed ionisation charge per unit	
379	length for our MC stopping proton sample. The different colors indicate	
380	how many consecutive dQ/dx pairs were grouped together. Right panel:	
381	distribution of the median change in dQ/dx per track after $N_{group} = 4$	
382	clusters were reclustered together.	146

LIST OF FIGURES

383	8.3 Distribution of the Geant4-simulated energy losses per unit length versus	
384	residual range for the stopping proton sample. The overlaid points	
385	represent the fitted most probable value of the dE/dx distribution in each	
386	residual range bin, whereas the curve is their best fit to the Bragg-Kleeman	
387	formula from Eq. (8.4).	148
388	8.4 Fitted most probable dQ/dx values for each dE/dx bin (red points),	
389	obtained from the stopping proton sample. The overlaid curve (black	
390	line) represents the best fit to the logarithmic calibration function from	
391	Eq. (8.5).	149
392	8.5 Fitted most probable dQ/dx values for each dE/dx bin for three different	
393	ADC bit limits, 10 (blue points), 12 (default, yellow points) and 16-bit	
394	(red points).	151
395	8.6 Top panel: area normalised dE/dx distributions for the true (solid grey)	
396	and the reconstructed energy deposits in the stopping proton sample,	
397	both after applying the calibration (blue) and the calibration and the	
398	normalisation correction (yellow). Also shown is the distribution obtained	
399	by applying a correction factor to the dQ/dx values but not the calibration	
400	(red). Bottom panel: fractional residuals for the uncorrected (blue),	
401	corrected (yellow) and uncalibrated (red) samples.	152
402	8.7 Left panel: fractional residuals between the true and the corrected dE/dx	
403	means (blue) and the 60% truncated means (yellow), for each event in the	
404	stopping proton sample. Right panel: fractional residuals between the	
405	true and the uncorrected (blue), corrected (yellow) and uncalibrated (red)	
406	dE/dx 60% truncated means, for each event in the stopping proton sample.	154
407	8.8 Estimated values of the mean dE/dx bias (left panel) and resolution	
408	(right panel) obtained using the corrected data from the stopping proton	
409	sample, for different values of the truncation factor.	155
410	8.9 Examples of the truncated mean dE/dx LanGauss fits for various $\beta\gamma$	
411	bins, from a simulated FHC neutrino sample.	157

LIST OF FIGURES

412	8.10 Resulting one and two dimensional projections of the posterior probability distributions of the ALEPH $\langle dE/dx \rangle$ parameters obtained by fitting the 60% truncated mean dE/dx values from a FHC neutrino sample in ND-GAr. The vertical dashed lines in the 1D distributions represent the 16th, 50th and 84th percentiles.	158
417	8.11 Truncated mean dE/dx obtained for the FHC neutrino sample as a function of the $\beta\gamma$ product (upper panel). Also shown are the fitted most probable values for each $\beta\gamma$ bin (red points) and the best fit obtained using the ALEPH parametrisation (black line). The residuals resulting from the fit are shown in the lower panel.	160
422	8.12 Estimated values of the mean dE/dx bias (left panel) and resolution (right panel) obtained using the corrected data from the stopping proton sample, for different values of the truncation factor.	161
425	8.13 Distributions of energy deposits in the ECal for a muon (left) and a charged pion (right) with similar momentum. The energy is projected onto the plane perpendicular to the principal component of the hit clusters, and the positions are relative to the center of the interaction.	162
429	8.14 Left panel: comparison between the precision (blue), sensitivity (yellow) and F_1 score (red) obtained for the default (horizontal lines) and new algorithms, both with the χ^2 -based direction estimator (squares) and cheating the directions (circles), for different values of the χ^2 cut. Right panel: comparison of the performance of the new algorithm when applying the cluster t_0 correction (squares) and when (circles).	163
435	8.15 Schematics of a possible option to deal with track-ECal associations in non-zero t_0 neutrino interaction events, trying to correct for the drift direction uncertainty in a cluster-by-cluster basis using the cluster time, $t_{cluster}$	165

LIST OF FIGURES

439	8.16 Mean values of the F_1 -score marginal distributions for the different 440 free parameters of the new clustering algorithm, with the error bars 441 representing one standard deviation around the mean. The F_1 -score 442 values were computed for the 6561 possible parameter configurations 443 using 1000 ν_μ CC interaction events.	171
444	8.17 Left panel: distributions of the number of ECal clusters per photon from 445 π^0 decays for the standard (red) and new (blue) clustering algorithms. 446 Right panel: reconstructed invariant mass distributions for photon pairs 447 from single π^0 events using the standard (red) and new (blue) ECal 448 clustering algorithms.	173

List of Tables

450	2.1	Summary of neutrino oscillation parameters determined in the Neutrino	
451		Global Fit of 2020 [16].	38
452	3.1	Summary of the two-phased plan for DUNE	43
453	3.2	Exposure and time required to achieve the different physics milestones of	
454		the two phases	44
455	6.1	<i>Characteristic parameters of the two monoenergetic muon events selected, relative to the U plane: projected angles in the xz' and $y'z'$ planes, S/N values for the raw and filtered waveforms, mean improvement of the S/N and peak asymmetry.</i>	95
459	8.1	Calibration parameters obtained from the fit of the ND-GAr simulated	
460		stopping proton sample to the calibration function from Eq. (8.5). The	
461		fits were performed for the 10, 12, and 16-bit ADC limits.	151
462	8.2	Summary of parameters and sampled values used in the optimisation of	
463		the clustering algorithm.	171

⁴⁶⁴ Chapter 1

⁴⁶⁵ Introduction

466 Chapter 2

467 Neutrino physics

468 Ever since they were postulated in 1930 by Wolfgang Pauli to explain the continuous
469 β decay spectrum [17] and later found by Reines and Cowan at the Savannah River
470 reactor in 1953 [18], neutrinos have had a special place among all other elementary
471 particles. They provide a unique way to probe a wide range of quite different physics,
472 from nuclear physics to cosmology, from astrophysics to colliders. Moreover, there is
473 compelling evidence to believe that the study of neutrinos may be key to unveil different
474 aspects of physics beyond the SM, difficult to test elsewhere.

475 In this Chapter I will review the basics of neutrino physics, from its role within the
476 SM to the main open questions related to the neutrino sector, paying special attention
477 to the phenomenology of neutrino oscillations.

478 2.1 Neutrinos in the SM

479 By definition, in the SM there are no right-handed neutrino fields. A direct implication
480 of this fact is that neutrinos are strictly massless within the SM. This follows from the
481 experimental observation that all neutrinos produced via weak interactions are pure
482 left-handed helicity states (and similarly antineutrinos are pure right-handed states).
483 The hypothetical existence of right-handed neutrinos could be indirectly inferred from
484 the observation of non-zero neutrino masses, nevertheless the existence neutrino masses

Chapter 2. Neutrino physics

485 is not a sufficient condition for the existence of such fields.

486 In the SM neutrinos appear in three flavours, namely ν_e , ν_μ and ν_τ . These are
487 associated with the corresponding charged leptons e , μ and τ , in such a way that the
488 charged current part of the Lagrangian coupling them is diagonal. As in the electroweak
489 theory neutrinos are coupled to the Z boson in a universal way, by measuring the so-called
490 invisible decay width of the Z we have an estimate of the number of light (i.e. lighter
491 than the Z boson) neutrino flavours. This number was measured by LEP in a combined
492 analysis of $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow$ hadrons to be $N_\nu = 2.9840 \pm 0.0082$ [19].

493 2.2 Neutrino oscillations

494 The evidence for neutrino oscillation [20], and therefore the existence of non-zero neutrino
495 masses, constitutes one of the groundbreaking discoveries of modern Physics and has
496 acted as driving force for Beyond the Standard Model (BSM) Physics. The minimal
497 extension of the Standard Model (SM) we can do to address these phenomena is
498 introducing distinct masses for at least two of the neutrinos. This way, we are left with
499 three neutrino mass eigenstates ν_1 , ν_2 and ν_3 , with masses m_1 , m_2 and m_3 respectively,
500 which in general will not coincide with the flavour eigenstates ν_e , ν_μ and ν_τ .

501 The way to relate these two sets of neutrino eigenstates is via a 3×3 unitary matrix,
502 called the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [21, 22], as:

$$|\nu_\alpha\rangle = \sum_{i=1}^3 U_{\alpha i}^* |\nu_i\rangle, \quad (2.1)$$

503 where the Greek index α denotes the flavour $\{e, \mu, \tau\}$ and the Latin index i the associated
504 masses $\{1, 2, 3\}$. This leptonic mixing matrix may be parametrized in terms of 6
505 parameters, 3 of which are mixing angles θ_{12} , θ_{13} and θ_{23} , one CP-violating phase δ_{CP}

2.2. Neutrino oscillations

506 and 2 Majorana phases α and β :

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{CP}} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta_{CP}} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\beta} \end{pmatrix}, \quad (2.2)$$

507 where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$. This matrix is analogous to the Cabibbo-Kobayashi-
508 Maskawa (CKM) matrix in the quark sector. If neutrinos are Dirac fermions, we can
509 drop the Majorana phases in the PMNS matrix. But, in any case, these phases play no
510 role on the neutrino oscillations.

511 2.2.1 Oscillations in vacuum

512 Consider the case where a neutrino of flavour α is produced at $t = 0$, and then it
513 propagates through vacuum. Such a state will evolve in time according to the relation:

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^3 U_{\alpha i}^* e^{-iE_i t} |\nu_i(t=0)\rangle, \quad (2.3)$$

514 as the mass eigenstates are also eigenstates of the free Hamiltonian. Now, if we express
515 the mass eigenstates as a superposition of flavour eigenstates, the last expression can be
516 rewritten as:

$$|\nu_\alpha(t)\rangle = \sum_{i=1}^3 U_{\beta i} e^{-iE_i t} U_{\alpha i}^* |\nu_\beta\rangle. \quad (2.4)$$

517 This way, the probability for the neutrino to transition from flavour α to flavour β
518 will be given by:

$$P(\nu_\alpha \rightarrow \nu_\beta) = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_{i=1}^3 U_{\beta i} e^{-iE_i t} U_{\alpha i}^* \right|^2. \quad (2.5)$$

519 A usual approximation to take at this point is to consider ultra-relativistic neutrinos,
520 i.e. $E \approx |\vec{p}|$, so we can write the dispersion relations as:

$$E_i = \sqrt{p^2 + m_i^2} \approx E + \frac{m_i^2}{2E}, \quad (2.6)$$

Chapter 2. Neutrino physics

so we can write the oscillation probability as:

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\beta) &= \sum_{i,j} U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^* e^{-i \frac{\Delta m_{ij}^2}{2E} t} \\ &= \delta_{\alpha\beta} - 4 \sum_{i < j} \Re [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin^2 \left(\frac{\Delta m_{ij}^2}{4E} t \right) \\ &\quad + 2 \sum_{i < j} \Im [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin \left(\frac{\Delta m_{ij}^2}{2E} t \right), \end{aligned} \quad (2.7)$$

where Δm_{ij}^2 is the difference of the squared masses of the j th and i th neutrino mass eigenvalues. At this point, it is usual to write the phase responsible for the oscillations as (under the approximate assumption $t \approx L$):

$$\Delta_{ij} \equiv \frac{\Delta m_{ij}^2}{4E} L \simeq 1.27 \frac{\Delta m_{ij}^2}{(\text{eV}^2)} \frac{L}{(\text{km})} \frac{(\text{GeV})}{E}. \quad (2.8)$$

Notice that, in the case of antineutrinos the only difference would be the sign of the last term in the oscillation probability. This way, one can write the CP asymmetry as:

$$\begin{aligned} A_{CP}^{\alpha\beta} &= P(\nu_\alpha \rightarrow \nu_\beta) - P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) \\ &= 4 \sum_{i < j} \Im [U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*] \sin 2\Delta_{ij}. \end{aligned} \quad (2.9)$$

2.2.2 Oscillations in matter

When neutrinos propagate through matter, their oscillation can be affected in mainly two ways. First, neutrinos can inelastically scatter with nuclei, thus destroying the coherent propagation of their quantum state. Nevertheless, in most cases this effect is negligible (even in very dense mediums like the core of the Sun). Second, neutrinos can also experience coherent or forward scatterings, that can affect their oscillation but not lose the coherent propagation of the state.

The first proposed model to account for neutrino oscillations in matter was proposed by Mikhaev, Smirnov and Wolfenstein (MSW) [23]. It relies on the fact that, as the only charged lepton present in ordinary matter is the electron, electron neutrinos can

2.2. Neutrino oscillations

537 undergo both charged and neutral-current interactions with matter whereas for muon
538 and tau neutrinos just neutral currents are possible.

539 2.2.3 Current status of neutrino oscillations

540 A wide range of neutrino experiments provide experimental input to the neutrino
541 oscillation framework, both using natural or synthetic neutrino sources. The results
542 from one of the neutrino global fit analyses, shown in Tab. 2.1¹, summarise well our
543 current understanding of the different oscillation parameters.

544 **Solar neutrino experiments** detect neutrinos produced in thermonuclear reactions
545 inside the Sun, mainly from the so-called *pp* chain and the CNO cycle. These neutrinos
546 have a typical energy in the range from 0.1 to 20 MeV. These experiments (Homestake
547 [24], GALLEX [25], SAGE [26], Borexino [27], Super-Kamiokande [28] and SNO [29])
548 provide the best sensitivities to θ_{12} and Δm_{21}^2 .

549 **Atmospheric neutrino experiments** detect the neutrino flux produced when
550 cosmic rays scatter with particles in Earth's atmosphere. These collisions generate particle
551 showers that eventually produce electron and muon neutrinos (and antineutrinos). Their
552 energies range from few MeV to about 10^9 GeV. Experiments, like Super-Kamiokande
553 [30] and IceCube [31] use atmospheric neutrinos to measure oscillations and are specially
554 sensitive to θ_{23} and Δm_{32}^2 .

555 **Reactor neutrino experiments** look for the $\bar{\nu}_e$ spectrum produced by nuclear
556 reactors, with energies in the MeV scale. Depending on the distance to the source,
557 long-baseline experiments like KamLAND [32] are sensitive to the solar mass splitting
558 Δm_{21}^2 whereas much shorter baseline experiment such as RENO [33] or DayaBay [34]
559 measure θ_{13} and Δm_{31}^2 .

560 **Accelerator experiments** measure neutrino fluxes generated in particle accelerators.
561 Usually mesons are produced in the accelerator to be focused into a beam, then some
562 decay to muon neutrinos and the rest are absorbed by a target. Depending on the

¹These are the results reported during M. Tórtola's talk at Neutrino 2024 (see this link). I need to keep an eye and see if they publish these or other updated results in the near future.

Chapter 2. Neutrino physics

Table 2.1: Summary of neutrino oscillation parameters determined in the Neutrino Global Fit of 2020 [16].

Parameter	Best fit $\pm 1\sigma$	3σ range
Δm_{21}^2 [eV $^2 \times 10^{-5}$]	$7.55^{+0.22}_{-0.20}$	6.98 – 8.19
$ \Delta m_{31}^2 $ [eV $^2 \times 10^{-3}$] (NO)	$2.51^{+0.02}_{-0.03}$	2.43 – 2.58
$ \Delta m_{31}^2 $ [eV $^2 \times 10^{-3}$] (IO)	$2.41^{+0.03}_{-0.02}$	2.34 – 2.49
$\sin^2 \theta_{12}/10^{-1}$	3.04 ± 0.16	2.57 – 3.55
$\sin^2 \theta_{23}/10^{-1}$ (NO)	$5.64^{+0.15}_{-0.21}$	4.23 – 6.04
$\sin^2 \theta_{23}/10^{-1}$ (IO)	$5.64^{+0.15}_{-0.18}$	4.27 – 6.03
$\sin^2 \theta_{13}/10^{-2}$ (NO)	$2.20^{+0.05}_{-0.06}$	2.03 – 2.38
$\sin^2 \theta_{13}/10^{-2}$ (IO)	$2.20^{+0.07}_{-0.04}$	2.04 – 2.38
δ_{CP}/π (NO)	$1.12^{+0.16}_{-0.12}$	0.76 – 2.00
δ_{CP}/π (IO)	$1.50^{+0.13}_{-0.14}$	1.11 – 1.87

563 configuration one can obtain a beam made of mostly neutrinos or antineutrinos. The
 564 typical energies of these neutrinos are in the GeV range. Experiments such as NOvA
 565 [35], T2K [36], MINOS [37], OPERA [?] and K2K [38] (and in the future DUNE [39])
 566 are primarily sensitive to θ_{13} , θ_{23} and Δm_{32}^2 . Also, in the coming years DUNE [39] and
 567 Hyper-Kamiokande [40] will be sensitive to δ_{CP} .

568 2.3 Open questions in the neutrino sector

569 A crucial question that remains open these days, and is of vital importance for oscillation
 570 phenomena, is whether the mass eigenvalue ν_3 is the heaviest (what we call normal
 571 ordering) or the lightest (referred to as inverted ordering) of the mass eigenstates. In
 572 other words, this means that we do not know the sign of Δm_{32}^2 , so we can either have
 573 $m_1 < m_2 < m_3$ (NO) or $m_3 < m_1 < m_2$ (IO).

574 Another big puzzle is related to the value of δ_{CP} . Nowadays it is poorly constrained,
 575 with all values between π and 2π being consistent with data. A prospective measurement
 576 different from $\delta_{CP} = 0, \pi$ will predict CP-violation in the leptonic sector, and thus

2.3. Open questions in the neutrino sector

577 contribute along with the one measured in the quark sector to the total amount of
578 CP-violation. Although it is true that these two contributions by themselves are not
579 enough to explain the matter anti-matter asymmetry in our universe, the amount of
580 CP-violation in the leptonic sector can be key to explain such imbalance.

581 Both of these questions, because of their nature, could be understood thanks to
582 future oscillation experiments.

583 Notwithstanding, there are other mysteries that can not be unveiled just by conducting
584 oscillation experiments, as certain quantities do not influence these phenomena. Among
585 these there is the question of the absolute values of the neutrino masses. Depending
586 on the value of the lightest of the neutrino masses we can have different mass spectra,
587 from hierarchical $m_1 \ll m_2 < m_3$ (NO) or $m_3 \ll m_1 < m_2$ (IO) to quasi-degenerate
588 $m_1 \simeq m_2 \simeq m_3$.

589 Other open question concerns the nature itself of the neutrinos. If neutrinos are Dirac
590 particles then their mass term can be generated through the usual Higgs mechanism
591 by adding right-handed neutrino fields. However, if they are Majorana particles and
592 therefore their own antiparticles, there is no need to add extra fields to have the mass
593 term in the Lagrangian. Experiments like SuperNEMO [?], SNO+ [?] and NEXT [?],
594 which search for neutrino-less double beta decay, will be able to determine whether
595 neutrinos are Dirac or Majorana.

596 Chapter 3

597 The Deep Underground Neutrino
598 Experiment

599 The Deep Underground Neutrino Experiment (DUNE) is a next generation long-baseline
600 neutrino experiment [1]. It will aim to address several questions in neutrino physics,
601 study neutrinos from astrophysical sources and search for beyond the standard model
602 physics.

603 This chapter reviews the main goals of the DUNE experiment, the design of the far
604 detector modules and their data acquisition (DAQ) system, and the role that the near
605 detector plays in the physics program of DUNE.

606 3.1 Overview

607 The main physics goals of DUNE are:

- 608 • measure the neutrino mass hierarchy, the amount of CP violation in the leptonic
609 sector and the θ_{23} octant,
- 610 • detect rare low energy neutrino events, like neutrinos from supernova bursts, and
- 611 • search for proton decay and other beyond the standard model phenomena.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.1: Schematic diagram of the DUNE experiment and the LBNF beamline [1].

The design of DUNE has been tailored with these goals in mind. It will consist of two neutrino detectors. A near detector (ND) complex will be placed in Fermilab, 574 m downstream of the neutrino production point, whereas a larger far detector (FD) will be built in the Sandford Underground Research Facility (SURF), South Dakota, approximately 1300 km away. Fig. 3.1 shows a simplified view of the various components of DUNE (not to scale).

The beam neutrinos to be used in DUNE will be provided by the LBNF beamline, the multi-megawatt wide-band neutrino beam planned for Fermilab. First, an intense proton beam is extracted from the Fermilab Main Injector. Then, these protons with energies between 60 GeV and 120 GeV collide with a high-power production target and produce charged mesons. Two magnetic horns allow to focus the mesons and perform a sign selection (thus having the capability to switch between neutrino and antineutrino mode). Soon after that, the mesons decay and produce neutrinos (or antineutrinos) which are then aimed to SURF.

Before arriving to the FD, the neutrino beam meets the ND complex, which serves as the experiment's control. Its role is to measure the unoscillated neutrino energy spectra. From these we can predict the unoscillated spectra at the FD, which can be compared to the spectra measured at the FD in order to extract the oscillation parameters. Therefore, the design of the DUNE ND is mainly driven by the needs of the oscillation physics program.

The liquid Argon time projection chamber (LArTPC) technology has been chosen for

3.2. Physics goals of DUNE

Table 3.1: Summary of the two-phased plan for DUNE. Adapted from Ref. [41].

Parameter	Phase I	Phase II	Benefit
FD mass	20 kt fiducial	40 kt fiducial	FD statistics
Beam power	up to 1.2 MW	2.4 MW	FD statistics
ND config.	ND-LAr, TMS, SAND	ND-LAr, ND-GAr, SAND	Systematic constraints

the FD modules of DUNE. Its four modules will record neutrino interactions from the accelerator-produced beam arriving at predictable times. As it also aims at recording rare events, the FD requires trigger schemes which can deal with both kinds of physics, and also maximum uptime.

DUNE is planned to be built using a staged approach consisting on two phases, which are summarised in Tab. 3.1. Phase I consists of a FD with 50% of the total fiducial mass, a reduced version of the ND complex and a 1.2 MW proton beam. It will be sufficient to achieve some early physics goals, like the determination of the neutrino mass ordering. For its Phase II, DUNE will feature the full four FD modules, a more capable ND and a 2.4 MW proton beam. The physics milestones for the two phases are given in Tab. 3.2, in a staging scenario which assumes that Phase II is completed after 6 years of operation.

A summary of the DUNE science program can be found in the DUNE FD Technical Design Report (TDR) Volume I [1]. For a detailed discussion on the two-phased approach the reader is referred to the DUNE Snowmass 2021 report [41].

3.2 Physics goals of DUNE

As noted in the literature (see for instance Ref. [16] for a review), the parameter space of the neutrino oscillation phenomena within the three-flavour picture is quite constrained by current experimental data. However, there are still crucial open questions, like the mass ordering, the value of δ_{CP} or the θ_{13} octant. One of the main goals of DUNE is to shed some light on the values of these parameters [42].

To address these questions DUNE can look to the subdominant oscillation channel

Chapter 3. The Deep Underground Neutrino Experiment

Table 3.2: Exposure and time required to achieve the different physics milestones of the two phases. The predictions assume a Phase II staging scenario where FD modules 3 and 4 are deployed in years 4 and 6 and both the beam and ND are upgraded after 6 years. Adapted from Ref. [41].

Stage	Physics milestone	Exposure (kt-MW-years)	Years (staged)
Phase I	5σ MO ($\delta_{CP} = -\pi/2$)	16	1-2
	5σ MO (100% of the δ_{CP} values)	66	3-5
	3σ CPV ($\delta_{CP} = -\pi/2$)	100	4-6
Phase II	5σ CPV ($\delta_{CP} = -\pi/2$)	334	7-8
	δ_{CP} resolution of 10 degrees ($\delta_{CP} = 0$)	400	8-9
	5σ CPV (50% of the δ_{CP} values)	646	11
	3σ CPV (75% of the δ_{CP} values)	936	14
	$\sin^2(2\theta_{13})$ resolution of 0.004	1079	16

655 $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) and study the energy dependence of the ν_e ($\bar{\nu}_e$) appearance probability.

656 When we focus on the antineutrino channel $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ there is a change in the sign of δ_{CP} ,
 657 thus introducing CP-violation. Moreover, due to the fact that there are no positrons in
 658 the composition of Earth, there is a sign difference for the matter effect contribution
 659 when looking to the antineutrino channel. This asymmetry is proportional to the baseline
 660 length L and is sensitive to the sign of Δ_{31} , and thus to the neutrino mass ordering.

661 Another of the main physics goals of DUNE is the search for baryon-number violating
 662 processes. Specifically, it will try to answer the question of whether protons are stable
 663 or not. There is no symmetry argument that forbids protons from decaying, but its
 664 apparent stability seems to suggest that baryon number is conserved [43]. However,
 665 proton decay is a usual feature of grand-unified theories, where electromagnetic, weak
 666 and strong interactions are unified above a certain energy scale [44].

667 As the energy deposition scale for this kind of searches is nearly the same as the one
 668 for long-baseline neutrino oscillations, DUNE will be able to look for them. It has several
 669 advantages over other experiments, such as excellent imaging and particle identification,
 670 which can be translated to lower backgrounds.

671 The last of the main objectives of DUNE is the detection of neutrinos originated in
 672 supernovae explosions, what is called a supernova neutrino burst (SNB). These neutrinos
 673 carry with them information about the core-collapse process, from the progenitor to the

3.3. Far Detector

explosion and the remnant; but also may have information about new exotic physics. So far, the only neutrino events ever recorded from such a process were a few dozens of $\bar{\nu}_e$ events from the 1987A supernova located in the Magellanic Cloud, 50 kpc away from Earth [45, 46].

DUNE aims to collect also some SNB events. Although these are quite rare, as the expected supernovae explosion events are about one every few decades for our galaxy and Andromeda, the long lifetime of the experiment (around a few decades as well) makes it reasonable to expect some. Nowadays the main sensitivity to SNB of most experiments is to the $\bar{\nu}_e$ through inverse beta decay. One of the advantages of DUNE is its expected sensitivity to ν_e , since the dominant channel will be ν_e CC scattering.

Moreover, due to the stringent requirements that the main physics goals set for DUNE, it will allow also to perform searches for all kind of BSM physics. Among others, DUNE will be able to look for: active-sterile neutrino mixing, non-unitarity of the PMNS matrix, non-standard interactions, Lorentz and CPT violations, neutrino trident production, light-mass DM, boosted DM and heavy neutral leptons. The reader is referred to the DUNE FD TDR Volume II [42] for a full discussion of the physics scope of DUNE.

3.3 Far Detector

The so-called DUNE FD complex will sit 1.5 km underground at SURF, South Dakota. Two caverns will host the four FD modules, two of them per cavern, each embedded in cryostats of dimensions 18.9 m (w) \times 17.8 m (h) \times 65.8 m (l). A central, smaller cavern will host the cryogenic system.

Three out of the four modules will be liquid argon (LAr) time projection chamber detectors, often refer to as LArTPCs, with a LAr fiducial mass of at least 10 kt each. The first and second FD modules, FD-1 and FD-2, will use a Horizontal Drift (HD) technology, whereas the third module, FD-3, will have a Vertical Drift (VD) direction. The technology for the fourth module is still to be decided,

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.2: Schematic diagram showing the operating principle of a LArTPC with wire readout. Figure taken from Ref. [1].

For each event, with energies ranging from a few MeV to several GeV, these detectors collect both the scintillation light and the ionisation electrons created when the charged particles produced in neutrino-nucleus interactions ionise the argon nuclei. In both HD and VD designs the characteristic 128 nm scintillation light of argon is collected by a photon detection system (PDS). This light will indicate the time at which electrons start to drift, thus enabling reconstruction over the drift coordinate when compared to the time when the first ionisation electron arrives to the anode. Reconstruction of the topology in the transverse direction is achieved using the charge readout. Fig. 3.2 illustrates the detection principle described, for the case of a HD detector with a wire readout.

3.3.1 Horizontal Drift

Within the HD design the ionisation electrons produced as charged particles traverse the LAr drift horizontally towards the anode planes, made out of three layers of wire readout, due to the effect of an electric field. This design, previously known as single-phase (SP), was tested by the ProtoDUNE-SP detector at CERN. The prototype collected data from

3.3. Far Detector




Figure 3.3: Proposed design for the FD-1 and FD-2 modules following the HD principle. Figure taken from Ref. [1].

716 a hadron beam and cosmic rays, providing high-quality data sets for calibration studies
 717 and proving the excellent performance of this design.

718 Each FD HD detector module is divided in four drift regions, with a maximum drift
 719 length of 3.5 m, by alternating anode and cathode walls. The surrounding field cage
 720 ensures the uniformity of the 500 V/cm horizontal electric field across the drift volumes.
 721 The three anode walls, which constitute the charge readout of the detector, are built by
 722 stacking anode plane assemblies (APAs), 2 high times 25 wide. The design of the HD
 723 modules is shown in Fig. 3.3.

724 Each APA is made of 2560 active wires arranged in three layers, plus an extra grid
 725 layer, wrapped around a metal frame. The two induction wire planes, U and V, sit at
 726 $\pm 35.7^\circ$ to the vertical on each side of the APA. The collection and shielding plane wires,
 727 X and G, run parallel to the vertical direction. The ionisation electrons drift past the
 728 induction planes, generating bipolar signals on those wires, and are collected by the
 729 collection plane, producing a monopolar positive signal. The spacing between the wires
 730 is ~ 5 mm, and it defines the spatial resolution of the APA.

731 The front-end readout electronics, or cold electronics as they are immerse in the LAr,

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.4: Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G. Figure taken from Ref. [1].

732 are attached to the top of the up APAs and the bottom of the down APAs. Mounted on
 733 the front-end mother boards we have a series of ASICs that digitize the signals from the
 734 collection and induction planes. Each wire signal goes to a charge-sensitive amplifier,
 735 then there is a pulse-shaping circuit and this is followed by the analogue-to-digital
 736 converter. This part of the process happens inside the LAr to minimise the number of
 737 cables penetrating the cryostat. The digitised signals come out finally via a series of
 738 high-speed serial links to the warm interface boards (WIBs), from where the data is sent
 739 to the back-end DAQ through optical fibers.




Figure 3.5: A PDS module containing 24 X-ARAPUCAs (left) and the location of the modules on the APAs (right). Figure taken from Ref. [1].

3.3. Far Detector




Figure 3.6: Proposed design for the FD-3 module following the VD principle. Figure adapted from Ref. [47].

The PDS uses modules of X-ARAPUCA devices, mounted on the APA frames between the wire planes. Each X-ARAPUCA consists of layers of dichroic filter and wavelength-shifter. They shift the VUV scintillation light into the visible spectrum, sending then the visible photons to silicon photomultiplier (SiPM) devices. The PDS modules are $209\text{ cm} \times 12\text{ cm} \times 2\text{ cm}$ bars, containing 24 X-ARAPUCAs. There are 10 of these PDS modules per APA. Fig. 3.5 shows a PDS module (left) and the placement of the modules on the APAs (right).

3.3.2 Vertical Drift

In the VD case the ionisation electrons will drift vertically until they meet a printed circuit board-based (PCB) readout plane. It is based on the original dual-phase (DP) design deployed at CERN, known as ProtoDUNE-DP, used a vertical drift design with an additional amplification of the ionization electrons using a gaseous argon (GAr) layer above the liquid phase. The VD module incorporates the positive features of the DP design without the complications of having the LAr-GAr interface.

The current design of the FD VD module counts with two drift chambers with a maximum drift distance of 6.5 cm. A cathode plane splits the detector volume along the

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.7: Schematic representation of the electrode strip configuration for a top (left) and bottom (right) CRU. Figure taken from Ref. [47].

drift direction while the two anode planes are connected to the bottom and top walls of the detector. The layout of the VD module is shown in Fig. 3.6. Compared with the HD design, the VD option offers a slightly larger instrumented volume and a more cost-effective solution for the charge readout.

As in the HD design, each drift volume features a 500 V/cm electric field and a field cage that ensures its uniformity. The anode planes are arrays of $3.4\text{ m} \times 3\text{ m}$ charge-readout planes (CRPs). These are formed by a pair of charge-readout units (CRUs), which are built from two double-sided perforated PCBs, with their perforations aligned. The perforations allow the drift electrons to pass between the layers.

The PCB face opposite to the cathode has a copper guard plane which acts as shielding, while its reverse face is etched with electrode strips forming the first induction plane. The outer PCB has electrode strips on both faces, the ones facing the inner PCB form the second induction plane while the outermost ones form the collection plane. Fig. 3.7 shows the layout of the electrode strips for the top (left) and bottom (right) CRUs.

3.3. Far Detector




Figure 3.8: Detailed diagram of the DUNE FD DAQ system. Figure taken from Ref. [2].

770 The magenta and blue lines represent the first and second induction planes respectively,
 771 and the green lines correspond to the collection plane.

772 The PDS in the VD module will use the same X-ARAPUCA technology developed
 773 for the HD design. The plan is to place the PDS modules on the cryostat walls and on
 774 the cathode, in order to maximise the photon yield.

775 **3.3.3 FD Data Acquisition System**

776 The task of the data acquisition (DAQ) system is to receive, process and store data from
 777 the detector modules. In the case of DUNE the DAQ architecture is designed to work
 778 for all FD modules interchangeably, except some aspects of the upstream part which
 779 may depend on the specific module technology.

780 The enormous sample rate and the number of channels in TPC and PD readouts
 781 will produce a very large volume of data. These pose really strong requirements and
 782 challenges to the DUNE FD DAQ architecture. It will be required to read out data of
 783 the order of ten thousand or more channels at rates of a few MHz. In order to cope
 784 with the huge data volume, segmented readouts and compression algorithms are used to
 785 reduce the data rate to manageable levels.

Chapter 3. The Deep Underground Neutrino Experiment

The DAQ system of the DUNE FD is composed of five different subsystems. The first one is the upstream DAQ, which receives the raw data from the detector, buffers it and perform some low-level pre-processing. The minimally processed data is then fed into a hierarchical data selection system, which then performs a module level trigger decision. In case of a positive decision a trigger command is produced and executed by the data flow orchestrator, located in the back-end (BE) DAQ subsystem. Subsequently the DAQ BE retrieves the relevant data from the buffers located in the upstream DAQ, adds all the data into a cohesive record and saves it to permanent storage. Watching over all the other subsystems we also have the control, configuration and monitoring subsystem and the time and synchronization subsystem. Fig. 3.8 shows a schematic diagram of the DAQ system, showing the different subsystems and their relations.

A notorious challenge for the DUNE DAQ system comes from its broad physics goals. We must be prepared to process events spanning a wide range of time windows (from 5 ms in the case of beam and cosmic neutrinos and nucleon decay to 100 s in the case of SNBs) and therefore this requires a continuous readout of the detector modules. Moreover, because of the off-beam measurements we need to ensure the capabilities of online data processing and self-triggering. Having this into account, together with the technical constraints, the DUNE FD DAQ faces a series of challenges: it needs to be fault tolerant and redundant to reduce downtime, accommodate new components while it keeps serving the operational modules, have large upstream buffers to handle SNB physics, be able to support a wide range of readout windows and last reduce the throughput of data to permanent storage to be at most 30 PB/year.

3.4 Near Detector

In order to estimate the oscillation parameters we measure the neutrino energy spectra at the FD. This reconstructed energy arises from a convolution of the neutrino flux, cross section, detector response and the oscillation probability. Using theoretical and empirical models to account for the other effects, one can extract the oscillation probability using

3.4. Near Detector




Figure 3.9: Representation of the ND hall in Phase II, showing the different subcomponents. From right to left, in the direction of the beam, we have ND-LAr, ND-GAr and SAND. Figure taken from Ref. [48].

813 the measurement. However, these models have associated a number of uncertainties that
 814 are then propagated to the oscillation parameters.

815 One of the main roles of the ND is to measure the neutrino interaction rates before
 816 the oscillation effects become relevant, i.e. close to the production point. By measuring
 817 the ν_μ and ν_e energy spectra, and that of their corresponding antineutrinos, at the ND
 818 we can constrain the model uncertainties. A complete cancellation of the uncertainties
 819 when taking the ratio between the FD and ND measurements is not possible, as that
 820 would require both detectors to have identical designs and the neutrino fluxes to be
 821 the same. Because of the distance, the flux probed by the FD will have a different
 822 energy and flavour composition than that at the ND, as neutrinos oscillate and the beam
 823 spreads. The differences in the flux also determine the design of the detectors, therefore
 824 the ND is limited in its capability to match the FD design.

825 Nevertheless, having a highly capable ND DUNE can minimise the systematic
 826 uncertainties affecting the observed neutrino energy. The ND data can be used to
 827 tune the model parameters by comparison with the prediction. Then, one uses the
 828 tuned model to predict the unoscillated FD spectra. Comparing the prediction with the

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.10: Schematic representation of the external components of ND-LAr, including the cryostat and the PRISM movable system (left) and detailed drawing of one ArgonCube module (right). Figure adapted from Ref. [1].

829 measured spectra it is possible to extract the oscillation parameters.

830 Additionally, the ND will have a physics program of its own. In particular, it will
 831 measure neutrino cross sections that will then be used to constrain the model used in
 832 the long-baseline oscillation analysis. It will also be used to search for BSM phenomena
 833 such as heavy neutral leptons, dark photons, millicharged particles, etc.

834 The DUNE ND can be divided in three main components, a LArTPC known as ND-
 835 LAr, a magnetised muon spectrometer, which will be the Temporary Muon Spectrometer
 836 (TMS) in Phase I and ND-GAr in Phase II, and the System for on-Axis Neutrino
 837 Detection (SAND). The layout of the Phase II DUNE ND can be seen in Fig. 3.9. The
 838 first two components of the ND will be able to move off-axis, in what is called the
 839 Precision Reaction-Independent Spectrum Measurement (PRISM) concept. More details
 840 on the purpose and design of the ND can be found in the DUNE ND Conceptual Design
 841 Report (CDR) [48].

842 3.4.1 ND-LAr

843 ND-LAr is a LArTPC, as the ND needs a LAr component in order to reduce cross
 844 section and detector systematic uncertainties in the oscillation analysis. However, its

3.4. Near Detector




Figure 3.11: Schematic view of the TMS detector, highlighting its main parts. Figure adapted from Ref. [1].

845 design differs significantly from those proposed for the FD modules. Because of the
 846 high event rates at the ND, approximately 55 neutrino interaction events per $10 \mu\text{s}$ spill,
 847 ND-LAr will be built in a modular way. Each of the modules, based on the ArgonCube
 848 technology, is a fully instrumented, optically isolated TPC with a pixelated readout.
 849 The pixelisation allows for a fully 3D reconstruction and the optical isolation reduces
 850 the problems due to overlapping interactions. Fig. 3.10 shows a representation of the
 851 external parts of ND-LAr (left) and a detailed diagram of an ArgonCube module (right).

852 With a fiducial mass of 67 t and dimensions $7 \text{ m (w)} \times 3 \text{ m (h)} \times 5 \text{ m (l)}$, ND-LAr
 853 will be able to provide high statistics and contain the hadronic systems from the beam
 854 neutrino interactions, but muons with a momentum higher than 0.7 GeV will exit the
 855 detector.

856 3.4.2 TMS/ND-GAr

857 In order to accurately estimate the neutrino energy, the momentum of the outgoing
 858 muons needs to be determined. That is the reason why a muon spectrometer is needed
 859 downstream of ND-LAr.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.12: Cross section of the ND-GAr geometry, showing the HPgTPC, ECal and magnet. Figure adapted from Ref. [1].

860 In Phase I that role will be fulfilled by TMS. It is a magnetised sampling calorimeter,
861 with alternating steel and plastic scintillator layers. Fig. 3.11 shows a schematic view of
862 the TMS detector. The magnetic field allows a precise measurement of the sign of the
863 muon, so one can distinguish between neutrino and antineutrino interactions.

864 After the Phase II upgrade, TMS will be replaced with ND-GAr. This detector is
865 a magnetised, high-pressure GAr TPC (often denoted as HPgTPC) surrounded by an
866 electromagnetic calorimeter (ECal) and a muon tagger. A cross section of its geometry
867 can be seen in Fig. 3.12. ND-GAr will be able to measure the momenta of the outgoing
868 muons while also detect neutrino interactions inside the GAr volume. This allows
869 ND-GAr to constrain the systematic uncertainties even further, as it will be able to
870 accurately measure neutrino interactions at low energies thanks to the lower tracking
871 thresholds of GAr.

872 3.4.3 PRISM

873 In general, the observed peak neutrino energy of a neutrino beam decreases as the
874 observation angle with respect to the beam direction increases. This feature has been
875 used in other long-baseline neutrino experiments, like T2K (2.5° off-axis) and NOvA

3.4. Near Detector




Figure 3.13: Predicted beam muon neutrino flux at the ND location for different off-axis positions. Figure taken from Ref. [48].

(0.8° off-axis), in order to achieve narrower energy distributions. The DUNE PRISM concept exploits this effect using a movable ND. Within PRISM both ND-LAr and the muon spectrometer (TMS in Phase I and ND-GAr in Phase II) can be moved up to 3.2° off-axis, equivalent to move the detectors 30.5 m laterally through the ND hall.

This allows to record additional data samples with different energy compositions. Fig. 3.13 compares the on-axis muon neutrino flux at the ND with the fluxes at different off-axis positions. As the off-axis position increases the neutrino flux becomes closer to a monoenergetic beam with a lower peak energy. These samples can be used to perform a data-driven determination of the relation between true and reconstructed neutrino energy, in order to reduce the dependence on the interaction model. The off-axis samples are linearly combined to produce a narrow Gaussian energy distribution centered on a target true energy. From the combination coefficients one can build a sample of reconstructed neutrino events that will determine the energy mapping.

The PRISM samples can also be used to form a flux at the ND location similar in shape to the oscillated flux measured by the FD. This method can be used to extract the oscillation parameters with minimal input from the neutrino interaction model.

Chapter 3. The Deep Underground Neutrino Experiment




Figure 3.14: Schematic longitudinal section of the LBNF beamline at Fermilab (not to scale). Figure taken from Ref. [49].

892 3.4.4 SAND

893 The role of SAND is to monitor the beam stability by measuring the on-axis neutrino
 894 energy spectra. As the PRISM program requires that ND-LAr and its downstream
 895 muon spectrometer spend about half of the time in off-axis positions, it is not possible
 896 to monitor the stability with the movable detectors. Moreover, for the success of PRISM
 897 it is essential to have a stable beam configuration, or, at least, a quick assessment and
 898 modeling of the distortions.

899 The SAND detector is magnetised, and it counts with an inner low density tracker,
 900 a LAr target with optical readout and surrounding sampling calorimeter.

901 3.5 LBNF beamline

902 The Long-Baseline Neutrino Facility (LBNF) project is responsible for producing the
 903 neutrino beam for the DUNE detectors. A detailed discussion of the LBNF program
 904 can be found in the DUNE/LBNF CDR Volume III [49].

905 The LBNF beamline will provide a high-intensity neutrino beam within the adequate
 906 energy range in order to meet the long-baseline oscillation physics goals of DUNE. A
 907 schematic diagram of the longitudinal section of the LBNF beamline is shown in Fig.
 908 3.14. First, a beam of $60 - 120$ GeV protons is extracted from the Fermilab Main

3.5. LBNF beamline

909 Injector. This beam is aimed towards the target area, where it collides with a cylindrical
 910 graphite target to produce pions and kaons.

911 The diffuse, secondary beam of particles is focused by a pair of magnetic horns.
 912 These select the positively charged particles when operated in Forward Horn Current
 913 (FHC) mode, or the negatively charged ones when the current is reversed, also known as
 914 Reverse Horn Current (RHC) mode. The focused secondary beam then enters a 194 m
 915 decay pipe where the pions and kaons will predominantly produce $\mu^+\nu_\mu$ pairs when in
 916 FHC mode (or $\mu^-\bar{\nu}_\mu$ in RHC mode).

917 At the end of the decay pipe a hadron absorber removes the undecayed hadrons and
 918 muons from the beam, which reduces the ν_e ($\bar{\nu}_e$) and $\bar{\nu}_\mu$ (ν_μ) contamination coming
 919 from the μ^+ (μ^-) decays. The resulting neutrino flux at the FD is shown in Fig. 3.15,
 920 both for FHC (left) and RHC (right) modes. These predictions show the intrinsic $(\bar{\nu}_e)$
 921 contamination and wrong sign component from wrong sign and neutral meson decays,
 922 as well as muons decaying before reaching the absorber.




Figure 3.15: Predicted neutrino fluxes at the FD in FHC mode (left panel) and RHC mode (right panel). Figures taken from Ref. [42].

₉₂₃ **Chapter 4**

₉₂₄ **ND-GAr**

₉₂₅ ND-GAr is a magnetised, high-pressure gaseous argon TPC (HPgTPC), surrounded by
₉₂₆ an electromagnetic calorimeter (ECal) and a muon detector (commonly refer to as μ ID).
₉₂₇ A detailed discussion on the requirements, design, performance and physics of ND-GAr
₉₂₈ can be found in the DUNE ND CDR [48] and the ND-GAr whitepaper (cite).

₉₂₉ In DUNE Phase II ND-GAr will fulfill the role of TMS, measuring the momentum
₉₃₀ and sign of the charged particles exiting ND-LAr. Additionally, it will be able to measure
₉₃₁ neutrino interactions inside the HPgTPC, achieving lower energy thresholds than those
₉₃₂ of the ND and FD LArTPCs. By doing so ND-GAr will allow to constrain the relevant
₉₃₃ systematic uncertainties for the LBL analysis even further.

₉₃₄ The goal of the present chapter is to review the requirements that the physics program
₉₃₅ of DUNE impose on ND-GAr, present the current status of its design and describe the
₉₃₆ GArSoft package, its simulation and reconstruction software.

₉₃₇ **4.1 Requirements**

₉₃₈ The primary requirement for ND-GAr is to the measure the momentum and charge of
₉₃₉ muons from ν_μ and $\bar{\nu}_\mu$ CC interactions in ND-LAr, in order to measure their energy
₉₄₀ spectrum. To achieve the sensitivity to the neutrino oscillation parameters described
₉₄₁ in the DUNE FD TDR Volume II [42] ND-GAr should be able to constrain the muon

Chapter 4. ND-GAr

942 energy within a 1% uncertainty or better. The main constraint will come from the
943 calibration of the magnetic field, performed using neutral kaon decays in the HPgTPC.

944 Another requirement for ND-GAr is the precise measurement of neutrino interactions
945 on argon for the energies relevant to the neutrino oscillation program. The goal is to
946 constrain the cross section systematic uncertainties in the regions of phase space that
947 are not accessible to ND-LAr. This requires the kinematic acceptance for muons in
948 ND-GAr to exceed that of ND-LAr, being comparable to the one observed in the FD.

949 ND-GAr should also be able to the relationship between true and reconstructed energy
950 from neutrino interactions on argon with low thresholds, being sensitive to particles that
951 are not observed or may be misidentified in ND-LAr. In particular, ND-GAr needs to
952 have low tracking thresholds in order to measure the spectrum of pions and protons
953 produced in final-state interactions (FSI). It also must be able to accurately measure
954 the pion multiplicity in 1, 2 and 3 pions final states, to inform the pion mass correction
955 in the LArTPCs.

956 4.2 Reference design

957 The final design of ND-GAr is still under preparation. However, a preliminary baseline
958 design was in place at the time of the ND CDR. This section summarises the main
959 features of that design, as it is also the one used for the default geometry in our simulation.
960 A DUNE Phase II whitepaper, discussing the different options under consideration for
961 the ND-GAr design, is in progress.

962 4.2.1 HPgTPC

963 The reference design for the ND-GAr HPgTPC follow closely that of the ALICE TPC.
964 It is a cylinder with a central high-voltage cathode, generating the electric field for
965 the two drift volumes, with a maximum drift distance of 2.5 m each. The anodes will
966 be instrumented with charge readout chambers. The original design repurposed the
967 multi-wire proportional readout chambers of ALICE, however the current R&D efforts

4.2. Reference design




Figure 4.1: Diagram of the ALICE TPC, showing the two drift chambers, inner and outer field cages and readout chambers. Figure taken from Ref. [1].

968 focus on a gas electron multiplier option instead. Fig. 4.1 shows a schematic diagram of
 969 the ALICE TPC design. The basic ND-GAr geometry will resemble this, except for the
 970 inner field cage.

971 It will use a 90-10 molar fraction argon-CH₄ mixture at 10 bar. With this baseline
 972 gas mixture light collection is not possible, as the quenching gas absorbs most of the
 973 VUV photons. Additional R&D efforts are underway, to understand if different mixtures
 974 allow for the light signal to be used to provide a t_0 while maintaining stable charge gain.

975 4.2.2 ECal

976 The main role of the ND-GAr ECal is the calorimetric measurement of the electron
 977 energies and the reconstruction of photons, in particular those from neutral pion decays.
 978 Also, the ECal is able to provide a t_0 timestamp for neutrino interactions, by associating
 979 its activity to the tracks in the HPgTPC. The ECal will also be able to perform
 980 neutron reconstruction using time of flight and reject external backgrounds, thanks to
 981 its sub-nanosecond time resolution.

982 The ECal design features three independent subdetectors, two end caps at each side
 983 and a barrel surrounding the HPgTPC. Each of the detectors is divided in modules,

Chapter 4. ND-GAr




Figure 4.2: View of the 12-sided ECal barrel and outer muon tagger geometries (left) and layout of the ECal tile layers for the 2 mm Cu, 10 mm scintillator option (right). Figure adapted from Ref. [1].

which combine alternating layers of plastic scintillator and absorber material readout by SiPMs. The inner scintillator layers consist of $2.5 \times 2.5 \text{ cm}^2$ high-granularity tiles, whereas the outer ones are made out of 4 cm wide cross-strips spanning the whole module length. The current barrel geometry consists of 8 tile layers and 34 strip layers, while the end caps feature 6 and 36 respectively. The thickness of the scintillator layers is 7 mm and 5 mm for the Pb absorber layers. The 12-sided geometry of the ECal barrel (left) and the layout of the tile layers (left)¹ can be seen in Fig. 4.2.

4.2.3 Magnet

The ND-GAr magnet design, known as the Solenoid with Partial Yoke (SPY), consists of two coupled solenoids with an iron return yoke. The idea behind the design is to have a solenoid as thin as possible, as well as a return yoke mass distribution that minimises the material budget between ND-LAr and ND-GAr. The magnet needs to provide a 0.5 T field in the direction perpendicular to the beam, parallel to the drift electric field. It needs to host the pressure vessel and the surrounding ECal, which points to an inner diameter of $\sim 6.4 \text{ m}$.

The solenoid is a single layer coil, based on niobium titanium superconducting

¹The figure shows the layout of the tile layers for a previous design with 2 mm Cu absorber and 10 mm plastic scintillator, as mentioned in the text the current choice is 5 mm Pb absorber and 7 mm scintillator.

4.3. GArSoft

1000 Rutherford cable. The total length of the coil is 7.5 m. The bobbin will be split in four
1001 segments grouped in pairs with two identical cryostats, connected in series. The iron
1002 yoke features an aperture in the upstream side to allow the muons coming from ND-LAr.
1003 Still, its material will be enough to reduce the magnetic field reaching SAND, and also
1004 stop the charged pions produced inside the HPgTPC.

1005 4.2.4 Muon system

1006 The design of the ND-GAr muon system is still in a preliminary stage. Its role is to
1007 distinguish between muons and pions punching through the ECal. This is especially
1008 important for wrong-sign determination, to separate these from neutral current events.

1009 In its current form, the muon system consists of three layers of longitudinal sampling
1010 structures. It alternates 10 cm Fe absorber slabs with 2 cm plastic scintillator strips.
1011 The transverse granularity required is still under study.

1012 4.3 GArSoft

1013 GArSoft is a software package developed for the simulation and reconstruction of events
1014 in ND-GAr. It is inspired by the LArSoft toolkit used for the simulation of LArTPC
1015 experiments, like the DUNE FD modules. It is based on `art`, the framework for event
1016 processing in particle physics experiments [?]. Other of its main dependencies are `ROOT`,
1017 `NuTools`, `GENIE` and `Geant4`. It allows the user to run all the steps of a generation-
1018 simulation-reconstruction workflow using FHiCL configuration files.

1019 4.3.1 Event generation

1020 The standard generator FHiCLs in GArSoft run the event generation and particle
1021 propagation simulation (i.e. Geant4) in the same job by default. However, it is possible
1022 to split them up if needed. The current version of GArSoft provides five different event
1023 generators, each of them producing `simb::MCTruth` products defined in `NuTools`. The
1024 available modules are:

Chapter 4. ND-GAr

- 1025 • **SingleGen**: particle gun generator. It produces the specified particles with a given
1026 distribution of momenta, initial positions and angles.
 - 1027 • **TextGen**: text file generator. The input file must follow the `hepevt` format², the
1028 module simply copies this to `simb::MCTruth` data products.
 - 1029 • **GENIEGen**: GENIE neutrino event generator. The module runs the neutrino-nucleus
1030 interaction generator using the options specified in the driver FHiCL file (flux file,
1031 flavour composition, number of interactions per event, t_0 distribution, ...). Current
1032 default version is `v3_04_00`.
 - 1033 • **RadioGen**: radiological generator. It produces a set list of particles to model
1034 radiological decays. Not tested.
 - 1035 • **CRYGen**: cosmic ray generator. The module runs the CRY event generator with a
1036 configuration specified in the FHiCL file (latitude and altitude of detector, energy
1037 threshold, ...). Not tested.
- 1038 The module `GArG4` searches for all the generated `simb::MCTruth` data products, using
1039 them as inputs to the Geant4 simulation with the specified detector geometry. A constant
1040 0.5 T magnetic field along the drift coordinate is assumed. The main outputs of this step
1041 are `simb::MCParticle` objects for the generated Geant4 particles, `gar::EnergyDeposit`
1042 data products for the energy deposits in the HPgTPC and `gar::CaloDeposit` data
1043 products for the energy deposits in the ECal and muon system.

1044 4.3.2 Detector simulation

- 1045 The standard detector simulation step in GArSoft is all run with a single FHiCL, but
1046 the different modules can be run independently as well. First the `IonizationReadout`

²In brief, each event contains at least two lines. The first line contains two entries, the event number and the number of particles in the event. Each following line contains 15 entries to describe each particle. The entries are: status code, pdg code for the particle, entry of the first mother for this particle, entry of the second mother for this particle, entry of the first daughter for this particle, entry of the second daughter for this particle, x component of the particle momentum, y component of the particle momentum, z component of the particle momentum, energy of the particle, mass of the particle, x component of the particle initial position, y component of the particle initial position, z component of the particle initial position and time of the particle production.

4.3. GArSoft

1047 module simulates the charge readout of the HPgTPC, and later the `SiPMReadout` module
1048 runs twice, once for the ECal and then for the muon system, with different configurations.

1049 The `IonizationAndScintillation` module collects all the `gar::EnergyDeposit`
1050 data products, to compute the equivalent number of ionization electrons for each energy
1051 deposit. The `ElectronDriftAlg` module simulates the electron diffusion numerically
1052 both in the longitudinal and transverse directions and applies an electron lifetime
1053 correction factor. The induced charge on the nearest and neighbouring readout pads
1054 is modeled using the provided pad response functions. The digitisation of the data is
1055 then simulated with the `TPCReadoutSimAlg` module. By default, the ADC sampling
1056 rate used is 50.505 MHz. The resulting raw waveforms for each channel are stored with
1057 zero-suppression, in order to save memory and CPU time. The algorithms keep blocks
1058 of ADC values above a certain threshold, plus some adjustable additional early and late
1059 tick counts. The results of these three steps are `gar::raw::RawDigit` data products.

1060 For the ECal and the muon system the `SiPMReadout` module calls either the
1061 `ECALReadoutSimStandardAlg` or `MuIDReadoutSimStandardAlg` modules. These take
1062 all the `gar::CaloDeposit` data products in the corresponding detector and do the
1063 digitisation depending on whether the hit was in a tile or strip layer. They include single
1064 photon statistics, electronic noise, SiPM saturation and time smearing. The resulting
1065 objects are `gar::raw::CaloRawDigit` data products.

1066 4.3.3 Reconstruction

1067 The reconstruction in GArSoft is also run as a single job by default. It first runs the hit
1068 finding, clustering, track fitting and vertex identification in the HPgTPC, followed by
1069 the hit finding and clustering in the ECal and muon system. After those it produces the
1070 associations between the associations between the tracks and the ECal clusters.

1071 Focusing first on the HPgTPC reconstruction, the `CompressedHitFinder` module
1072 takes the zero-suppressed ADCs from the `gar::raw::RawDigit` data products. The
1073 reconstructed hits largely correspond to the above threshold blocks, however the hit
1074 finder identifies waveforms with more than one maximum, diving them in multiple hits

Chapter 4. ND-GAr

1075 if they dip below a certain threshold. The data products produced are of the form
1076 `gar::rec::Hit`. These are the inputs to the clustering of hits in the `TPCHitCluster`
1077 module. Hits close in space and time are merged, and the resulting centroids are found.
1078 This module outputs `gar::rec::TPCClusters` objects and associations to the input
1079 hits.

1080 The following step prior to the track fitting is pattern recognition. The module
1081 called `tpcvecchitfinder2` uses the `gar::rec::TPCClusters` data products to find track
1082 segments, typically called vector hits. They are identified by performing linear 2D fits
1083 to the positions of the clusters in a 10 cm radius, one fit for each coordinate pair. A
1084 3D fit defines the line segment of the vector hit, using as independent variable the one
1085 whose sum of (absolute value) slopes in the 2D fits is the smallest. The clusters are
1086 merged to a given vector hit if they are less than 2 cm away from the line segment. The
1087 outputs are `gar::rec::VecHit` data products, as well as associations to the clusters. The
1088 `tpcpatrec2` module takes the `gar::rec::VecHit` objects to form the track candidates.
1089 The vector hits are merged together if their direction matches, their centers are within
1090 60 cm and their direction vectors point roughly to their respective centers. Once
1091 the clusters of vector hits are formed they are used to make a first estimation of the
1092 track parameters, simply taking three clusters along the track. The module produces
1093 `gar::rec::Track` data products and associations between these tracks and the clusters
1094 and vector hits.

1095 The track is fitted by means of a Kalman filter in the `tpctrackfit2` module, using
1096 the position along the drift direction as the independent variable. Two different fits are
1097 performed per track, a forward and a backwards fit, each starting from one of the track
1098 ends. The Kalman filter state vector ($y, z, R, \phi, \tan\lambda$) is estimated at each point along
1099 the track using a Bayesian update. The track parameters reported in the forward and
1100 backwards fits are the ones computed at the opposite end where the fit started. The
1101 main outputs of the track fit are the `gar::rec::Track` objects. Additionally, the module
1102 stores the fitted 3D positions along the track in the `gar::rec::TrackTrajectory` data
1103 products and the total charge and step sizes for each point also get stored in the form of

4.3. GArSoft

1104 `gar::rec::TrackIonization` objects.

1105 After the tracking step, the `vertexfinder1` module looks at the reconstructed
1106 `gar::rec::Track` products, creating vertex candidates with the track ends that are
1107 within 12 cm of each other. The vertices are then fitted using linear extrapolations from
1108 the different track ends associated. The results are `gar::rec::Vertex` data products,
1109 and associations to the tracks and corresponding track ends.

1110 For the ECal and muon tagger, the `SiPMHitFinder` module runs twice with different
1111 configurations, adapted to the particular capabilities of both. The module simply takes
1112 the `gar::raw::CaloRawDigit` products, applies a calibration factor to convert the ADC
1113 counts to MeV and for the strip layer hits it calculates the position along the strip using
1114 the times recorded of both SiPMs. This module produces `gar::rec::CaloHit` data
1115 products. Next, these objects are used as inputs to the `CaloClustering` module. It
1116 merges the hits based on a simple nearest neighbours (NN) algorithm. For the resulting
1117 clusters it also computes the total energy and position of the centroid. The results are
1118 stored as `gar::rec::Cluster` data products, with associations to the hits.

1119 The last step in the reconstruction is associating the reconstructed tracks in the
1120 HPgTPC to the clusters formed in the ECal and muon system. The `TPCECALAssociation`
1121 module checks first the position of the track end points, considering only the points
1122 that are at least 215 cm away from the cathode or have a radial distance to the center
1123 greater than 230 cm. The candidates are propagated up to the radial position, in the
1124 case of clusters in the barrel, or the drift coordinate position, for the end cap cluster, of
1125 the different clusters in the collection using the track parameters computed at the end
1126 point. The end point is associated to the cluster if certain proximity criteria are met.
1127 This module creates associations between the tracks, the end points and the clusters.
1128 The criteria for the associations are slightly different for the ECal and the muon tagger.

₁₁₂₉ Chapter 5

₁₁₃₀ FWTPG offline software

₁₁₃₁ Chapter 6

₁₁₃₂ Matched Filter approach to
₁₁₃₃ induction wire Trigger Primitives

₁₁₃₄ 6.1 Motivation

₁₁₃₅ The filter implemented in the firmware of the upstream DUNE FD DAQ is a 32nd-order
₁₁₃₆ low-pass finite impulse-response (FIR) filter. The output of such filter for a discrete
₁₁₃₇ system can be written as:

$$y[i] = \sum_{j=0}^N h[j]x[i-j], \quad (6.1)$$

₁₁₃₈ where N is the order of the filter, y is the output sequence, x is the input sequence and h
₁₁₃₉ is the set of coefficients of the filter. The current implementation within `dtp-firmware`
₁₁₄₀ [50] uses a set of 16 non-zero integer coefficients.

₁₁₄₁ Filtering is a vital step in the hit finder chain. It helps to suppress the noise and
₁₁₄₂ enhance the signal peaks with respect to the noiseless baseline. A good filtering strategy
₁₁₄₃ allows us to use lower thresholds when forming the trigger primitives (TPs) and thus
₁₁₄₄ increasing the sensitivity of our detector to low energy physics events. In such events,
₁₁₄₅ the hits produced by the ionisation electrons tend to have lower amplitudes than those
₁₁₄₆ of interest to the baseline physics programme of the DUNE experiment.

₁₁₄₇ This is particularly important for the induction planes. In general, signal peaks in

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.1: Schematic representation of an APA. The black lines represent the APA steel frame. The green and magenta lines correspond to the direction of the U and V induction wires respectively. The blue lines indicate the direction of the X collection wires and the wire shielding G.

the induction wires have smaller amplitude than the ones in the induction plane. This, together with the fact that the pulse shapes are bipolar, reduces our capacity to detect the hits on these channels. The inefficiency of detecting TPs in the induction planes (denoted as U and V planes) lead trigger algorithms to focus mainly on the TPs from the collection plane (so-called X plane). As a result, the possibility of making trigger decisions based on the coincidence of TPs across the three wire planes remains nowadays unexploited in DUNE. Fig. 6.1 shows a schematic view of an anode plane assembly (APA), with the different wire plane orientations highlighted.

A possible improvement of the current hit finder chain could require optimising the existing or choosing a new filter implementation. A filter strategy which improves the induction signals may be able to enhance the detection efficiency of TPs from the induction planes and ideally make it comparable to that of the collection plane.

The goal is to implement a better finite-impulse response filter design and to evaluate its performance relative to the current filter. To do so, we need to take into account the limitations of the firmware: the FIR filter shall have maximum 32 coefficients (so-called taps) whose values are 12-bit unsigned integers. Although it is technically possible to include non-integer coefficients, it would be a technical challenge as we have 40 FIR

6.2. Signal-to-noise ratio definition




Figure 6.2: Left panel: Zoomed unfiltered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44` (blue line). The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840, where the green shaded region represents $\pm \sigma_{\text{noise}}$.

1165 instances per APA, as there are 4 FIR per optical link and 10 optical links per APA.
 1166 With these restrictions, the task is to provide a set of 32 coefficients which yield an
 1167 optimal filter performance for the induction wires.

1168 6.2 Signal-to-noise ratio definition

1169 I introduce the signal to noise ratio (S/N) as a measure of the FIR filter performance
 1170 and demonstrate how to extract its value for a set of ProtoDUNE-SP data. The S/N
 1171 metrics allow us to compare different filter implementations and serve as a basis for more
 1172 detailed studies presented later in this document. Specifically, I use the ADC capture
 1173 `felix-2020-07-17-21:31:44` (data capture taken for firmware validation purposes). I
 1174 defined S/N as the height of the signal peaks relative to the size of the noise peaks.
 1175 To quantify this quantity channel by channel one first need to estimate the standard
 1176 deviation of the ADC data for each channel, σ_{ADC} . Then, I define the corresponding

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

noise waveform to be the ADC values in the range $\pm 3\sigma_{ADC}$. From this new noise data one can estimate again the mean and standard deviation, μ_{noise} and σ_{noise} , so I can write the S/N for any given channel as:

$$S/N = \frac{\max [ADC] - \mu_{noise}}{\sigma_{noise}}, \quad (6.2)$$

where $\max [ADC]$ is simply the maximum ADC value found in the corresponding channel.

One can apply this definition of the S/N with a waveform from one of the channels of the data capture¹. Fig. 6.2 shows a zoomed region of the waveform corresponding to channel 7840 (blue line), where one can clearly see three signal peaks and continuous additive noise (we actually see 6 peaks, 3 positive and 3 negative, but, because by design for induction channels the expected signal pulse shapes are bipolar, I treat them as a collection of 3 individual signal peaks). I estimated the standard deviation of this raw waveform to be $\sigma_{raw} = 7.84$ ADC, so I am able to define the noise waveform (red line) as the ADC values in the range ± 23.52 ADC. This way one obtains $\mu_{noise} = 0$ and $\sigma_{noise} = 5.58$ ADC, which gives $S/N = 30.84$.

We can repeat this calculation now for the corresponding filtered waveform (using the current firmware FIR filter). In Fig. 6.3 I plotted the same time window for the filtered waveform from channel 7840 (blue line). In this case, the standard deviation of the waveform is larger than before, giving $\sigma_{raw} = 10.99$ ADC. The resulting noise waveform (red line) results from selection the ADC values in the range ± 32.91 ADC, giving now $\mu_{noise} = -0.47$ ADC and $\sigma_{noise} = 7.03$ ADC. Finally, one obtains $S/N = 24.68$. Notice that the value of S/N decreases after the filtering. Clearly, one can see that the noise baseline has increased by a factor of 1.35 when we applied the FIR filter and at the same time the amplitude of the signal peaks has remained almost unchanged, leading to this poorer S/N value.

¹All the original work was done within the `dtp-simulation` package [51], which offers a variety of tools to read raw data and emulate the TPG block (pedestal subtraction, filtering and hit finder). However, the results shown in this report were re-worked later using the C++ based `dtpemulator` package [52]. Its main purpose is the emulation of the TPG block and, in the same way as its predecessor, it has been cross-checked against the current firmware implementation.

6.3. Low-pass FIR filter design




Figure 6.3: Left panel: Zoomed filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44` (blue line). The filter used was the current implementation of the low-pass FIR filter in `dtp-firmware`. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$

1200 6.3 Low-pass FIR filter design

1201 In general, when one uses a method to optimize the frequency response of a digital filter,
 1202 such as the Parks-McClellan algorithm, one finds a set of N real coefficients that give
 1203 the best response for the specified pass-band and order of the filter [53].

1204 In our case, as the sampling frequency is defined as 1 ticks^{-1} , the Nyquist frequency
 1205 will simply be $1/2 \text{ ticks}^{-1}$. The current implementation of the filter seems to have as
 1206 pass-band the range $[0, 0.1] \text{ ticks}^{-1}$. This can be seen in Fig. 6.4, where I show the
 1207 power spectrum, in decibels, of such filter implementation (blue solid line). For instance,
 1208 the Park-McClellan algorithm finds the optimal Chebyshev FIR filter taking as input
 1209 the boundaries of the target pass-band and stop-band, which can be written in the form:

$$\left\{ \begin{array}{l} [0, f_c] \\ [f_c + \delta f, f_N] \end{array} \right. , \quad (6.3)$$

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

where f_c is the cut-off frequency, δf is the transition width and f_N is the aforementioned Nyquist frequency. A similar behaviour to the one in the current filter can be obtained by setting $f_c = 0$ and $\delta f = 0.1 \text{ ticks}^{-1}$. The response of the resulting filter is also shown in Fig. 6.4 (blue solid line). Notice that the suppression of the stop-band is enhanced for this optimal filter. For comparison I included the power response of the filter obtained by taking the integer part of the coefficients resulting from the Parks-McClellan method (red dashed line). One can see that it does not suppress that much the stop-band, in a similar way to the current implementation of the filter.




Figure 6.4: Power spectrum in decibels for the current implementation of the low-pass FIR filter in `dtp-firmware` (blue line), compared to the response of an optimal filter obtained using the Parks-McClellan algorithm for the same pass-band (red line). Also for comparison I include the spectrum of the optimal filter when taking only the integer part of the coefficients (red dashed line).

At this point, I tried to improve the performance of the FIR filter using the Parks-McClellan method, i.e. maximize the overall S/N, using the available data captures. I did so by varying the values of the two quantities that parametrize the pass-band and stop-band, the cut-off frequency f_c and the transition width δf .

Fig. 6.5 shows the average relative change in the S/N (i.e. the ratio between the value of the S/N after and before the filtering) for capture `felix-2020-07-17-21:31:44`,

6.3. Low-pass FIR filter design




Figure 6.5: Relative change in the S/N for the ProtoDUNE-SP raw data capture `felix-2020-07-17-21:31:44`, using different values of the cutoff frequency f_c and the transition width δf . The optimal Chebyshev filters were applied using just the integer part of the coefficients given by the Parks-McClellan algorithm.

when using filters designed with the Parks-McClellan algorithm for the specified values of the cut-off frequency f_c and the transition width δf , restricted to integer values for the filter coefficients. One can clearly distinguish different regions where we get an improvement of up to a factor of 1.35 for the U plane. For large values of $f_c + \delta f$ the ratio tends to 1, as expected (in that limit the width of the stop-band goes to 0, meaning that no frequencies are filtered out and thus the waveform remains the same).

Using the configuration which gives the best mean performance for the three planes (see bottom right panel of Fig. 6.5), i.e. $f_c = 0.068 \text{ ticks}^{-1}$ and $\delta f = 0.010 \text{ ticks}^{-1}$, we can see how such filter affects the different channels. Fig. 6.6 shows the distribution of the S/N improvement values for all the channels in the raw ADC capture `felix-2020-07-17-21:31:44`, separated by wire plane, after the optimal Chebyshev filter was applied. One can see that there is a clear improvement for both U and V induction wire planes, obtaining a mean change of 1.25 and 1.30 for them respectively. However, in the case of the collection plane X the mean of this distribution is roughly 1, meaning that a good fraction of channels in that plane get a slightly worse S/N after the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.6: Distribution of the relative change of the S/N on the different wire planes from the ProtoDUNE-SP raw data capture *felix-2020-07-17-21:31:44* after the optimal Chebyshev filter was applied. The filter was computed with the Parks-McClellan algorithm using a cutoff of $f_c = 0.068 \text{ ticks}^{-1}$ and a transition width $\delta f = 0.010 \text{ ticks}^{-1}$.

1239 filter is applied. In any case, this is not a big issue as the S/N for collection channels is
 1240 usually much higher than the one for induction channels.

1241 The results I obtained optimising the low pass filter with the Parks-McClellan method
 1242 are promising. Nonetheless, the improvement found is rather marginal so I wondered
 1243 if there could be an alternative approach to the filtering problem which yields better
 1244 outputs. At this point, I found a possible alternative in matched filters. By construction,
 1245 this kind of filters offer the best improvement on the S/N.

1246 6.4 Matched filters

1247 In the context of signal processing, a matched filter is the optimal linear filter for
 1248 maximising the signal-to-noise ratio (S/N) in the presence of additive noise, obtained
 1249 by convolving a conjugated time-reversed known template with an unknown signal to
 1250 detect the presence of the template in the signal [54].

1251 Given a known signal sequence $s(t)$ and another (a priori unknown) noise sequence

6.4. Matched filters




Figure 6.7: Left panel: Zoomed match filtered waveform corresponding to channel 7840 from the ProtoDUNE-SP raw data capture felix-2020-07-17-21:31:44 (blue line). The filter used was directly extracted from the data, being the 32 values around the first peak in the original waveform. The green dashed lines mark the region $\pm 3\sigma_{\text{raw}}$. The resulting noise waveform is also shown (red line). Top right panel: ADC distribution for channel 7840 after match filtering, where the green shaded region represents $\pm \sigma_{\text{raw}}$. Bottom right panel: noise ADC distribution for channel 7840 after match filtering, where the green shaded region represents $\pm \sigma_{\text{noise}}$

1252 $n(t)$, the input signal can be written as:

$$x(t) = s(t) + n(t). \quad (6.4)$$

1253 Now, considering a linear time-invariant filter, whose impulse-response function I
1254 will refer to as $h(t)$, one can write the output signal as:

$$\begin{aligned} y(t) &= x(t) * h(t) \\ &= (s(t) + n(t)) * h(t) \\ &= y_s(t) + y_n(t), \end{aligned} \quad (6.5)$$

1255 where $y_s(t)$ and $y_n(t)$ are simply the outputs of the filter due to the signal and the noise
1256 components respectively.

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1257 The goal of the matched filter is to detect the presence of the signal $s(t)$ in the input
 1258 sample $x(t)$ at a certain time t_0 , which effectively means we need to maximise the S/N.
 1259 This way, what one wants is to have a filter which gives a much bigger output when the
 1260 known signal is present than when it is not. Putting it in other words, the instantaneous
 1261 power of the signal output $y_s(t)$ should be much larger than the average power of the
 1262 noise output $y_n(t)$ at some time t_0 .

1263 For the case of the filtered signal, one can easily re-write it as an inverse Fourier
 1264 transform:

$$y_s(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega H(\omega)S(\omega)e^{i\omega t}, \quad (6.6)$$

1265 where $H(\omega)$ and $S(\omega)$ are the Fourier transforms of the impulse-response function (i.e.
 1266 the transfer function of the filter) and of the input signal, respectively.

1267 Now focusing on the noise, we can use the Wiener-Khinchin theorem [55] to write
 1268 the mean power of the noise after filtering as:

$$E|y_n(t)|^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega |H(\omega)|^2 S_n(\omega), \quad (6.7)$$

1269 where $S_n(\omega)$ is the power spectral density of the noise.

1270 Having these, one can write the instantaneous S/N at time t_0 as:

$$\begin{aligned} \left(\frac{S}{N} \right)_{t_0} &= \frac{|y_s|^2}{E|y_n(t)|^2} \\ &= \frac{1}{2\pi} \frac{\left| \int_{-\infty}^{\infty} d\omega H(\omega)S(\omega)e^{i\omega t_0} \right|^2}{\int_{-\infty}^{\infty} d\omega |H(\omega)|^2 S_n(\omega)}. \end{aligned} \quad (6.8)$$

1271 Once we have this expression, we need to find the upper limit of it to determine what
 1272 would be the optimal choice for the transfer function. One can use the Cauchy-Schwarz
 1273 inequality, which in the present case takes the form:

$$\left| \int_{-\infty}^{\infty} dx f(x)g(x) \right|^2 \leq \int_{-\infty}^{\infty} dx |f(x)|^2 + \int_{-\infty}^{\infty} dx |g(x)|^2, \quad (6.9)$$

6.4. Matched filters

1274 for any two analytical functions $f(x)$ and $g(x)$. One can prove that making the choice:

$$\begin{aligned} f(x) &= H(\omega) \sqrt{S_n(\omega)} e^{i\omega t_0}, \\ g(x) &= \frac{S(\omega)}{\sqrt{S_n(\omega)}}, \end{aligned} \quad (6.10)$$

1275 leads to the following upper bound for the S/N:

$$\left(\frac{S}{N} \right)_{t_0} \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{|S(\omega)|^2}{S_n(\omega)}. \quad (6.11)$$

1276 From Eqs. (6.8), (6.9) and (6.10) one can also derive the form of the transfer function

1277 such that the upper bound is exactly reached [56]:

$$H(\omega) \propto \frac{S^*(\omega) e^{-i\omega t_0}}{S_n(\omega)}. \quad (6.12)$$

1278 From this last expression we can clearly see the way the matched filter acts. As the

1279 transfer function is proportional to the Fourier transform of the signal it will try to only

1280 pick the frequencies present in the signal [57].

1281 The matched filter transfer function can be greatly simplified if the input noise is
 1282 Gaussian. In that case, the power spectral density of the noise is a constant, so it can be
 1283 re-absorbed in the overall normalisation of the transfer function. Moreover, considering
 1284 that the input signal is a real function, one can simply set $S^*(\omega) = S(-\omega)$, which gives:

$$H(\omega) \propto S(-\omega) e^{-i\omega t_0}. \quad (6.13)$$

1285 For a discrete signal, one can think of the input and impulse-response sequences as

1286 vectors of \mathbb{R}^N . Then, the matched filter tries to maximise the inner product of the signal

1287 and the filter while minimising the output due to the noise by choosing a filter vector

1288 orthogonal to the later. In the case of additive noise, that leads to the impulse-response

1289 vector:

$$h = \frac{1}{\sqrt{s^\dagger R_n^{-1} s}} R_n^{-1} s, \quad (6.14)$$

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.8: Relative improvement in the S/N for the raw data capture `felix-2020-07-17-21:31:44`, using the matched filter following the parametrisation in Eq. (6.17). The black crosses in both panels denote the location of the maximum ratio value.

where s is a reversed signal template sequence of length N equal to the order of the filter and R_n is the covariance matrix associated with the noise sequence n . For the Gaussian noise case, the covariance matrix is simply the unit matrix, so the above expression simplifies again to:

$$h = \frac{s}{|s|}. \quad (6.15)$$

For this first stage of the study, I use a definition of the S/N per channel given by:

$$\text{S/N} = \frac{\max [ADC] - \mu_{noise}}{\sigma_{noise}}, \quad (6.16)$$

where the subscript *noise* refers to a subset of the data obtained by only taking into account waveform values within a $\pm 3\sigma$ range around the mean of the data and $\max [ADC]$ is the maximum of the original waveform. This definition is further discussed in App. 6.2, where I also show examples of its application to raw data and to a waveform filtered with the current low-pass FIR filter.

To test whether this choice of filter is appropriate one needs to choose a signal template. As an example of how a matched filter would affect our signal, I simply took the filter coefficients to be the 32 ADC values around a signal peak present in the data.

6.4. Matched filters

1303 In Fig. 6.7 (left panel) I plotted a zoomed region for channel 7840 in the raw data capture
1304 `felix-2020-07-17-21:31:44`, after applying the matched filter described before (blue
1305 line). When compared to the raw and FIR filtered case (see App. 6.2), after applying
1306 the match filter the standard deviation of the noise waveform (red line) decreases and at
1307 the same time the signal peaks are enhanced. This leads to an improvement of the S/N
1308 by a factor of 1.92 when compared to the raw waveform.

1309 In order to obtain the matched filter that is more suitable for our data, I explored
1310 different configurations of signal templates. In order to perform this exploration, I
1311 parametrised the signal using the bipolar function:

$$f(x) = -A(x + \delta) e^{-x^2/\sigma^2}, \quad (6.17)$$

1312 where the parameter δ controls the asymmetry between the positive and negative peaks
1313 and σ controls their width. The amplitude parameter A is set such that it keeps the
1314 height of the biggest peak to be less than 200 ADC in absolute value.

1315 As this parametrisation is only adequate for bipolar signals I will focus exclusively
1316 on the induction channels. Also, the optimal configurations I found for the U and V
1317 plane will be kept separate, i.e. I will have two sets of coefficients that will be applied to
1318 either the U and V planes of wires. I do so as I found this was the choice giving the
1319 best performance. Even so, as I will discuss, the differences are not very pronounced. In
1320 case it is not technically possible to separate channels in the firmware according to the
1321 wire plane they come from and use different sets of filter coefficients for them, we can
1322 just find a common unique set of coefficients. In such case, I do not expect our results
1323 to change dramatically.

1324 In Fig. 6.8 I present the results of our parameter scan, for channels in the induction
1325 planes U (left panel) and V (right panel). For each configuration of σ and δ the resulting
1326 matched filter was applied to all channels in the corresponding plane within the data
1327 capture `felix-2020-07-17-21:31:44`, the S/N improvement was computed with respect
1328 to the raw waveforms and then the S/N mean value was kept as a score for such filter.

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1329 One can see that the improvement obtained for the U plane is in general higher than the
1330 one for the V plane. In any case, I got substantially higher ratios than the ones obtained
1331 for the low-pass FIR filters. For the optimal configurations I attained improvements up
1332 to a factor of 1.85 for the U plane and 1.65 for the V plane.

1333 The sets of optimal matched filter coefficients were obtained for the parameters
1334 $\delta = 0.035$, $\sigma = 0.191$ for the U plane and $\delta = 0.018$, $\sigma = 0.191$ for the V plane. I
1335 show these two sets of coefficients in Fig. 6.9 (left panel). Also in Fig. 6.9 (right
1336 panel) I plot the distribution of the S/N improvement after the optimal match filters
1337 for the U and V were applied to the corresponding channels in the raw data capture
1338 `felix-2020-07-17-21:31:44`. As mentioned before, the mean improvement achieved
1339 for the U plane channels is slightly bigger than the one for the V channels. Note, however,
1340 that the spread of the distribution for the V plane is also smaller than the one for the U
1341 plane.

1342 I also performed a similar scan for the case of a low-pass FIR filter using the Parks-
1343 McClellan algorithm. In that case, the parameters to check were the cutoff frequency
1344 and the transition width of the filter. A summary of the results is given in App. 6.3.

1345 Overall, one can see that the improvements on the S/N are much more significant in
1346 the case of the matched filter than it is for the low-pass FIR filters. The analysis of this
1347 and other raw data captures from ProtoDUNE-SP suggest that matched filters increase
1348 the S/N of induction channels by a factor of 1.5 more than the optimal low-pass FIR
1349 filters.

1350 Although these results are by themselves great points in favour of the matched
1351 filter, more studies are needed to completely assess the robustness of this approach. I
1352 proceeded then to test the matched filter with simulated data samples.

1353 6.5 Using simulated samples

1354 In order to further test the matched filter, the next step was to generate and process
1355 data samples using *LArSoft* [58]. In this way, one can control the particle content of

6.5. Using simulated samples




Figure 6.9: Left panel: Optimal matched filter coefficients for the U (blue line) and V (red line) planes. The filters were computed with our parametrisation in Eq. (6.17) for the parameter values $\delta = 0.035$, $\sigma = 0.191$ and $\delta = 0.018$, $\sigma = 0.191$ respectively. Right panel: Distribution of the relative change of the S/N on the two induction wire planes from the ProtoDUNE-SP raw data capture *felix-2020-07-17-21:31:44* after their respective optimal matched filters were applied.

1356 the samples, the orientation of the tracks and their energy, and therefore see how the
1357 matched filter behaves in various situations.

1358 To begin with, I prepared different monoenergetic and isotropic samples containing
1359 a single particle per event. Each sample contains a different particle species, namely
1360 electrons, muons, protons and neutral pions all with a kinetic energy of $E_k = 100$ MeV.
1361 I chose these because of the fairly different topologies they generate in the liquid argon,
1362 ranging from shower-like to track-like. The procedure I followed to generate the samples
1363 and process them is discussed in detail in App. ??.

1364 These were generated with the single particle gun and the Geant4 stage of the
1365 *LArSoft* simulation [58] was performed with the standard configuration for the DUNE
1366 FD 10kt module.

1367 For simplicity, I restricted the particles to start drifting in a single TPC volume
1368 (in this case TPC 0), so I can focus exclusively on the signals coming from one APA.
1369 The chosen kinetic energy for all the particles in my first trial is $E_k = 100$ MeV, so a
1370 necessary check is to see if all our tracks will be typically contained in one TPC volume.
1371 Fig. 6.10 (left panel) shows the distributions of the track lengths in the liquid argon

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.10: Left panel: distributions of the particles track length in the liquid argon for the generated $E_k = 100$ MeV monoenergetic samples, electrons (blue), muons (red), protons (green) and neutral pions (purple). Right panel: distribution of the length of the longest photon in the neutral pion sample after the decay process $\pi^0 \rightarrow \gamma\gamma$.

of all generated particles with $E_k = 100$ MeV. One can see that, in the case of the track-like particles (i.e. muons and protons), their length distributions are quite sharp and centered at relatively low distances (30 and 8 cm, respectively). For electrons, the distribution is quite broad but it does not extend past ~ 30 cm. The case of neutral pions can be misleading, as they decay promptly the track length associated with the true Monte Carlo particle is always < 1 cm. In Fig. 6.10 (right panel) I show the effective length distribution of the longest photon after the pion decays as $\pi^0 \rightarrow \gamma\gamma$, highlighting the number of fully contained photons in the TPC volume per event (either zero, one or both). One can see that the vast majority of events has both photons contained and that just a negligible number of them has none of them contained in the TPC volume. In any case, for the sake of caution, I will only keep the pion events with both photons contained.

Once I have prepared a sample at the Geant4 level, I need to process it through the detector simulation. In order to make adequate estimations of the noise levels and run the filtering and hit finder as I did with the ProtoDUNE data, one needs to turn off the default zero-suppression of the waveforms produced by the simulation. At this first stage I am only concerned with the waveforms with the noise added, so I keep the noise

6.5. Using simulated samples

addition option as true in the configuration. However, for studies related to the hit finder performance one will also need to store the noiseless waveforms in order to retrieve the truth information of the hits. I will discuss this approach next.

After the detector simulation stage, one needs to extract the no zero-suppressed noisy waveforms, along with their offline channel numbers, and store them in a certain format to be analysed later. To reduce the amount of data that will go for processing, I used the information from the Geant4 step of the simulation to select only the active channels, i.e. the channels where some ionisation electrons arrive. Moreover, as said previously, I only extract the waveforms from APA 0 and exclusively the ones coming from induction channels. The resulting ROOT file contains a tree with two branches, one containing the waveforms for each event and channel and the other with the corresponding offline channel numbers.

Finally, to extract the truth values for the orientation of the tracks and the energies of the particles I used a modified analysis module. This gives a ROOT file with a single tree, containing several branches with different information such as the components of the initial momentum of the particles, initial and final xyz location, track length, etc.

For the analysis of the resulting waveforms and truth values I used a custom set of Python libraries (available at [??]). Among other functionalities, these enable the user to read the ROOT files, export the raw data as pandas objects, apply the filters and compute the S/N of both the raw and filtered signals. So far, the default configuration for the filtering uses the set of optimal matched filter coefficients that I found using the ProtoDUNE data samples.

Additionally, for the analysis of the samples it was necessary to use two different reference frames, to study separately the signals coming from the U and V induction wire planes. As I am focussing on a single APA, the U and V wires have a different orientation in the yz plane. In the case of U wires, these are tilted 35.7° clockwise from the vertical (y direction), whereas the V wires are at the same angle but in the counter clockwise direction. Because of this, the best option is to deal with two new coordinate systems rotated by $\pm 35.7^\circ$ along the x axis, so the new y' and y'' directions are aligned with the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.11: Left panel: schematic representation of the two new rotated reference frames used in this analysis (denoted as prime and double prime), viewed from the yz plane. The magenta stack of lines represent the wires in the U plane, whereas the green lines correspond to the wires in the V plane. Right panel: 3D representation of the momentum of one of the generated monoenergetic muons (red arrow) in the original reference frame (black lines), along with the new reference frame used for the U plane waveforms (blue lines). In the yz plane I added the projection of these three.

1418 U and V induction wires. Fig. 6.11 (left panel) shows a schematic representation of
 1419 the original reference frame together with the two rotated ones (denoted by primed and
 1420 double primed). This way, one can easily understand how parallel was a track to the
 1421 wires in the two induction planes. Fig. 6.11 (right panel) shows a 3D representation of
 1422 the momentum of a track (red arrow) in the original reference frame (black lines), along
 1423 with the new reference frame for U wires (blue lines). I added the projection in the yz
 1424 plane of this three, to show the usefulness of the new reference frame to tell whether a
 1425 track is parallel or normal to the wires in the induction plane.

1426 Fig. 6.12 shows the distribution of the average S/N improvement per event when one
 1427 applies the optimal matched filters. I produced separate distributions for the channels
 1428 in the U (red) and V (blue) induction wire planes. Notice that the S/N distributions
 1429 for the track-like particles, i.e. muons (top left panel) and protons (bottom left panel),
 1430 have significantly larger mean values than the distributions of the shower like particles,
 1431 i.e. electrons (top right panel) and neutral pions (bottom right panel). An important

6.5. Using simulated samples




Figure 6.12: Distributions of the mean S/N improvement per event for the corresponding sample after applying the matched filters. Here I separated the change in the U plane (blue) and the V plane (red) channels. From top left to the right: muon, electron, proton and neutral pion. All the events have a fixed kinetic energy of $E_k = 100 \text{ MeV}$.

difference between these results and the ones seen before for ProtoDUNE data is that, overall, the improvements that I get for simulated data are bigger. This could be due either to the default noise model used in the *LArSoft* simulation or to the simulated hits having higher energy than the ones in the recorded data. Nonetheless, the concluding message is that the previously optimised matched filters give an overall significant improvement of the S/N for the different samples.

About the convention I followed for the plots and results, in the case of the raw and filtered S/N of each event in the sample I simply took the average of the quantities over all the active channels in the event. That is, if a certain event has N_{chan} active channels these two quantities are computed as:

$$(S/N_{fir})_{event} = \frac{\sum_{i=0}^{N_{chan}} (S/N_{fir})_i}{N_{chan}},$$

$$(S/N_{raw})_{event} = \frac{\sum_{i=0}^{N_{chan}} (S/N_{raw})_i}{N_{chan}}.$$
(6.18)

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.13: Angular dependence of the mean S/N and the S/N improvement, for the different monoenergetic samples considered (from top to bottom: electrons, muons, protons and neutral pions). The two columns on the left represent the values for the U plane waveforms. The top subplots show the mean S/N for raw (green) and filtered (red) waveforms whereas the bottom subplots depict the averaged S/N improvement (black).

1442 However, for the ratio of the raw and filtered S/N (what I called the S/N improvement)
 1443 per event I am not just taking the ratio of the previous two quantities but computing
 1444 the average of the individual ratios per channel in the event:

$$\left(\frac{S/N_{fir}}{S/N_{raw}} \right)_{event} = \frac{\sum_{i=0}^{N_{chan}} \left(\frac{S/N_{fir}}{S/N_{raw}} \right)_i}{N_{chan}}, \quad (6.19)$$

1445 and so:

$$\left(\frac{S/N_{fir}}{S/N_{raw}} \right)_{event} \neq \frac{(S/N_{fir})_{event}}{(S/N_{raw})_{event}}. \quad (6.20)$$

1446 6.5.1 Angular dependence

1447 Having these monoenergetic samples, one can also study the angular dependence of the
 1448 performance of the matched filter. This is an important point, as it is a well established
 1449 fact that for certain configurations (an extreme case configuration being signals normal

6.5. Using simulated samples




Figure 6.14: Angular dependence of the mean S/N and the S/N improvement, for the different monoenergetic samples considered (from top to bottom: electrons, muons, protons and neutral pions). The two columns on the left represent the values for the U plane waveforms. The top subplots show the mean S/N for raw (green) and filtered (red) waveforms whereas the bottom subplots depict the averaged S/N improvement (black).

to the wire plane and perpendicular to the induction wires at the same time) the S/N is much lower than average as the corresponding waveforms are severely distorted. In this sense, I am interested to see how the matched filter behaves for these cases and how the S/N improvement on those compare to the average.

Fig. 6.13 shows the angular dependence of the S/N for the monoenergetic $E_k = 100$ MeV isotropic muons, for the different induction wire planes and projections. The angles for each event are given by the components of the initial value of the momentum of the particles, taking the angles of the projections on the xz and yz planes with respect to the z axis (more accurately, one needs to compute these angles twice for each event, a pair for the $xy'z'$ coordinate system and the other for the $xy''z''$). The top row shows the dependence on the angles corresponding to the U plane, i.e. $\theta_{xz'}$ and $\theta_{y'z'}$, whereas the bottom row shows the angular dependence viewed from the V plane, $\theta_{xz''}$ and $\theta_{y''z''}$. In each plot, the top subplot represents the mean values of the S/N for the raw (blue) and

Chapter 6. Matched Filter approach to induction wire Trigger Primitives

1463 matched filtered (red) signals, and the bottom subplot the averaged S/N improvement
1464 (black). The solid lines represent the mean value obtained for the corresponding angular
1465 value, whereas the semitransparent bands represent one standard deviation around the
1466 mean at each point.

1467 As expected, the S/N is in general higher when tracks are parallel to the APA (i.e.
1468 $\theta_{xz} \sim 0$) and lower when it is normal to the plane ($\theta_{xz} \sim \pm 90^\circ$). In the same way, tracks
1469 parallel to the wires ($\theta_{yz} \sim \pm 90^\circ$) tend to have higher S/N than those perpendicular to
1470 these ($\theta_{yz} \sim \pm 0$).

1471 Fig. 6.14 shows the corresponding angular dependence information for the $E_k =$
1472 100 MeV electrons sample. Notice that, in this case, the S/N behaviour discussed above
1473 does not hold. A possible explanation can be that, because most hits in these events
1474 are produced by the secondary particles generated in the EM shower, the signal peaks
1475 whose S/N ratios were computed do not correspond to the directional information of
1476 the primary electron.

1477 6.5.2 Distortion and peak asymmetry

1478 As a little case of study, I selected two of the simulated $E_k = 100$ MeV monoenergetic
1479 muon events. With respect to the U induction plane, one is parallel to the APA (low
1480 $\theta_{xz'}$) and to the wires (high $\theta_{y'z'}$) and the other is normal to the APA plane (high $\theta_{xz'}$)
1481 and perpendicular to the wires (low $\theta_{y'z'}$). As expected from the results on the angular
1482 dependence discussed above, the former has a higher S/N (before and after the filtering)
1483 when compared to the latter. An interesting thing to notice about these two samples
1484 is that, even though one has a much bigger S/N than the other, it is the one with the
1485 smallest S/N the one that got the biggest averaged S/N improvement. In Table 6.1
1486 I included all the relevant parameters of these two $E_k = 100$ MeV muon events I am
1487 considering, namely, the angles with respect to the $xy'z'$ reference frame, the values of
1488 the S/N, the S/N improvement and also the so-called peak asymmetry Δ_{peak} that I will
1489 discuss next.

1490 One can try to understand better what is going on with these two events by looking

6.5. Using simulated samples




Figure 6.15: Selected consecutive waveforms corresponding to two monoenergetic $E_k = 100$ MeV muon events, one is parallel to the APA and to the wires in the U plane (left panel) and the other is normal to the APA plane and perpendicular to the U plane wires (right panel). The solid lines represent the raw waveforms whereas the dashed lines correspond to the waveforms after the matched filter was applied. The waveforms on the left panel have been scaled by a factor of 0.15 to have similar amplitude to the ones on the right panel.

Table 6.1: Characteristic parameters of the two monoenergetic muon events selected, relative to the U plane: projected angles in the xz' and $y'z'$ planes, S/N values for the raw and filtered waveforms, mean improvement of the S/N and peak asymmetry.

	$\theta_{xz'}$ (°)	$\theta_{y'z'}$ (°)	S/N _{raw}	S/N _{MF}	$\frac{S/N_{MF}}{S/N_{raw}}$	Δ_{peak} (ADC)
High ("parallel")	-1.58	-77.76	41.65	112.44	2.83	-35.73
Low ("normal")	76.92	-2.56	8.07	25.46	3.12	-10.38

at the raw and filtered data from some of their active channels. Fig. 6.15 shows a selection of consecutive raw and filtered U plane waveforms from the event with high S/N (left panel) and the one with low S/N (right panel). Notice that to show both collections of waveforms at a similar scale I had to apply a factor of 0.15 to the waveforms with high S/N. Additionally, next to each waveform I included the values of the raw and

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.16: Left panel: peak asymmetry distribution for the case of the monoenergetic $E_k = 100$ MeV muon sample. Each value corresponds to a single bipolar signal peak from a channel in any event. The blue distribution represents the peaks on U plane channels, whereas the red corresponds to signal peaks in V wires. Right panel: relation between the mean peak asymmetry per event with the S/N for U channel waveforms from the $E_k = 100$ MeV muon sample. The top subplot shows the decimal logarithm of the mean S/N for the raw (red) and the matched filtered (blue) waveforms. The bottom subplot contains the mean S/N improvement ratio after the matched filter was applied.

1496 matched filtered S/N for the corresponding channel. The first thing to notice in this plot
 1497 is that the amplitude of the signal peaks from the normal track have a much smaller
 1498 amplitude, and also appear quite distorted when compared to the others. On the other
 1499 hand, although the matched filtered S/N is still smaller, the relative improvement is
 1500 bigger than in the parallel case.

1501 A way I found to quantify the difference between the shapes within these two events
 1502 is their different peak asymmetry. One can define the peak asymmetry as the (signed)
 1503 difference between the positive and the negative peaks of the bipolar shape, i.e.:

$$\Delta_{peak} \equiv h_+ - h_-, \quad (6.21)$$

1504 where both heights h_+ and h_- are positive defined. Fig. 6.16 (left panel) shows the
 1505 distribution of this peak asymmetry for all the waveforms corresponding to channels
 1506 in the U (blue) and V (red) planes for the monoenergetic muon sample. One can
 1507 see that these distributions are clearly shifted to negative values (with mean values

6.5. Using simulated samples

1508 $\mu_{\Delta}^U = -10.57$ ADC and $\mu_{\Delta}^V = -5.72$ ADC respectively). It is interesting to notice
1509 that the peak asymmetry value of the sample with high S/N sits at the left tail of the
1510 distribution whereas the corresponding value of the sample with low S/N lies around
1511 the mean.

1512 Now, one can try to correlate the peak asymmetry with the S/N and the S/N change
1513 per event. Fig. 6.16 (right panel) shows the result of comparing (minus) the mean
1514 peak asymmetry per event to the averaged raw (red) and matched filtered (blue) S/N
1515 per event (top subplot). The horizontal lines sit at the mean value obtained in the fit
1516 and represent the width of the $-\Delta_{peak}$ bins used, while the vertical lines indicate one
1517 standard deviation around that mean value. Notice that, when taking decimal logarithm
1518 on both, there is an approximate linear relation between these quantities, except for
1519 peak asymmetry values bigger than -5 ADC where the S/N remains constant.

1520 Also, in the bottom subplot of Fig. 6.16 (right panel) I show the relation between
1521 the peak asymmetry and the mean S/N improvement. In this case, one see that there is
1522 a maximum at $\Delta_{peak} \sim -10$ ADC. As mentioned previously, this is also the value of the
1523 mean of the peak asymmetry distribution. In fact, it is expected that our filter favours
1524 the signal peaks with the most common values of the peak asymmetry, as this was one
1525 of the features I target in our filter coefficient optimisation through the parameter δ .

1526 These results suggest that events with poorer values of the mean S/N, usually
1527 associated to non-favourable track orientations, tend to have smaller values of the mean
1528 peak asymmetry (in absolute value). Nonetheless, because our matched filters have
1529 been optimised to account for these asymmetries, the improvement on the S/N for these
1530 events is sizeable if not better than the one for events which already had a high S/N.

1531 6.5.3 Hit sensitivity

1532 One of the advantages of the matched filter, directly related to increasing the S/N, is
1533 the capability of picking hits that before fell below the threshold. For instance, Fig. 6.17
1534 shows the raw ADC data from an example event (electron, $E_k = 100$ MeV) with the
1535 produced true hits superimposed (black boxes), together with the hits produced by the

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.17: Raw data display in the plane time (in firmware ticks) vs. offline channel number for an $E_k = 100$ MeV electron event. The produced true hits are superimposed (black boxes) as well as the hits coming from the standard hit finder chain (blue circles) and the hit finder using the matched filter (green triangles).

standard hit finder chain (blue circles), i.e. using the current FIR filter, and the hits obtained using the matched filters (green triangles). Both the standard and the matched filter hit finders were run with a threshold of 10 ADC. Notice that the standard hits match well the true ones at the initial part of the event (where we have a track-like object), but they miss most of the hits produced by the EM shower at later times. On the other hand, the hits produced with the matched filter have a better agreement with the true hits even for the more diffuse shower activity.

Notwithstanding that now I get more hits with this combination of matched filter and low threshold as a results of the enhancement of the signal peaks relative to the noise level, it is also true that I pick some spurious hits not related to any real activity if one lowers the thresholds too much. Therefore, some optimisation of the threshold is needed. Basically one will need to make a trade-off between precision and sensitivity.

Having this in mind, I tried to compare the produced hits one gets from the standard hit finder and the ones resulting from applying the matched filter with the true hits.

6.5. Using simulated samples

1550 By running the hit finders on our samples with different values of the threshold one
1551 can understand, for instance, how low one can set the threshold without getting mostly
1552 spurious hits and then evaluate the gains obtained from this.

1553 Because now I am also interested in seeing how the hit sensitivity changes with the
1554 energy, I prepared new isotropic samples with the same types of particles as before
1555 (muons, electrons, protons and neutral pions) but with a flat kinetic energy distribution
1556 ranging from 5 to 100 MeV.

1557 In order to estimate the hit sensitivity, given a certain sample, one needs to recover
1558 the set of true hits to be able to compare these with the ones produced. To do so,
1559 a modification in the procedure I was using to extract the raw waveforms is needed.
1560 For this kind of study I run the detector simulation in two steps, first I produce the
1561 waveforms without noise and extract them in the same format I used for the raw data,
1562 then the noise is added and the noisy waveforms are then written to a file as well.

1563 To have a better comparison between the true hits and the ones produced from
1564 the raw waveforms after applying the two filters, I applied also the FIR filter and the
1565 matched filters to the noiseless waveforms and then I run the hit finder with a minimal
1566 threshold (in this case I used 1 ADC) on these noiseless filtered waveforms. In this way
1567 I generated two sets of true hits, I will refer to them as standard true hits (with the
1568 current/default FIR filter) and matched filter true hits respectively. This allows a more
1569 precise matching between the different groups of hits produced, as it will account for
1570 any delays and distortions introduced by the FIR and the matched filters.

1571 In the case of the raw waveforms (with noise), I run the hit finder on them, with
1572 different values of the threshold, after applying either the FIR or the matched filters. I
1573 will name them simply standard hits and matched filter hits respectively. Then, I match
1574 the generated hits to the true hits (the standard hits with the standard true hits and
1575 the matched filter hits with the matched filter true hits). The matching is performed by
1576 comparing the channel number and the timestamp of the hits. To count as a match,
1577 I require that all hits with the same channel number and timestamp have overlapping
1578 hit windows, i.e. the time windows between their hit end and hit start times need to

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.18: Dependence of the precision (blue), sensitivity (red) and F_1 (green) scores on the threshold values used in the hit finder, for the FIR (left panel) and matched filter (right panel) cases. The results were obtained after matching the hits to the true hits in the case of the isotropic muon sample with kinetic energy in the range 5 to 100 MeV, taking only into account the induction plane channels. The points represent the mean value while the error bars indicate one standard deviation around that mean value.

overlap. If more than one hit in one of the groups have hit overlap with the same hit in the other group I only count the hit with closer hit peak time value.

The generation of the samples, the procedure to produce the standard hits (with the default FIR filter) and matched filter hits and the matching of these with the true hits is described in detail in App. ??.

To quantify the performance of the two hit finder approaches, I use a classical method from statistical classification known as confusion matrix [59]. This is basically a way of sorting the outputs of a binary classifier, considering the true values of the classification and the predicted values. It divides the outputs in four categories: true positive (TP, both true and predicted values are 1), false negative (FN, true value is 1 but predicted is 0), false positive (FP, true value is 0 but predicted is 1) and true negative (TN, both true and predicted values are 0)).

The contents of the confusion matrix allow us to compute other derived scores to judge the performance of our classifiers. In this study, I will make use of three of these

6.5. Using simulated samples

1593 metrics, namely the precision or positive predictive value:

$$\text{PPV} = \frac{\text{TP}}{\text{TP} + \text{FP}}, \quad (6.22)$$

1594 the sensitivity or true positive rate:

$$\text{TPR} = \frac{\text{TP}}{\text{TP} + \text{FN}}, \quad (6.23)$$

1595 and the F_1 score [60]:

$$F_1 = \frac{2\text{TP}}{2\text{TP} + \text{FP} + \text{FN}}, \quad (6.24)$$

1596 which is the harmonic mean of the precision and the sensitivity.

1597 In our specific case I am not going to make use of the true negative value, as its
1598 definition in this context can be ambiguous because one does not have clear instances in
1599 the classification process. This way, I will only count the number of true positives as the
1600 total amount of hits I can match between true and raw populations, the number of false
1601 negatives will be the number of missing true hits and the false positive the number of
1602 hits which do not match any true hit.

1603 In Fig. 6.18 I show the precision (blue), sensitivity (red) and F_1 (green) scores I
1604 obtained for different values of the threshold used in the hit finder for the case of the
1605 muon sample. Because the matched filters are only applied to induction channels, I only
1606 consider here hits coming from the U and V planes. The panel on the left corresponds
1607 to the scores I got when I ran the hit finder on the FIR filtered waveforms, whereas the
1608 right panel contains the scores for the matched filter case. The points are centered at
1609 the threshold value used and represent the mean value obtained for each score using all
1610 the generated events, while the error bars indicate one standard deviation around the
1611 mean value.

1612 One can see that the precision for the matched filter case is lower when the thresholds
1613 are very low, as the noise baseline is slightly amplified, but then rises to high values
1614 quicker than for the FIR case. The other difference one can spot is that the sensitivity

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.19: Dependence of the averaged hit sensitivity on the kinetic energy of the events for the matched filter (blue) and standard (red) hits, for the case of the muon (left panel) and electron (right panel) samples, separated between U (top plots) and V (bottom plots) induction wire planes. The top subplots contain the hit sensitivities for the two hit finder alternatives, while the bottom subplots show the ratio between the two. The horizontal lines sit at the mean value and represent the size of the energy bins, while the vertical error bars indicate one standard deviation around that mean value.

in the FIR case starts dropping faster at around the same threshold values where the precision stabilizes around 1, while in contrast for the matched filter this rapid decrease starts at higher threshold values. A similar scan for the same thresholds was performed for the electron sample in the same energy range, yielding similar results.

In Fig. 6.19 I show the averaged hit sensitivity versus the kinetic energy of the events, both for the matched filter hits (blue) and the standard hits (red). The left panel corresponds to the muon sample, whereas the one on the right corresponds to the electron sample, both with kinetic energies between 5 and 100 MeV. In each panel the top plot corresponds to hits in the U plane, while the bottom plot contains the same information for the V plane. Each plot contains two subplots, the one on the top shows the hit sensitivity values for the matched filter and standard hits separate, while the bottom subplot depicts the ratio between the matched filter and standard sensitivities.

6.5. Using simulated samples




Figure 6.20: Distributions of the hit sensitivity in the U (top panels) and V (bottom panels) planes versus the hit sensitivity in the X plane, both for the standard hits (left panels) and the matched filter hits (right panels), in the case of the electron sample and a threshold of 30 ADC.

1627 The horizontal lines are placed at the mean value obtained in the fit and represent the
 1628 width of the E_k bins used, while the vertical error bars indicate one standard deviation
 1629 around that mean value. In both cases the threshold used was 30 ADC, as I required
 1630 the precision to be higher than 0.99 for both matched filter and standard cases.

1631 One can see that, in general, the improvements are better for the U than for the V
 1632 plane. While for the U channels I achieved a mean improvement of 50% and 80% for
 1633 muons and electrons respectively, the improvement in the V plane is stalled at 10% and
 1634 25%. Nevertheless, if I look at the sensitivities for the matched filter hits in both planes
 1635 one can see these have similar mean values for each energy bin, while on the contrary

Chapter 6. Matched Filter approach to induction wire Trigger Primitives




Figure 6.21: Top panels: standard residual plots of the hit sensitivities between the X and U planes. Bottom panels: quantile-quantile plots of the hit sensitivity standard residuals between the X and U planes. In all cases, the left panel corresponds to the standard hits while the right panel represents the matched filter case, all from the electron sample with a 30 ADC threshold.

for the standard hits the sensitivity remains relatively high for the V plane. This way, it looks there was a less significant gain because the hit sensitivity was already high.

Another interesting observation is the different behaviors for muons and electrons. While hit sensitivity for muons grows significantly with energy, in the case of electrons this slightly decreases the higher the kinetic energy of the event is. In any case, when it comes to the improvement on the sensitivities, this remains almost constant in all cases.

Furthermore, we can look at how the concurrence of hits between the different wire planes has changed. For any given event, I expect to have a similar number of hits in the three planes. As the ionisation electrons need to cross the U and V planes prior to reach the collection plane X they will induce current in those wire planes. A way to check the concurrence of hits across planes is looking at the relation between the hit sensitivities for each individual event. One cannot expect the sensitivities to be exactly equal across

6.5. Using simulated samples

1648 planes, but ideally they should be normally distributed around the diagonal.

1649 Fig. 6.20 shows the hit sensitivity in the U (top panels) and V (bottom panels)
1650 planes versus the hit sensitivity in the X plane, for the case of the standard hits (left
1651 panels) and the matched filter hits (right panels). All plots were generated for the
1652 electron sample and a threshold of 30 ADC. From these one can see a clear trend,
1653 when I use the standard hit finder chain the sensitivities in the induction planes are
1654 systematically lower than the hit sensitivity in the X plane, i.e. most of the points sit
1655 below the diagonal (red dashed line). In contrast, when the matched filters are applied,
1656 the majority of the events are distributed around the diagonal. This points out that the
1657 concurrence of hits across planes has improved.

1658 To exemplify the improvement I obtained, one can consider the residuals of the hit
1659 sensitivities for the X and U planes. Assuming the diagonal hypothesis, i.e. given a
1660 dataset of the form (x, y) for any x I take the predicted y value to be equal to the value
1661 of x , I can compute the standard residuals for the hit sensitivities in U given the ones for
1662 X. In Fig. 6.21 (top panels) I show these standard residuals against the corresponding
1663 values of the hit sensitivity in the U plane, for our electron sample with kinetic energy
1664 between 5 and 100 MeV. If I compare the scatter points in the case of the standard
1665 hits (left panel) and the matched filter hits (right panel), I see that the residuals of the
1666 standard hit finder case follow a certain pattern and their mean deviates from 0.

1667 To see clearly if the residuals are normally distributed, in Fig. 6.21 (bottom panels)
1668 I plot the corresponding quantile-quantile plot for both the standard (left panel) and
1669 matched filter (right panel) standard residuals. One can clearly see that the points for
1670 the standard case follow a strongly non-linear pattern, suggesting that the residuals
1671 do not follow a normal distribution. In contrast, for the matched filter hits the points
1672 conform to a roughly linear path, implying that in this case the normality condition is
1673 fulfilled.

1674 All these results hint at the fact that the concurrence of hits across the wire planes
1675 can be strengthened by applying the matched filters.

1676 **Chapter 7**

1677 **DM searches with neutrinos from
1678 the Sun**

1679 **7.1 Motivation**

1680 The idea of detecting neutrino signals coming from the Sun’s core to probe DM is not new.
1681 The main focus of these searches has usually been high-energy neutrinos originated from
1682 DM annihilations into heavy particles [61–64], although recent studies have proposed to
1683 look at the low-energy neutrino flux arising from the decay of light mesons at rest in the
1684 Sun [65–68] previously thought undetectable.

1685 In this chapter I try to demonstrate the capability of DUNE to constrain different
1686 DM scenarios. I used the neutrino fluxes arising from DM annihilations in the core
1687 of the Sun to compute the projected limits that DUNE would be able to set on the
1688 annihilation rates in the Sun and the DM scattering cross sections.

1689 **7.2 Gravitational capture of DM by the Sun**

1690 The Sun and the centre of the Earth are possible sources of DM annihilations, specially
1691 interesting because of their proximity. Their gravitational attraction ensured the capture
1692 of DM from the local halo through repeated scatterings of DM particles crossing them.

Chapter 7. DM searches with neutrinos from the Sun

1693 Only neutrinos produced from DM annihilations can escape the dense interior of these
1694 objects. Therefore, neutrino telescopes are the most useful experimental layouts to
1695 pursue DM searches from their cores.

1696 The neutrino flux from DM annihilations inside the Sun depends on the DM capture
1697 rate, which is proportional to the DM scattering cross section, and the annihilation rate,
1698 which is proportional to the velocity-averaged DM annihilation cross-section. The total
1699 number of DM particles inside the Sun follows the Boltzmann equation [65]:

$$\frac{dN_{DM}}{dt} = C_{\odot} - A_{\odot} N_{DM}^2, \quad (7.1)$$

1700 where C_{\odot} and A_{\odot} are the total Sun DM capture and annihilation rates respectively.
1701 In this expression I neglected the evaporation term, proportional to N_{DM} , which only
1702 contribute for $m_{DM} \lesssim 4$ GeV [69]. As the current threshold of neutrino telescopes is
1703 a few GeV, this region falls below the probed range but can be important in future
1704 low-energy projects.

1705 This equation has an equilibrium solution:

$$N_{DM}^{eq} = \sqrt{\frac{C_{\odot}}{A_{\odot}}}, \quad (7.2)$$

1706 which represents the amount of DM inside the Sun if the capture and annihilation have
1707 reached equilibrium. As the Sun is approximately 4.6 Gyr old, it is usually assumed that
1708 equilibrium has been achieved. Therefore, the anomalous neutrino flux from the Sun
1709 would only depend on the DM scattering cross section, enabling us to set limits on this
1710 quantity. If one does not assume equilibrium, some assumptions on the DM annihilation
1711 cross section are necessary to extract predictions from neutrino signals.

1712 Here, I am going to consider three possible scenarios for the DM interactions: DM
1713 scattering off electrons, spin-dependent (SD) and spin-independent interactions off nuclei.
1714 For the case of these last two, the cross sections will be given in terms of the SD and
1715 SI elastic scattering DM cross section off protons (assuming that DM interactions off

7.2. Gravitational capture of DM by the Sun

1716 protons and neutrons are identical), σ_p^{SD} and σ_p^{SI} , as [4, 65]:

$$\sigma_i^{\text{SD}} = \left(\frac{\tilde{\mu}_{A_i}}{\tilde{\mu}_p} \right)^2 \frac{4(J_i + 1)}{3J_i} |\langle S_{p,i} \rangle + \langle S_{n,i} \rangle|^2 \sigma_p^{\text{SD}}, \quad (7.3)$$

$$\sigma_i^{\text{SI}} = \left(\frac{\tilde{\mu}_{A_i}}{\tilde{\mu}_p} \right)^2 A_i^2 \sigma_p^{\text{SI}}, \quad (7.4)$$

1717 where $\tilde{\mu}_{A_i}$ is the reduced mass of the DM-nucleus i system, $\tilde{\mu}_p$ is the reduced mass of
1718 the DM-proton system, A_i and J_i the mass number and total angular momentum of
1719 nucleus i and $\langle S_{p,i} \rangle$ and $\langle S_{n,i} \rangle$ the expectation value of the spins of protons and neutrons
1720 averaged over all nucleons, respectively (see Ref. [70] for a review on spin expectation
1721 values).

1722 Since the Sun is mainly composed of Hydrogen, the capture of DM from the halo
1723 is expected to occur mainly through spin-dependent scattering. However, since the
1724 spin-independent cross section is proportional to the square of the atomic mass, heavy
1725 elements can contribute to the capture rate (even though they constitute less than 2%
1726 of the mass of the Sun). Heavy elements can also contribute to the spin-dependent cross
1727 section if the DM has also momentum-dependent interactions.

1728 DM particles can get captured by the Sun if after repeated scatterings off solar
1729 targets their final velocity is lower than the escape velocity of the Sun. In the limit of
1730 weak cross sections, this capture rate can be approximately written as [4]:

$$C_{\odot}^{\text{weak}} = \sum_i \int_0^{R_{\odot}} dr 4\pi r^2 \int_0^{\infty} du_{\chi} \frac{\rho_{\chi}}{m_{\chi}} \frac{f_{v_{\odot}}(u_{\chi})}{u_{\chi}} \omega(r) \int_0^{v_e(r)} dv R_i^-(\omega \rightarrow v) |F_i(q)|^2, \quad (7.5)$$

1731 where the summation extends over all possible nuclear targets. In this expression, R_{\odot}
1732 is the radius of the Sun, ρ_{χ} is the local DM density, m_{χ} the mass of the DM particle,
1733 $f_{v_{\odot}}(u_{\chi})$ the DM velocity distribution seen from the Sun's reference frame, $R_i^-(\omega \rightarrow v)$
1734 is the differential rate at which a DM particle with velocity v scatters a solar target of
1735 mass m_i to end up with a velocity ω and $|F_i(q)|$ is the nuclear form factor of target i .

1736 The differential scattering rate takes a rather simple form when considering velocity-

Chapter 7. DM searches with neutrinos from the Sun

¹⁷³⁷ independent and isotropic cross sections. In that case, this quantity is given by [4, 71]:

$$R_i^-(\omega \rightarrow v) = \frac{2}{\sqrt{\pi}} \frac{\mu_{i,+}^2}{\mu_i} \frac{v}{\omega} n_i(r) \sigma_i \left[\chi(-\alpha_-, \alpha_+) + \chi(-\beta_-, \beta_+) e^{\mu_i(\omega^2 - v^2)/u_i^2(r)} \right], \quad (7.6)$$

¹⁷³⁸ where μ_i is the ratio between the DM mass and the mass of target i , $\mu_{i,\pm}$ is defined as:

$$\mu_{i,\pm} \equiv \frac{\mu_i \pm 1}{2}, \quad (7.7)$$

¹⁷³⁹ $n_i(r)$ is the density profile of target i in the solar medium, $u_i(r)$ is the most probable

¹⁷⁴⁰ velocity of target i given by:

$$u_i(r) = \sqrt{\frac{2T_\odot(r)}{m_i}}, \quad (7.8)$$

¹⁷⁴¹ where $T_\odot(r)$ is the temperature of the Sun, the quantities α_\pm and β_\pm are defined as:

$$\alpha_\pm \equiv \frac{\mu_{i,+}v \pm \mu_{i,-}\omega}{u_i(r)}, \quad (7.9)$$

$$\beta_\pm \equiv \frac{\mu_{i,-}v \pm \mu_{i,+}\omega}{u_i(r)}, \quad (7.10)$$

¹⁷⁴² and the function $\chi(a, b)$ is a Gaussian integral of the form:

$$\chi(a, b) \equiv \int_a^b dx e^{-x^2}. \quad (7.11)$$

¹⁷⁴³ Finally, if one assumes the DM halo velocity distribution in the galactic rest frame

¹⁷⁴⁴ to be a Maxwell-Boltzmann distribution, one can write the halo velocity distribution for

¹⁷⁴⁵ an observer moving at the speed of the Sun with respect to the DM rest frame as:

$$f_{v_\odot}(u_\chi) = \sqrt{\frac{3}{2\pi}} \frac{u_\chi}{v_\odot v_d} \left(e^{-\frac{3(u_\chi - v_\odot)^2}{2v_d^2}} - e^{-\frac{3(u_\chi + v_\odot)^2}{2v_d^2}} \right), \quad (7.12)$$

¹⁷⁴⁶ where:

$$\omega^2 = u_\chi + v_e(r)^2, \quad (7.13)$$

¹⁷⁴⁷ is the DM velocity squared, v_\odot the relative velocity of the Sun from the DM rest frame

7.2. Gravitational capture of DM by the Sun




Figure 7.1: Input solar parameters used in our capture rate computation as functions of the Sun's radius, from left to right: temperature (with respect to the temperature at the core), mass (in solar masses) and electron number density (with respect to the electron density at the core). All quantities shown correspond to the standard solar model BS2005-OP [3].

1748 and $v_d \simeq \sqrt{3/2}v_\odot$ the velocity dispersion.

1749 For the case of strong scattering cross section, Eq. (7.5) ceases to be valid, as it
 1750 escalates indefinitely with the cross section. In that limit, the capture rate saturates to
 1751 the case where the probability of interaction is equal to one, which can be written as:

$$C_\odot^{\text{geom}} = \pi R_\odot^2 \left(\frac{\rho_\chi}{m_\chi} \right) \langle v \rangle \left(1 + \frac{3}{2} \frac{v_e^2(R_\odot)}{v_d^2} \right) \xi(v_\odot, v_d), \quad (7.14)$$

1752 where $v_d = \sqrt{8/3\pi}v_d$ is the mean velocity in the DM rest frame and the factor $\xi(v_\odot, v_d)$
 1753 accounts for the suppression due to the motion of the Sun:

$$\xi(v_\odot, v_d) = \frac{v_d^2 e^{-\frac{3v_\odot^2}{2v_d^2}} + \sqrt{\frac{\pi}{6}} \frac{v_d}{v_\odot} (v_d^2 + 3v_e^2(R_\odot) + 3v_\odot^2) \text{Erf} \left(\sqrt{\frac{3}{2}} \frac{v_\odot}{v_d} \right)}{2v_d^2 + 3v_e^2(R_\odot)}. \quad (7.15)$$

1754 Having these into account, one can write the total capture rate as a combination of
 1755 both contributions, allowing a smooth transition between the two, as:

$$C_\odot = C_\odot^{\text{weak}} \left(1 - e^{C_\odot^{\text{geom}}/C_\odot^{\text{weak}}} \right). \quad (7.16)$$

1756 I computed the capture rate from Eq. (7.16) in the case of interactions with
 1757 electrons. To do so, I used the standard solar model BS2005-OP [3]. Fig. 7.1 shows the

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.2: Capture rates as a function of the DM mass for the DM-electron interactions (red lines), SD DM-nucleons interactions (green lines) and SI DM-nucleons interactions (blue lines). Solid lines represent the values computed in this work while the dashed lines are the one given in Ref. [4]. All the rates are shown for a choice of scattering cross section of $\sigma_i = 10^{-40} \text{ cm}^2$.

1758 three parameters from the solar model that are needed for the computation, the solar
 1759 temperature (left panel), mass (central panel) and electron density (right panel) profiles.

1760 For the case of the interactions off nuclei, the computations are more convoluted
 1761 as one needs to add up the contributions of the different most abundant nuclei in
 1762 the Sun. Also, in contrast to the electron scenario where the form factor is trivially
 1763 $|F_e(q)|^2 = 1$, for any nucleus i one would need to consider some appropriate nuclear
 1764 density distribution (either a Gaussian approximation, a Woods-Saxon distribution, etc)
 1765 which would complicate the calculations even further.

1766 That is the reason why, at this stage of our study, I decided to take an alternative
 1767 approach to the computation of the DM-nucleus capture rates. I used the **DarkSUSY**
 1768 software, that allows us to compute these quantities performing a full numerical
 1769 integration over the momentum transfer of the form factors. The default standard
 1770 solar model used by **DarkSUSY** is BP2000¹ [72].

¹This is what they say in their manual, but I fear it is somewhat outdated. It appears to me this

7.2. Gravitational capture of DM by the Sun

1771 In Fig. 7.2 I show the results I obtained for the capture rates, for the case of
 1772 interactions off electrons (red solid line), SD (green solid line) and SI (blue solid line)
 1773 interactions of nucleons. In all cases I used a value of the scattering cross sections of
 1774 $\sigma_i = 10^{-40} \text{ cm}^2$. Note here one of the limitations of the **DarkSUSY** approach, one can
 1775 not extend the computation below $m_{\text{DM}} = 1 \text{ GeV}$. Nevertheless, this is not something
 1776 to worry about in this case, as I will discuss next. As a comparison, I added also the
 1777 values computed in Ref. [4] (same color scheme, dashed lines). One can see there is good
 1778 agreement between these and the **DarkSUSY** computation of the SD and SI interactions
 1779 for $m_{\text{DM}} \geq 1 \text{ GeV}$. In this regime their computations also matches quite well our
 1780 result for the electron capture rate. However, these start to differ significantly below
 1781 $m_{\text{DM}} = 1 \text{ GeV}$, being their estimate up to a factor of 5 bigger than ours for low masses.
 1782

1782 Let us comment briefly about the assumption I made before about not including
 1783 an evaporation term in the Boltzmann equation. If I include this term in the equation
 1784 (which will be proportional to the number of DM particles) the equilibrium solution
 1785 takes the form:

$$N_{\text{DM}}^{eq} = \sqrt{\frac{C_\odot}{A_\odot}} \frac{1}{\kappa + \frac{1}{2} E_\odot \tau_{eq}}, \quad (7.17)$$

1786 where E_\odot is the total evaporation rate, τ_{eq} is the equilibrium time in the absence of
 1787 evaporation:

$$\tau_{eq} = \frac{1}{\sqrt{C_\odot A_\odot}}, \quad (7.18)$$

1788 and κ is defined as:

$$\kappa \equiv \sqrt{1 + \left(\frac{E_\odot \tau_{eq}}{2} \right)^2}. \quad (7.19)$$

1789 Now, it is easy to proof that in case evaporation dominates $\kappa \gg 1$ and therefore:

$$N_{\text{DM}}^{eq} \simeq \frac{C_\odot}{E_\odot}. \quad (7.20)$$

1790 In contrast, if evaporation is irrelevant $\kappa \simeq 1$ and one recovers Eq. (7.2).

model is relatively old and do not see why they are not using others like [3]. Maybe one can double-check in the code to make sure.

Chapter 7. DM searches with neutrinos from the Sun

1791 In this way, one can define the evaporation mass as the mass for which the number
1792 of DM particles in equilibrium approaches Eq. (7.20) at 10% level:

$$\left| N_{DM}^{eq}(m_{\text{evap}}) - \frac{C_{\odot}(m_{\text{evap}})}{E_{\odot}(m_{\text{evap}})} \right| = 0.1 N_{DM}^{eq}(m_{\text{evap}}). \quad (7.21)$$

1793 This can be regarded as the minimum testable mass one can reach using the annihilation
1794 products of the DM in the Sun.

1795 It was reported in Ref. [4] that, in the case of both SD and SI DM interactions
1796 off nuclei, this value ranges from 2 to 4 GeV depending on the specific scattering
1797 cross section value, compatible with the usual assumptions in the literature. What is
1798 interesting is the case of the electron capture. It was found that, when one applies a
1799 cutoff in the velocity distribution of the DM trapped in the Sun slightly below the escape
1800 velocity, the evaporation mass for the DM-electron interaction decreases remarkably. For
1801 a moderate choice of $v_c(r) = 0.9v_e(r)$ one gets an evaporation mass of around 200 to
1802 600 MeV. This possibility opens a region of the parameter space that could be tested
1803 with neutrino detectors.

1804 7.3 Neutrino flux from DM annihilations

1805 When WIMPs annihilate inside the Sun a flux of high-energy neutrinos is expected from
1806 heavy quarks, gauge bosons and $\tau^+\tau^-$ final states, which decay before losing energy
1807 in the dense solar medium, as they will produce a continuum spectra up to $E_{\nu} \sim m_{\chi}$
1808 (in the case of direct annihilation to neutrinos one would have a line at $E_{\nu} = m_{\chi}$) [66].
1809 This kind of signal has been extensively studied in the literature, allowing to put strong
1810 limits on the SD WIMP-proton cross section for large m_{χ} . However, the number of
1811 high-energy neutrinos per WIMP annihilation is small and the spectrum depends on the
1812 unknown final state. Moreover, background rejection is easier for large m_{χ} but neutrinos
1813 with $E_{\nu} \gtrsim 100$ GeV are significantly attenuated by interactions in the Sun.

1814 Nevertheless, most WIMP annihilation final states eventually produce a low-energy
1815 neutrino spectrum. In this case one does not just consider the more massive final

7.4. Computing limits from solar neutrino fluxes

states but also annihilations into e^+e^- , $\mu^+\mu^-$ and light quarks [65]. In particular, light mesons would be produced and stopped in the dense medium, thus decaying at rest and producing a monoenergetic neutrino signal. The decay-at-rest of kaons will produce a $E_\nu = 236$ MeV ν_μ while in the case of pions one would have a $E_\nu = 29.8$ MeV ν_μ . In practice only K^+ and π^+ contribute to these signals, as K^- and π^- are usually Coulomb-captured in an atomic orbit and get absorbed by the nucleus. There is also a low-energy neutrino signal coming from muon decays, which are produced in kaon or pion decays, leptonic decays of other hadrons and heavy leptons or even directly from WIMP annihilations, which can decay at rest and contribute to the previous low-energy neutrino flux with a well known spectrum below 52.8 MeV.

These monoenergetic MeV neutrinos were previously considered undetectable but, due to the large yield, the known spectra and the modern advances in the detector technology, these low-energy neutrino flux can be a good probe of the SD WIMP-proton cross-section in standard solar WIMP capture scenario, as it is sensitive to low WIMP masses and insensitive to the particular final state. A good place to look for these signals are next-generation neutrino experiments such as DUNE.

7.4 Computing limits from solar neutrino fluxes

In order to use the neutrino fluxes from DM annihilations in the Sun, the first thing I need to do is to determine the expected number of atmospheric background events, for a given exposure, after directionality selection has been applied. I can write this number as:

$$N_B = \eta_B \int d\Omega \int_{E_{min}}^{E_{max}} dE_\nu \frac{d^2\Phi_{atm}^\mu}{dE_\nu d\Omega} \times \left(A_{eff}^{(\mu)}(E_\nu) T \right), \quad (7.22)$$

where η_B is the background efficiency, E_{min} and E_{max} the minimum and maximum energies to integrate over, $d^2\Phi_{atm}^\mu/dE_\nu d\Omega$ the differential flux of atmospheric muon neutrinos, $A_{eff}^{(\mu)}$ is the effective area of DUNE to muon neutrinos and T is the exposure time. The effective area can be expressed as the product of the neutrino-nucleus scattering cross section and the number of nuclei in the fiducial volume of the detector. This way

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.3: *NuWro* computed $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section as a function of the neutrino energy E_μ . The black line shows to the total cross section, whereas the others correspond to the different contributions (in red quasi-elastic scattering, in green resonant pion exchange, in blue deep inelastic scattering and in purple meson exchange current).

1842 for DUNE I can write:

$$A_{eff}^{(\mu)}(E_\nu) = (6.0 \times 10^{-10} \text{ m}^2) \left(\frac{\sigma_{\nu - \text{Ar}}^{(\mu)}(E_\nu)}{10^{-38} \text{ cm}^2} \right) \left(\frac{M_{target}}{40 \text{ kT}} \right), \quad (7.23)$$

1843 where $\sigma_{\nu - \text{Ar}}^{(\mu)}$ is the $\nu_\mu - {}^{40}\text{Ar}$ charged-current scattering cross section. In Fig. 7.3 I
 1844 show the computed value of this cross section as a function of the neutrino energy E_ν ,
 1845 in the range of interest both for the atmospheric background and signal events. It was
 1846 computed using the NuWro Monte Carlo neutrino event generator [73], including the
 1847 charged-current contributions of the quasi-elastic scattering (red line), resonant pion
 1848 exchange (green line), deep inelastic scattering (blue line) and meson exchange current
 1849 (purple line).

1850 The background rejection will depend on the resolution of the detector and the
 1851 selection one applies on the events. A geometry argument can be used to estimate
 1852 the maximum background rejection one can achieve in this case, considering one can

7.4. Computing limits from solar neutrino fluxes

1853 efficiently discriminate all events coming from a direction different from that of the
 1854 Sun. In that case, the optimal background efficiency will simply be the relative angular
 1855 coverage of the Sun. Taking the angular diameter of the Sun as seen from the Earth to
 1856 be 0.5° , I have:

$$\eta_B^{(opt)} \approx \frac{\pi \left(\frac{0.5}{2}\right)^2}{360 \times 180} \simeq 3.03 \times 10^{-6}. \quad (7.24)$$

1857 This value will give a very optimistic estimate of the number of background events.
 1858 However, it can be regarded as an lower limit, as it represents the best case scenario.

1859 In Fig. 7.4 I show the fluxes of atmospheric neutrinos at the Homestake mine during
 1860 solar minimum, taken from Ref. [5]. The values are averaged over the two angular
 1861 directions. In blue I have the flux of muon neutrinos while in red I indicate the flux
 1862 of electron neutrinos. Additionally, the dashed lines correspond to both antineutrino
 1863 species.

1864 Using these values for the muon neutrino and the corresponding total CC cross
 1865 section, one can compute the number of expected background events by integrating over
 1866 the given energy range (as in this case the angular integral is trivial). As for the energy
 1867 range to integrate over, I choose the range for DUNE specified in [42], $E_{min} = 10^{-1}$ GeV
 1868 and $E_{max} = 10$ GeV. Taking all these into account, I found the number of background
 1869 events to be:

$$N_B \simeq \eta_B \times (3.827 \times 10^4) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right). \quad (7.25)$$

1870 In order to estimate the sensitivity of DUNE to this kind signal, one can consider a
 1871 hypothetical data set where the number of observed neutrinos is taken to be the expected
 1872 number of background events rounded to the nearest integer, $N_{obs} = \text{round}(N_B)$ [74].
 1873 Now, if I assume that the number of signal and background events seen by DUNE are
 1874 given by Poisson distributions with means equal to the expected number of signal and
 1875 background events, N_S and N_B , one can denote by N_S^{90} to the number of expected
 1876 signal events such that the probability of having an experimental run with a number of
 1877 events greater than N_{obs} is 90%. This number can be obtained as the numerical solution

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.4: Expected atmospheric neutrino flux as a function of the neutrino energy E_ν at Homestake at solar minimum, taken from Ref. [5]. The blue solid (dashed) line correspond to muon neutrinos (antineutrinos) and the red solid (dashed) line correspond to electron neutrinos (antineutrinos).

1878 to the equation:

$$1 - \frac{\Gamma(N_{obs} + 1, N_S^{90} + N_B)}{N_{obs}!} = 0.9, \quad (7.26)$$

1879 where $\Gamma(x, y)$ is the upper incomplete gamma function.

1880 The number of signal events is related to the neutrino flux from DM annihilations in
 1881 a similar way as the background events to the atmospheric neutrino flux. In this case I
 1882 have:

$$N_S = \eta_S \Gamma_A^{eq} \int_{z_{min}}^{z_{max}} dz \frac{dN_\nu}{dAdN_Adz} \times (A_{eff}^\mu(z)T), \quad (7.27)$$

1883 where η_S is the signal efficiency, Γ_A^{eq} is the total annihilation rate of DM particles at
 1884 equilibrium, $\Gamma_A^{eq} = A_\odot (N_{DM}^{eq})^2$, z_{min} and z_{max} the minimum and maximum relative
 1885 energies to integrate over (in such a way that $z_{min,max} \leq E_{min,max}/m_{DM}$ for each m_{DM})
 1886 and $dN_\nu/dAdN_Adz$ the muon neutrino flux per DM annihilation in the Sun.

1887 Knowing N_S^{90} one can use the relation in Eq. (7.27) to obtain $\Gamma_A^{eq,90}$ for different
 1888 values of the DM mass. From there I can directly translate those values into the

7.5. Example: Kaluza-Klein Dark Matter

upper limits for DUNE on the DM scattering cross sections, for a given exposure. The relation between the annihilation rate and the DM-nucleon cross section comes from the equilibrium condition through the solar DM capture rate. The details of the evolution of the number of DM particles inside the Sun and the computation of the capture rates are discussed in App. 7.2.

7.5 Example: Kaluza-Klein Dark Matter

Even though there are plenty of BSM theories which provide viable dark matter candidates, Kaluza-Klein type of models [75, 76] within the universal extra dimensions (UED) paradigm naturally predict the existence of a massive, stable particle that can play the role of the dark matter. In the UED scenario all the SM fields can propagate in one or more compact extra dimensions [77], as opposed to the idea of brane worlds [78, 79], where just gravity can propagate in the bulk while SM particles live at fixed points.

Furthermore, in UED there is no violation of the translational invariance along the extra dimensions, thus leading to degenerate KK modes masses and also the conservation of the KK number in the effective four dimensional theory. At loop level, radiative corrections and boundary terms shift the masses of the KK modes and break KK number conservation into a KK parity. As a result, this theory only contains interactions between an even number of odd KK modes and therefore the lightest among the first KK excitations will be stable. This particle is usually denoted as the lightest Kaluza-Klein particle (LKP) and its mass is proportional to $1/R$, being R the size of the extra dimension.

A viable DM candidate needs to be electrically neutral and non-baryonic, therefore good candidates among the first Kaluza-Klein excitations would be the KK neutral gauge bosons and the KK neutrinos [80]. Another possible candidate is the first KK excitation of the graviton, which receives negligible radiate contributions and therefore has a mass almost equal to $1/R$, but it has been shown that the lightest eigenstate from

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.5: Feynman diagrams for $B^1 B^1$ annihilation into SM fermions.




Figure 7.6: Feynman diagrams for $B^1 B^1$ annihilation into a Higgs boson pair.

7.5. Example: Kaluza-Klein Dark Matter




Figure 7.7: Computed spectra of muon neutrinos at the DUNE FD site from B^1 annihilations in the Sun for three different values of M_{LKP} , plotted in relative energy units for legibility.

the mixing of the gauge mass states (B^1, W_3^1) would be lighter, as B^1 and W_3^1 receive negative radiative corrections [81]. It is also understood that, when these corrections become sizeable, the eigenstates become approximately pure B^1 and W_3^1 states as the Weinberg mixing angle grows small with the KK number [81]. In that case, the LKP can be well-approximated as being entirely B^1 .

I need to compute the neutrino flux produced by the annihilations of the LKP in the core of the Sun, taking into account their propagation in the solar medium, as well as neutrino oscillations. To this end I used `WimpSim` [82, 83] to generate one million annihilation events in the Sun over a time span of four years and propagate them to the DUNE FD location ($44^\circ 20' \text{ N}, 103^\circ 45' \text{ W}$), for different values of M_{LKP} . In Fig. 7.7 I show the obtained muon neutrino spectra arriving to the detector from LKP annihilations in the Sun, per unit area and per annihilation, plotted in relative energy units for different values of the mass. As one could expect the spectra get steeper the higher is the mass, due to the absorption of high-energy neutrinos in the solar medium. Also, one can see the peak at $z = 1$ due to the direct annihilation into neutrinos $\chi\chi \rightarrow \nu\bar{\nu}$.

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.8: Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin-dependent B^1 -proton scattering cross section as a function of M_{LKP} (green dots). I also show the previous limits from IceCube [6] (blue line) and Antares [7] (red line) on the LKP cross section. The shaded area represents the disfavoured region (at 95% confidence level) on the mass of the LKP from LHC data [8].

Now, one can estimate the sensitivity of DUNE to this particular model by using the methods I previously discussed. To begin with, I will use the optimistic estimation of the background efficiency in Eq. (7.24) to get our upper bound. Using it, one can directly compute the number of expected background events to be $N_B = 0.1101$ for an exposure of 400 kT yr. Then, Eq. (7.26) give us a value of $N_S^{90} = 2.20$ for the 90% exclusion number of expected signal events. By using the NuWro generated cross sections and the computed neutrino fluxes from B^1 annihilations in the Sun I can estimate the limits on the SD and SI DM-nucleus cross section using the relation in Eq. (7.2) and the capture rates I computed with DarkSUSY.

In Fig. 7.8 I show the projected sensitive for DUNE on the spin-dependent B^1 -proton scattering cross section versus the mass of the DM particle, for a exposure of 400 kT yr (green dots). I also include the previous results from IceCube [6] (blue line) and Antares [7] (red line). The shaded area represents the disfavoured region from combined searches for UED by ATLAS and CMS [8].

7.6. High energy DM neutrino signals




Figure 7.9: Computed spectra of muon neutrinos at the DUNE FD site from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilations in the Sun for the DM masses $m_{\text{DM}} = 10 \text{ GeV}$ (red line), 50 GeV (green line) and 100 GeV (blue line), plotted in relative energy units.

From the experimental point of view, this estimation lacked a detailed simulation of the detector response and thus this must be consider as a mere optimistic sensitivity computation. However, it shows the potential of DUNE to constrain this kind of exotic scenarios, showing the region where it will be in a position to compete with other neutrino telescopes. A more detailed analysis is needed if I am to make a realistic estimation. Even though the region of the parameter space where DUNE would be sensitive to this particular model is quite constrained by collider searches [8] and other rare decay measurements [84, 85], it still constitutes an alternative indirect probe.

7.6 High energy DM neutrino signals

To have better estimates on the capability of the DUNE FD to constrain the parameter space of DM using solar neutrino fluxes, I need to start accounting for the detector resolution effects and the topologies of the different signatures. As a starting point, I will focus on specific annihilation channels. For the case of DUNE, the relevant ones are mainly the hard channels $\tau^+\tau^-$ and $\nu\bar{\nu}$ and the soft channel $b\bar{b}$. These are the open annihilation channels for relatively low mass WIMPs that will actually give neutrino fluxes. Other channels, like W^+W^- and ZZ , are open for more massive WIMPs, but

Chapter 7. DM searches with neutrinos from the Sun

1961 those will produce usually a higher energy neutrino flux that will be out of reach for
1962 DUNE (usually the maximum neutrino energy is taken to be $E_{max} = 10$ GeV).

1963 In Fig. 7.9 I show the `WimpSim` [82, 83] generated muon neutrino spectra at the
1964 DUNE FD location ($44^\circ 20' N, 103^\circ 45' W$) from $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel)
1965 annihilations in the core of the Sun, for different DM masses. Here, one can clearly see
1966 the meaning of the previous distinction between hard and soft channels. For the same
1967 DM mass value, the muon neutrino spectrum from the $\tau^+\tau^-$ channel is more flat and
1968 reaches higher energies than the one from the $b\bar{b}$ channel, which drops faster.

1969 In this case, I prepared two sets of files, one for $\tau^+\tau^-$ and the other for $b\bar{b}$, for DM
1970 masses in the range from 5 to 100 GeV (actually for $b\bar{b}$ the first mass point I took is
1971 7.5 GeV, as a WIMP with $m_{DM} = 5$ GeV can not kinematically self annihilate into $b\bar{b}$).
1972 Then, I prepared the `WimpSim` output fluxes in a specific way to use them as inputs to
1973 `NuWro`, which simulates the neutrino interaction with the argon.

1974 Because `WimpSim` outputs an event list together with the fluxes, I can use the former
1975 to generate the events. The direction of these is given in terms of the azimuth and
1976 altitude angles viewed from the specified location, so first I need to convert these into the
1977 DUNE FD coordinates. Once I have done it, each event can be processed with `NuWro`.
1978 To increase the number of samples and optimise the computation time, I generate 100
1979 interactions (i.e. `NuWro` events) for each `WimpSim` event². I restrict the event generation
1980 to charged current interactions, but I allow all the different contributions to the CC
1981 cross section, i.e. quasielastic scattering (QEL), meson exchange current process (MEC),
1982 resonant pion production (RES) and deep inelastic scattering (DIS). I just take into
1983 account the CC contribution because I am only interested in final states with charged
1984 leptons, as we have better chances of reconstructing the kinematics of CC events.

1985 For the atmospheric fluxes I follow a similar procedure, only that this time I do not
1986 have a set of events but the fluxes binned in azimuth and altitude angles. This way, I
1987 transform these to DUNE coordinates and process the fluxes for each bin separated with

²This also solves a problem related with the generation of the neutrino interactions in `NuWro`, as if you only produce one event each time you launch `NuWro` it will always produce an interaction of the dominant interaction type for that particular energy.

7.6. High energy DM neutrino signals




Figure 7.10: Distribution of the muon neutrino energies from the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) annihilation channels, for $m_{\text{DM}} = 10 \text{ GeV}$, separated by CC interaction type: QEL (blue), MEC (orange), RES (green) and DIS (red).

1988 NuWro.

1989 At this point, I have two sets of events with different energies and final states.
 1990 In Fig. 7.10 one can see the distribution of the muon neutrino energies for the case
 1991 $m_{\text{DM}} = 10 \text{ GeV}$, both for the $\tau^+\tau^-$ (left panel) and $b\bar{b}$ (right panel) channels, separated
 1992 by interaction. One can clearly see that there are different energy regimes where the
 1993 primary interaction type is different. This leads to a plurality of event topologies,
 1994 therefore making it difficult to implement a general approach to the selection of events
 1995 in detriment of the background. As a way to proceed, I decided to split our samples,
 1996 based on the different interaction modes and contents of the final state, into a CC DIS
 1997 sample and a single proton CC QEL sample.

1998 **7.6.1 DIS events**

1999 To begin with, I consider the high energy part of the spectrum. In this region DIS events
 2000 dominate, i.e. interactions of the form $\nu_\mu + q_d(\bar{q}_u) \rightarrow \mu^- + q_u(\bar{q}_d)$. Therefore, our final
 2001 estates will contain a muon and a hadronic jet from the fragmentation of the outgoing
 2002 quark. As all these events have $E_\nu \gtrsim 1 \text{ GeV}$ the momentum transfer to the remnant
 2003 nucleus is negligible, for this reason the neutrino energy can be effectively reconstructed

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.11: Distributions of θ_μ (left panel), θ_j (central panel) and θ_{plane} (right panel) for the $b\bar{b}$ sample with $m_{\text{DM}} = 10$ GeV (blue) and the atmospheric background (red).

just taking into account the momenta of the muon and the jet. This technique was successfully used in Ref. [86] to select monoenergetic DM solar neutrino events from $\nu\bar{\nu}$ annihilation channels.

Using momentum conservation one sees that the plane generated by the momenta of the muon and the jet needs to also contain the momentum of the neutrino. As we are interested in neutrinos coming from the Sun, the momentum of the neutrino can be regarded as known beforehand. This will allow us to define the angle of the outgoing muon and jet with respect to the incoming neutrino. Moreover, one can also use that information to reject poorly reconstructed jets, checking for deviations of these from the momentum conservation plane.

To account for the limited angular resolution of the detector, I smeared the momenta of the muons and hadrons. In a liquid argon TPC muons are expected to be tracked with high precision, therefore I take the associated angular resolution to be 1° . In the case of jets, it is expected that for the hadrons dominating the cascade a detector like DUNE has an angular resolution between 1° to 5° [42], so I take the latter, more conservative, estimate.

As a first selection step, I will just take into account particles with kinetic energies above the detection threshold of DUNE. For muons and photons the specified threshold energy is 30 MeV, for charged pions 100 MeV and for other hadrons 50 MeV [42]. This way, if the outgoing muon in a certain event has an energy lower than the required

7.6. High energy DM neutrino signals




Figure 7.12: Left panel: signal efficiencies (blue lines) and background rejections (red lines) for events passing the cuts $\theta < \theta_{\text{cut}}$ for the jet (solid lines) and muon (dashed lines) angles. Right panel: signal efficiency (blue line) and background rejection (red line) for events passing the cut $\theta_{\text{plane}} < \theta_{\text{cut}}$ for the momentum conservation plane deviation.

threshold I will drop such event. For the case of hadrons and photons, I will only require to have at least one particle above the energy threshold, so then one can compute the jet momentum using the (smeared) momenta of the N particles above threshold as:

$$\vec{p}_j = \sum_{i=1}^N \vec{p}_i. \quad (7.28)$$

Additionally, I will also define an estimation of the deposited hadronic energy as:

$$E_j^{\text{dep}} = m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}} + \sum_{i=1}^N \sqrt{|\vec{p}_i|^2 + m_i^2}. \quad (7.29)$$

This quantity is useful to select events with enough hadronic visible energy in the detector. For events where most of the hadronic energy is scattered across plenty of hadrons with individual energies below the detection threshold, this estimation will give $E_j^{\text{dep}} \leq 0$. In these cases it could be expected that the jet momentum is poorly reconstructed, and therefore I require events to pass the cut $E_j^{\text{dep}} > 0$.

For the events I can compute the angles for the muon and jet with respect to the

Chapter 7. DM searches with neutrinos from the Sun

2034 incoming neutrino as:

$$\cos \theta_\mu = \hat{p}_\nu \cdot \hat{p}_\mu, \quad (7.30)$$

$$\cos \theta_j = \hat{p}_\nu \cdot \hat{p}_j, \quad (7.31)$$

2035 and the deviation from the momentum conservation plane as:

$$\sin \theta_{plane} = \left| \frac{\hat{p}_\mu \times \hat{p}_\nu}{|\hat{p}_\mu \times \hat{p}_\nu|} \cdot \hat{p}_j \right|. \quad (7.32)$$

2036 In Fig. 7.11 I show some distributions of these quantities for the case of the $b\bar{b}$ sample
 2037 with $m_{DM} = 10$ GeV (blue histograms) and for the atmospheric backgrounds (red).
 2038 In order to select the atmospheric events I followed the same criteria as for the signal
 2039 events. However, because in the signal case I used the true direction of the neutrino
 2040 as input, as it should be that of the Sun at that time and therefore known, in the
 2041 atmospheric case I used a set of solar positions as our ansatz for the neutrino direction.
 2042 From the distributions, one can see that the muon and the jet for the signal events are
 2043 predominantly forward and also that the deviations from the momentum conservation
 2044 plane are peaked at zero, as one should expect.

2045 Now, I can start applying cuts to maximise our signal selection efficiency while at
 2046 the same time I try to minimise the amount of atmospheric background events passing
 2047 the selection. To this end, I will need to find some lower and upper cuts for θ_j and
 2048 θ_μ and an upper bound for θ_{plane} . In Fig. 7.12 I show how upper bound cuts in the
 2049 different angular variables affect the signal efficiency (blue lines) and the background
 2050 rejection (red lines). Notice that the signal efficiency behaves in a quite different way
 2051 when I apply cuts in the jet and the muon angles. On the contrary, the cuts on both
 2052 variables have a similar effect on the background rejection.

2053 In order to obtain the optimal set of cuts, I perform a multidimensional scan. I
 2054 do this separately for the $\tau^+\tau^-$ and the $b\bar{b}$ samples. For each case, I scan the possible
 2055 cuts for each mass point and then I take the mean value of the signal efficiency for

7.6. High energy DM neutrino signals




Figure 7.13: Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) DIS samples as functions of the DM mass, m_{DM} , obtained by applying the optimal angular cuts $\theta_\mu < 27^\circ$, $4^\circ < \theta_j < 26^\circ$ and $\theta_{\text{plane}} < 3.5^\circ$.

2056 each configuration, to get the mean efficiency for each set of cuts. I do a similar scan
 2057 for the atmospheric sample independently. Then, I take the sets of cuts such that
 2058 the background rejection achieved is greater than 99.8% and search for the one which
 2059 maximises the $\tau^+\tau^-$ and $b\bar{b}$ sample mean efficiencies. I found that with the cuts $\theta_\mu < 27^\circ$,
 2060 $4^\circ < \theta_j < 26^\circ$ and $\theta_{\text{plane}} < 3.5^\circ$ I get a background rejection of 99.80% while achieving
 2061 a 49.40% and 44.92% mean signal efficiencies for the $\tau^+\tau^-$ and $b\bar{b}$ signals respectively.

2062 In Fig. 7.13 I show the signal efficiencies as a function of the DM mass for the $\tau^+\tau^-$
 2063 (blue line) and the $b\bar{b}$ (red line) DIS events, after applying the cuts discussed above, as
 2064 well as the energy threshold and hadronic visible energy selections. One can see that
 2065 the efficiency grows with the mass, as annihilations of more massive DM particles will
 2066 produce a neutrino spectrum centered at higher energies, where DIS events dominate.
 2067 Notice also that the efficiency is higher for the $\tau^+\tau^-$ case at every mass point, as in
 2068 general this channel produces neutrinos at higher energies than the corresponding $b\bar{b}$
 2069 channel.

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.14: Distributions of $\cos \theta_\mu$ (left panel), $\cos \theta_p$ (central panel) and $\cos \theta_N$ (right panel) for the $\tau^+\tau^-$ QEL sample with $m_{\text{DM}} = 5 \text{ GeV}$ (blue) and the atmospheric background (red).

2070 7.6.2 Single proton QEL events

2071 Now, one can try to explore the low energy tail of the neutrino energy distributions. This
 2072 regime is dominated by the QEL interactions, i.e. events of the type $\nu_\mu + n \rightarrow \mu^- + p$.
 2073 In this case, as the typical energies are $E_\nu \lesssim 1 \text{ GeV}$, the momentum transfer to the
 2074 remnant nucleus is sizeable. Therefore, I can not make the approximation I did before
 2075 and assume that the momentum of the muon and the proton will give an adequate
 2076 estimation of the reconstructed neutrino energy.

2077 In any case, as before, I can take the direction of the incoming neutrino as known.
 2078 That way, one can estimate the energy of the neutrino as:

$$E_\nu^{reco} = E_\mu + E_p + m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}}, \quad (7.33)$$

2079 and using momentum conservation I can write the momentum of the remnant nucleus
 2080 as:

$$\vec{p}_N = \hat{p}_\nu (E_\mu + E_p + m_{^{39}\text{Ar}} - m_{^{40}\text{Ar}}) - \vec{p}_\mu - \vec{p}_p. \quad (7.34)$$

2081 As in the previous case, I need to drop the events where the muon or the proton fall
 2082 below the kinetic energy detection threshold [42]. Also, I again apply a smearing to the
 2083 momenta of the particles, a 1% for muons and 5% for protons.

2084 Having done that, one can compute the following angular variables for our selected

7.6. High energy DM neutrino signals




Figure 7.15: Left panel: value of the loss function for the training sample (blue line) and accuracy for the validation sample (red line) versus the number of iterations for the MLP classifier training. Right panel: distributions of the predicted probabilities assigned by the MLP classifier to the test sample for the $\tau^+\tau^-$ QEL signal with $m_{\text{DM}} = 5 \text{ GeV}$ (blue) and the atmospheric background (red).

2085 events:

$$\cos \theta_\mu = \hat{p}_\nu \cdot \hat{p}_\mu, \quad (7.35)$$

$$\cos \theta_p = \hat{p}_\nu \cdot \hat{p}_p, \quad (7.36)$$

$$\cos \theta_N = \hat{p}_\nu \cdot \hat{p}_N. \quad (7.37)$$

2086 Fig. 7.14 shows the distributions of these angular variables for the $\tau^+\tau^-$ QEL
 2087 sample with $m_{\text{DM}} = 5 \text{ GeV}$ (blue) and the atmospheric background (red). Again, for
 2088 the atmospheric events I used a random solar position as the ansatz for the incoming
 2089 neutrino direction. Notice that now, opposed to the DIS case where the signal had very
 2090 sharp distributions for the variables considered, the shapes of the angular distributions
 2091 for signal and background are not that much different.

2092 This effectively means that the usual approach of applying simple angular cuts would
 2093 not work as well as in the previous situation. Therefore, as a possible solution, I tried to
 2094 use a multilayer perceptron (MLP) classifier to separate between signal and background
 2095 events. Thus, the power of the hypothesis test will serve as an estimate of the signal

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.16: Signal efficiencies for the $\tau^+\tau^-$ (blue line) and $b\bar{b}$ (red line) single proton QEL samples as functions of the DM mass, m_{DM} , obtained by requiring a minimum predicted probability from the MLP classifier of 0.97 in order to achieve a background rejection greater than 99.8%.

efficiency, and in the same way one can take the size of the test to be our background rejection.

For each DM mass value and channel, as well as for the background sample, I divide our events into training, validation and test samples. The input variables for the classifier were the reconstructed neutrino energy from Eq. (7.33) and the angular variables defined in Eqs. (7.35 - 7.37). I used the MLP classifier implemented in `scikit-learn` [87], with a total of five hidden layers, the rectified linear unit activation function and adaptive learning rate. In order to account for fluctuations due to artifacts in the training process I repeated the training a thousand times for each sample, redefining each time the training, validation and test subsets, so one can take as our signal efficiency and background rejection the mean values of the powers and sizes of the tests.

The results of one of these training processes for the $\tau^+\tau^-$ QEL signal with $m_{\text{DM}} = 5$ GeV is shown in Fig. 7.15. On the left panel I show the loss function values (blue) and accuracy (red) at each iteration for the training and the validation samples respectively.

7.6. High energy DM neutrino signals

2110 The training stops either when the maximum number of iterations is reached (1000 in
2111 this case) or when the accuracy for the validation sample reaches a certain tolerance
2112 (I chose 10^{-4} as our tolerance). On the right panel I have the distributions for the
2113 predicted probability by the model, separated in true signal (blue) and background
2114 (red) events, for the test sample. One can see that both populations are well separated,
2115 obtaining a power of 44.97% and a size of 0.17% when I require a predicted probability
2116 greater than 0.97.

2117 Applying this criteria for each sample, I obtain the mean signal efficiencies shown in
2118 Fig. 7.16. Notice that the efficiencies for the channel $\tau^+\tau^-$ (blue line) are consistently
2119 lower than the ones for the $b\bar{b}$ channel (red line). This can be due to the fact that, for
2120 each DM mass point, the neutrino spectrum coming from the $b\bar{b}$ annihilation channel is
2121 centered at lower energies when compared to the $\tau^+\tau^-$ spectrum. This directly translates
2122 into more low energy neutrinos undergoing QEL interactions, which give signals that
2123 can be easily separated from the atmospheric background. This explanation also help us
2124 understand why in both cases the signal acceptance drops when the DM mass increases.
2125 In all cases, the background rejection took values between 99.8% to 99.9%. I will assume
2126 a 99.8% background rejection value in all cases to keep our estimation conservative.

2127 7.6.3 Results

2128 In order to estimate the DM-nucleon cross section sensitivities in the present case I need
2129 again to compute the expected number of background events. As I am now separating
2130 events by interaction type Eq. (7.25) does not hold anymore, as in that case I integrated
2131 over the total neutrino-argon cross section. In this instance, the expected background
2132 events for DIS events is approximately given by:

$$N_B^{DIS} \simeq \eta_B^{DIS} \times (4.655 \times 10^3) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right), \quad (7.38)$$

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.17: Projected 90% confidence level upper limit for DUNE (400 kT yr) on the spin-dependent DM-nucleon scattering cross section as a function of m_{DM} , for the annihilation channels $\tau^+\tau^-$ (blue) and $b\bar{b}$ (red) separated by interaction type (up triangles denote DIS interactions whereas down triangles represent QEL interactions). I also show the previous limits from IceCube [9] (solid lines) and the projected sensitivities for Pingu [10] (dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines), as well as the direct detection limits from PICASSO [12] (solid green line) and PICO-60 C₃F₈ [13] (dashed green line).

2133 whereas for QEL events we have:

$$N_B^{QEL} \simeq \eta_B^{QEL} \times (2.248 \times 10^4) \times \left(\frac{\text{exposure}}{400 \text{ kT yr}} \right). \quad (7.39)$$

2134 Now, using these together with Eqs. (7.26) and (7.27) one can obtain the 90% C.L.
 2135 upper limit on the total annihilation rate at equilibrium for both kind of events. Then,
 2136 applying the computed DM-nucleons capture rates I can translate these into limits on
 2137 the DM-nucleon cross section by means of Eqs. (7.2), (7.5) and (7.6).

2138 Fig. 7.17 shows the obtained limits on the SD DM-nucleon cross section for DUNE,
 2139 using the DIS (up triangles) and QEL (down triangles) events both for the $\tau^+\tau^-$ (blue)
 2140 and the $b\bar{b}$ (red) samples, for an exposure of 400 kT yr. I also include the corresponding

7.7. Example: Leptophilic Dark Matter

2141 current limits from IceCube [9] (solid lines), as well as the projected sensitivities of Pingu
2142 [10] (dashed lines) and Hyper-Kamiokande [11] (dash-dotted lines). For comparison, I
2143 also show the reported direct detection limits from PICASSO [12] (solid green line) and
2144 PICO-60 C₃F₈ [13] (dashed green line).

2145 Notice that, for most of the mass range, the limits one can set by using the DIS
2146 events are stronger than those of the QEL interactions, except for the low mass part
2147 of both the $\tau^+\tau^-$ and the $b\bar{b}$ curves where the QEL events dominate. In general, the
2148 expected sensitivity of DUNE for DM masses $\lesssim 25$ GeV surpasses the stronger current
2149 indirect limits. However, experiments like Hyper-Kamiokande are foreseen to have an
2150 overall better sensitivity in this kind of searches, as they have a bigger active volume
2151 and accept a broader energy range.

2152 A pending question is what happens when we add the RES and MEC charged-current
2153 interaction contributions. In that case it would probably be more convenient to split
2154 the samples by final state interaction topologies. Also, another necessary improvement
2155 would be adding a full detector simulation and reconstructions. This will also require
2156 considering the effect of poorly reconstructed events or final states containing neutral
2157 particles such that they mimic the desired topology at the reconstruction level.

2158 7.7 Example: Leptophilic Dark Matter

2159 In general, the capture rate of DM particles by the Sun via interactions with electrons is
2160 several orders of magnitude smaller than the capture via DM-nucleus scattering. Thus,
2161 it would be sub-leading even when nucleon capture is loop suppressed. As I showed in
2162 Fig. 7.2, the capture rate via scattering off electrons only surpasses the capture rates
2163 via DM-nucleons interactions for DM masses $\lesssim 100 - 500$ MeV.

2164 However, if one considers a model where DM-nucleon interactions are forbidden even
2165 at loop level, then electron interactions will be the sole contributor to DM capture in
2166 the Sun. One can describe such scenario where the DM particles couple to leptons but
2167 not to the quark sector using effective operators.

Chapter 7. DM searches with neutrinos from the Sun

2168 In general, assuming that the DM particle is a Dirac fermion, the dimension six
 2169 operators describing the interaction between two DM particles and two leptons can be
 2170 written as:

$$\mathcal{L}_{eff} = G \sum_i (\bar{\chi} \Gamma_\chi^i \chi) (\bar{\ell} \Gamma_\ell^i \ell), \quad (7.40)$$

2171 where $G = 1/\Lambda^2$ is the effective coupling strength, Λ the cut-off of the effective field
 2172 theory and ℓ denotes any lepton. In principle, one should consider all the possible
 2173 Lorentz structures Γ_f^i in order to have a complete set of effective operators.

2174 However, some combinations will induce interactions with nucleons at loop level. As
 2175 we are specifically interested in interactions which forbid any communication with the
 2176 quark sector, I will not consider those [88]. In addition, some of the effective operators
 2177 give rise to velocity-suppressed scattering cross sections between DM particles and
 2178 leptons. I will also neglect those, as the suppression goes with the square of the DM
 2179 halo velocity which in units of the speed of light is $\sim 10^{-6}$.

2180 This way, the only Lorentz tensor structure that do not induce interactions with
 2181 quarks at loop level and gives a contribution to the scattering cross section that is not
 2182 velocity suppress is the axial-axial interaction. The effective Lagrangian is then given
 2183 by:

$$\mathcal{L}_{eff} = \frac{c_A^\chi c_A^\ell}{\Lambda^2} (\bar{\chi} \gamma^\mu \gamma^5 \chi) (\bar{\ell} \gamma_\mu \gamma^5 \ell), \quad (7.41)$$

2184 where c_A^χ and c_A^ℓ are the couplings for the different species. As the DM coupling appears
 2185 as a common factor for any lepton choice, I will redefine the corresponding coupling c_A^ℓ
 2186 to absorb c_A^χ . Also, for simplicity, I will assume that the couplings between the DM
 2187 particles and the leptons are flavour independent, i.e. I have just two couplings, c_A^e for
 2188 charged leptons and c_A^v for neutrinos.

2189 In the case of a scalar DM particle, the lowest order effective interaction with
 2190 leptons happens through a dimension five operator, generating scalar and pseudoscalar
 2191 interactions. However, the former induces interactions with quarks at two loop level
 2192 whereas the latter gives a velocity suppressed scattering cross section.

2193 From the effective Lagrangian in Eq. (7.41) it can be shown that the axial-axial

7.7. Example: Leptophilic Dark Matter

2194 contribution to the scattering cross section for the fermionic DM and a charged lepton
2195 is given by:

$$\sigma_{\text{DM}-e}^{AA} = 3(c_A^e)^2 \frac{m_e^2}{\pi \Lambda^4}. \quad (7.42)$$

2196 If the DM interacts exclusively with fermions, then the only annihilation channels
2197 that will give us a measurable neutrino flux coming out of the Sun are $\tau^+\tau^-$ and $\nu\bar{\nu}$. The
2198 former channel, already explored previously in the more mainstream scenario of the DM
2199 capture via scattering off nucleons, is open only for $m_{\text{DM}} > m_\tau \simeq 1776.86 \pm 0.12$ MeV
2200 [89], a mass region where the solar DM capture by electrons is at least one order of
2201 magnitude smaller than the capture via interactions with nucleons. On the contrary, the
2202 latter allows us to explore a region where the capture rate via scattering off electrons
2203 dominates over the rest.

2204 One downside of focusing in such low mass range is that it falls below the usual
2205 limit of $m_{\text{evap}} \sim 4$ GeV usually explored in the literature. The pretext to explore this
2206 region is the result discussed previously reported in Ref. [4], where DM evaporation in
2207 the Sun for the case of capture via electron scattering could be negligible for masses
2208 as low as $m_{\text{evap}} \sim 200$ MeV. This result is quite sensitive to the high velocity tail of
2209 the DM velocity distribution in equilibrium inside the Sun, and therefore full numerical
2210 simulations would be needed to assess the impact of this effect. However, this falls out of
2211 the scope of our work.

2212 In this case, as I have a specific realisation of the interaction between the DM
2213 and leptons, one can estimate the relic density of our DM for different values of the
2214 couplings and the effective field theory scale Λ . The first step to do so is compute the
2215 self-annihilation cross section. Because I consider cold relics, at the freeze-out time our
2216 DM particles were non-relativistic and so one can expand the annihilation cross section
2217 in terms of the relative velocity v between two annihilating DM particles as [90]:

$$\sigma_{\text{ann}}^{AA}|v| \approx \frac{1}{2\pi\Lambda^4} \sum_\ell \left(c_A^\ell\right)^2 m_\chi^2 \sqrt{1 - \frac{m_\ell^2}{m_\chi^2} \left[\frac{m_\ell^2}{m_\chi^2} + \frac{1}{12} \left(2 - \frac{m_\ell^2}{m_\chi^2}\right) v^2 \right]}, \quad (7.43)$$

Chapter 7. DM searches with neutrinos from the Sun




Figure 7.18: Left panel: Projected 90% confidence level sensitivity of DUNE (400 kT yr) to the scale Λ of an EFT containing only leptophilic DM axial-axial interactions (blue line). Right panel: In both cases the corresponding limits from DarkSide-50 [14] (dotted green line) and XENON1T [15] (dashed red line) are also shown, together with the configurations for which the correct relic density is achieved (black line), all for the coupling values $c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$.

2218 where the sum includes all the possible lepton final states with mass m_ℓ .
 2219 Solving the Boltzmann equation for the evolution of the DM density gives as a
 2220 solution a relic density of:

$$\Omega_\chi h^2 \approx \frac{(1.04 \times 10^9) x_F}{M_{Pl} \sqrt{g_*} (a + 3b/x_F)}, \quad (7.44)$$

2221 where $x_F = m_\chi/T_F$ being T_F the freeze-out temperature, g_* the number of relativistic
 2222 degrees of freedom at freeze-out and a and b the terms in the annihilation cross section
 2223 expansion $\sigma_{ann}|v| \approx a + bv^2 + \mathcal{O}(v^4)$. Using the current best fit for the relic DM density
 2224 $\Omega_\chi h^2 = 0.1198 \pm 0.0012$ [91] one can use these relations to compute the required effective
 2225 theory scale Λ at which the correct density is achieved for any combinations of m_χ and
 2226 c_A^ℓ .

2227 As discussed before, in the low DM mass region QEL interactions dominate. Moreover,
 2228 if I focus on direct annihilation to neutrinos, the energy of the muon neutrino flux is
 2229 known as it must be equal to the mass of the DM particle, $E_\nu = m_\chi$. That way, now
 2230 I do not need to use Eq. (7.33) in order to estimate the momentum transfer to the

7.7. Example: Leptophilic Dark Matter

2231 remnant nucleus, I can simply take:

$$\vec{p}_N = \hat{p}_\nu m_\chi - \vec{p}_\mu - \vec{p}_p. \quad (7.45)$$

2232 To estimate the signal efficiency and background rejection for this case I used again
2233 the MLP classifier from `scikit-learn`, using the same specifications as before. The
2234 only difference now is that I add also the reconstructed neutrino energy as one of the
2235 features to train the classifier with, because the characteristic monoenergetic flux for
2236 each m_χ value will help to distinguish between signal and background events.

2237 In this case, for masses below ~ 500 MeV I obtain a signal efficiency close to unity
2238 while keeping a background rejection of 99.9%. For bigger values of the mass, the signal
2239 efficiency drops significantly if I require to keep the background acceptance under 0.01%.
2240 However, because this kind of search is dominated by the background, sacrificing the
2241 signal acceptance to keep the background rejection to a minimum enhances the reach
2242 of the analysis. This way, for DM masses of the order of $m_\chi \sim 1$ GeV I end up with
2243 efficiencies as low as 1%.

2244 Now, estimating the number of background events using Eq. (7.39) one can go on
2245 and apply Eqs. (7.26) and (7.27) together with Eq. (7.42) to derive the sensitivity of
2246 DUNE to this kind of model. Fig. 7.18 (left panel) shows the potential reach of DUNE
2247 to constrain the EFT scale Λ this model containing only leptophilic DM axial-axial
2248 interactions (blue line), for a choice of couplings $c_A^e = 10^3$ and $c_A^\nu = 10^{-2}$. I also included
2249 the current limits on the DM-electron scattering cross section from DarkSide-50 [14]
2250 (dotted green line) and XENON1T [15] (dashed red line), reworked with Eq. (7.42) to
2251 show their implications for the EFT scale. The values of Λ for which the correct DM relic
2252 density value is achieved for each mass are also shown (black line). This tells us that,
2253 for that specific choice of couplings, DUNE would be sensitive to DM configurations
2254 allowed by the relic density constraint up to a mass of $m_\chi \sim 400$ MeV.

2255 In Fig. 7.18 (right panel) I show the same upper limits but for the DM-electron
2256 scattering cross section. From this view one can see that DUNE would be able to

Chapter 7. DM searches with neutrinos from the Sun

offer complementary information to the low energy DM-electron interaction searches performed by direct detection experiments, in a slightly higher mass range.

With the present example, although it focuses on a very specific realisation of the DM interactions, I show the potential of DUNE to constrain exotic DM scenarios. Thanks to its low backgrounds and superb angular resolution DUNE will be able to help with the systematic searches for dark sectors physics.

2263 Chapter 8

2264 Particle ID in GArSoft

2265 ND-GAr is a magnetised, high-pressure gaseous argon TPC (HPgTPC), surrounded by
2266 an electromagnetic calorimeter (ECal) and a muon detector (commonly refer to as μ ID).
2267 A detailed discussion on the requirements, design, performance and physics of ND-GAr
2268 can be found in the DUNE ND CDR [48] and the ND-GAr whitepaper (cite).

2269 In DUNE Phase II ND-GAr will fulfill the role of TMS, measuring the momentum
2270 and sign of the charged particles exiting ND-LAr. Additionally, it will be able to measure
2271 neutrino interactions inside the HPgTPC, achieving lower energy thresholds than those
2272 of the ND and FD LArTPCs. By doing so ND-GAr will allow to constrain the relevant
2273 systematic uncertainties for the LBL analysis even further.

2274 The goal of the present chapter is to review the requirements that the physics program
2275 of DUNE impose on ND-GAr, present the current status of its design and describe the
2276 GArSoft package, its simulation and reconstruction software.

2277 As decided during the DUNE Phase II workshop in June 2023 [reference], we want
2278 to build ND-GAr physics case by showing:

- 2279 • That ND-GAr can constrain systematic uncertainties that ND-LAr might miss.
- 2280 • The impact on the neutrino oscillation results if such systematic uncertainties are
2281 missed.
- 2282 • That ND-GAr is necessary to reach DUNE's main physics goals.

Chapter 8. Particle ID in GArSoft

2283 This way, the design of ND-GAr will be physics driven.

2284 In order to study the effects of final state interactions (FSI) in CC interactions,
2285 ND-GAr should be able to measure the spectrum of protons and charged pions at low
2286 energies. ND-GAr also needs to be able to measure the pion multiplicity, specially for
2287 energies above 100 MeV as at these energies the pions shower in the LAr, to inform the
2288 pion mass correction in the ND and FD LArTPCs.

2289 In order to correctly identify electrons, muons, pions, kaons and protons ND-GAr
2290 can use a combination of: dE/dx measurements in the HPgTPC, E_{ECAL}/p using the
2291 ECAL total energy and the momentum obtained from magnetic spectroscopy in the
2292 HPgTPC and penetration information through the ECAL and muon tagger.

2293 8.1 dE/dx measurement in the TPC

2294 Among the parameters extracted from the track fitting, ionisation is particularly useful
2295 for particle identification, as it is a function of the particle velocity. Although for the
2296 case of relativistic particles this dependence is not very strong, measuring the track on
2297 a large number of points may allow us to estimate the amount of ionisation accuratel.
2298 This, paired with a measurement of the momentum, may allow us to identify the particle
2299 type.

2300 The first calculation of the energy loss per unit length of relativistic particles using
2301 a quantum-mechanical treatment is due to Bethe [?]. Using this approach, the mean
2302 ionisation rate of a charged particle traveling through a material medium is (using
2303 natural units $G = \hbar = c = 1$):

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{4\pi Ne^4}{m_e \beta^2} z^2 \left(\log \frac{2m_e \beta^2 \gamma^2}{I} - \beta^2 \right), \quad (8.1)$$

2304 where N is the number density of electrons in the medium, e the elementary charge, m_e
2305 is the electron mass, z the charge of the particle in units of e , β is the velocity of the
2306 particle, $\gamma = (1 - \beta^2)^{-1}$ and I denotes the effective ionisation potential averaged over
2307 all electrons. This relation is known as the Bethe-Bloch formula.

8.1. dE/dx measurement in the TPC

From Eq. (8.1) one can see that the ionisation loss does not depend explicitly on the mass of the charged particle, that for non-relativistic velocities it falls as β^{-2} , then goes through a minimum and increases as the logarithm of γ . This behaviour at high velocities is commonly known as the relativistic rise. The physical origin of this effect is partly due to the fact that the transverse electromagnetic field of the particle is proportional to γ , therefore as it increases so does the cross section.

It was later understood that the relativistic rise could not grow indefinitely with γ . A way to add this feature in the Bethe-Bloch formula is by introducing the so-called density effect term. It accounts for the polarisation effect of the atoms in the medium, which effectively shield the electromagnetic field of the charged particle halting any further increase of the energy loss [?]. Denoting the correction as $\delta(\beta)$, one can rewrite Eq. (8.1) as:

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{4\pi Ne^4}{m_e \beta^2} z^2 \left(\log \frac{2m_e \beta^2 \gamma^2}{I} - \beta^2 - \frac{\delta(\beta)}{2} \right). \quad (8.2)$$

In general, the form of $\delta(\beta)$ depends on the medium and its state of aggregation, involving the usage of tabulated parameters and implicit relations [?].

Another standard method to compute the amount of ionisation a charged particle produces is the so-called photo-absorption ionisation (PAI) model proposed by Allison and Cobb [?]. Within their approach, the mean ionisation is evaluated using a semiclassical calculation in which one characterises the continuum material medium by means of a complex dielectric constant $\epsilon(k, \omega)$. However, in order to model the dielectric constant they rely on the quantum-mechanical picture of photon absorption and collision. Therefore, in the PAI model the computation of the ionisation loss involves a numerical integration of the measured photo-absorption cross-section for the relevant material.

In a particle physics experiment, the typical way of determining the energy loss per unit length as a function of the particle velocity is studying identified particles over a range of momenta. Once we have established this relation we can use it for other, unknown particles. In this sense, it makes sense to have a regular mathematical expression for this relation that one can use.

Chapter 8. Particle ID in GArSoft

2335 It happens that neither the Bethe-Bloch theory nor the PAI model from Allison and
2336 Cobb offer a close mathematical form for the ionisation curve. This is the reason why a
2337 full parametrisation of the ionisation curves can be useful. A parametrisation originally
2338 proposed for the ALEPH TPC [?] and later used by the ALICE TPC [?] group that
2339 manages to capture the features of the ionisation energy loss is:

$$f(\beta\gamma) = \frac{P_1}{\beta^{P_4}} \left(P_2 - \beta^{P_4} - \log \left[P_3 + \frac{1}{(\beta\gamma)^{P_5}} \right] \right), \quad (8.3)$$

2340 where P_i are five free parameters. Hereafter, we will refer to Eq. (8.3) as the ALEPH
2341 dE/dx parametrisation.

2342 8.1.1 Energy calibration

2343 In order to obtain the amount of energy loss by a charged particle due to ionisation
2344 in our TPC we need to determine the conversion between the charge deposited in our
2345 readout planes and the actual energy depositions. This procedure is known as energy
2346 calibration.

2347 In a general, the first step of the calibration involves a non-uniformity correction,
2348 to make sure that the detector response is uniform throughout the TPC. These are
2349 typically divided into three categories, non-uniformities in the transverse YZ plane,
2350 non-uniformities along the drift direction X and variations of the detector response
2351 over time (would not apply to us as the detector is not built yet). These would correct
2352 for effects such as electron diffusion and attenuation, space charge effects or channel
2353 misconfiguration. However, because at the moment I am only interested in making sure
2354 we recover a sensible result from our simulation, I will not apply uniformity corrections
2355 to our charge deposits.

2356 Other effects, like electron-ion recombination or ADC saturation, lead to a non-linear
2357 relation between the observed charge and the deposited energy in the detector, with the
2358 observed readout charge saturating at high ionisation energies. In this case, because we
2359 are dealing with gaseous argon and therefore recombination is not as important as in

8.1. dE/dx measurement in the TPC




Figure 8.1: Left panel: distribution of the fraction of Geant4-level energy deposits per track with residual range less than 20% of the total track length, for the isotropic proton sample. Right panel: distribution of the ionisation per unit length of the energy deposits in the proton sample after removing the tracks with less than 30% of their energy deposits in the last 20% of the track.

2360 liquid, we do not simulate recombination effects in the TPC. Even so, the simulation of
 2361 the electronic response will still introduce charge saturation, and one needs to correct
 2362 for it in order to obtain the exact amount of energy loss due to ionisation.

2363 By default, the track fitting algorithm in GArSoft provides a `TrackIonization`
 2364 object associated to each reconstructed track. It contains two collections of charge
 2365 deposits, one for each fitting direction, consisting on pairs of charge values (dQ , in ADC)
 2366 and step sizes (dx , in cm).

2367 In order to estimate the ionisation loss in the ND-GAr TPC, I have used an MC
 2368 sample consisting of single, isotropic protons propagating in the TPC. The starting points
 2369 of the protons were sampled inside a $50 \times 50 \times 25$ cm box centered at $(100, -150, 1250)$,
 2370 and their momenta are uniformly distributed in the range $0.25 - 1.75$ GeV. I ran the
 2371 simulated sample through GArSoft's default detector simulation and reconstruction, and
 2372 then a custom analyser module that extracts the ionisation data together with other
 2373 reconstructed track information from the Kalman fit.

2374 For studying the energy loss of the protons I select the reconstructed tracks that
 2375 range out (i.e. slow down to rest) inside the TPC. A characteristic feature of the energy
 2376 loss profile of any stopping ionising particle is the so-called Bragg peak, a pronounced

Chapter 8. Particle ID in GArSoft




Figure 8.2: Left panel: distribution of the reconstructed ionisation charge per unit length for our MC stopping proton sample. The different colors indicate how many consecutive dQ/dx pairs were grouped together. Right panel: distribution of the median change in dQ/dx per track after $N_{group} = 4$ clusters were reclustered together.

peak that occurs immediately before the particle comes to rest. From Eq. (8.1) we can see that this behaviour is expected, as the energy loss for non-relativistic particles is inversely proportional to β^2 . In data, a way of identifying the Bragg peak, and thus select the stopping particles, is checking the number of energy deposits towards the end of the track. In this case, I count the fraction of the Geant4 simulated energy deposits with a residual range value (the distance from a given energy deposit to the last deposit in the track trajectory) less than a 20% of the corresponding track length¹. The distribution of this fraction of energy deposits for our proton sample is shown in Fig. 8.1 (left panel). We can clearly see two well separated peaks in this distribution, one centered at 0.2 and another, narrower, one centered at a higher value. The first one corresponds to non-stopping protons, as in that case the number of energy deposits towards the end of the track is uniformly distributed due to the absence of the Bragg peak. In that way, I apply a cut in this distribution, requiring that at least 30% of the simulated energy deposits sit in the last 20% of the tracks, to ensure that the Bragg peak is present.

Fig. 8.1 (right panel) shows the distribution of the energy loss per unit length for the Geant4 simulated energy deposits of the selected stopping protons. We can see that it

¹As we are applying this selection at the Geant4 level we could have simply selected the stopping protons using the `EndProcess` labels from the simulation. However, the Bragg peak identification method displayed here could serve as a starting point for a selection of stopping protons in real data.

8.1. dE/dx measurement in the TPC

2394 follows the expected shape of a Landau distribution, which describes the fluctuations of
2395 the ionisation energy losses [?]. This distribution has a characteristic asymmetric PDF,
2396 with a long right tail that translates into a high probability for high-energy ionisation
2397 losses. The origin of these fluctuations is mainly the possibility of transferring a high
2398 enough energy to an electron, so it becomes a ionising particle itself.

2399 Now, from the point of view of the reconstruction, the objects that we have available
2400 to extract the ionisation information for the different reconstructed tracks are the
2401 collections of dQ and dx pairs, as stated before. The dQ values come from adding up
2402 the amplitude of all the reconstructed hits in a cluster, which is the input object to the
2403 Kalman fit.

2404 Fig. 8.2 (left panel) shows the distribution of the ionisation charge deposits per
2405 unit length for the track in the stopping proton sample (blue line). As one can notice,
2406 this distribution does not resemble the expected shape of the Landau PDF. This
2407 distribution peaks sharply at 0 and has a heavy tailed behaviour. Notice, however, how
2408 the distribution changes its shape as we group together N_{group} consecutive charge deposit
2409 pairs (red, purple and green lines). The distribution in the $N_{group} = 4$ case already has
2410 a shape which resembles that of the Geant4-level ionisation per unit length, so I will
2411 proceed using this amount of reclustering for the reconstruction-level depositions.

2412 An extra factor I need to account for, when reclustering is applied, is how the overall
2413 dQ/dx per track changes. To do so, we can look at the ratio between the median dQ/dx
2414 after and before the reclustering. Fig. 8.2 (right panel) shows the median enhancement
2415 in dQ/dx per track for the stopping proton sample in the case $N_{group} = 4$. Fitting a
2416 Landau distribution convolved with a Gaussian², I estimate the most probable value of
2417 this ratio to be $G_{group} = 2.24 \pm 0.05$.

2418 At this point, I am left with determining the conversion between the charge deposits
2419 per unit length dQ/dx and the energy deposits per unit length dE/dx . To this end, we
2420 need a way of comparing the two. I can use the residual range z to get a prediction of

²In the literature, this distribution is often referred to as Landau+Gaussian or langau. In the following, I will use LanGauss to refer to such PDF.

Chapter 8. Particle ID in GArSoft




Figure 8.3: Distribution of the Geant4-simulated energy losses per unit length versus residual range for the stopping proton sample. The overlaid points represent the fitted most probable value of the dE/dx distribution in each residual range bin, whereas the curve is their best fit to the Bragg-Kleeman formula from Eq. (8.4).

2421 the most probable dE/dx by using the following empirical parametrisation [?]:

$$\frac{dE}{dx}(z) = \frac{z^{\frac{1}{p}-1}}{p\Lambda^{\frac{1}{p}}}, \quad (8.4)$$

2422 which is quoted in the literature as the Bragg-Kleeman formula. In order to obtain the
 2423 p and Λ parameters I perform a fit using the energy losses and the residual ranges given
 2424 by the Geant4 stage of our proton sample.

2425 Within our simulation, the residual range is sampled with a maximum size of
 2426 5 mm. Therefore, to perform the fit to the Bragg-Kleeman formula, we can use a
 2427 fine-grained residual range binning. For each of the residual range bins I extract the
 2428 dE/dx distribution and fit it to a LanGauss distribution, to obtain the value of the
 2429 most probable dE/dx in the bin together with a statistical uncertainty. I then fit Eq.
 2430 (8.4) to these most probable values and the centres of the residual range bins. This

8.1. dE/dx measurement in the TPC




Figure 8.4: Fitted most probable dQ/dx values for each dE/dx bin (red points), obtained from the stopping proton sample. The overlaid curve (black line) represents the best fit to the logarithmic calibration function from Eq. (8.5).

procedure is depicted in Fig. 8.3, where I show the distribution of the energy loss per unit length versus the residual range, together with the most probable dE/dx values and their uncertainty in each bin (red points) and the curve with the best fit of the Bragg-Kleeman relation to those values (black line). The best fit is obtained for the parameter values $p = 1.8192 \pm 0.0005$ and $\Lambda = 0.3497 \pm 0.0008$ cm/MeV³.

Having an analytical expression that relates the residual range to dE/dx , I can take our reconstruction-level residual ranges from the stopping proton sample and compute the most probable energy loss associated.

In order to parametrise the charge saturation, we can use the following logarithmic

³These strange units for Λ come from dimensional analysis, just to keep the Bragg-Kleeman formula (8.4) consistent.

Chapter 8. Particle ID in GArSoft

2440 function inspired by the modified box model for recombination:

$$\frac{dE}{dx} = \frac{e^{\frac{dQ}{dx}B\frac{W_{ion}}{G_{group}C}} - A}{B}, \quad (8.5)$$

2441 where A and B are the calibration parameters we need to determine, W_{ion} is the average
 2442 energy to produce an electron-ion pair, G_{group} is the gain from the reclustering discussed
 2443 above and C is the calibration constant to convert number of electrons to ADC counts,
 2444 commonly refer to as gain (also to be obtained in the fit). In this case, I use a value for the
 2445 electron-ion production energy of $W_{ion} = 26.4$ eV [?]. This value, used in our simulation
 2446 as well, was measured for gaseous argon in normal conditions, and therefore should
 2447 be checked in the future to describe correctly the high-pressure argon-CH₄ mixture of
 2448 ND-GAr.

2449 For the calibration fit I follow a procedure similar to the previous one for Eq. (8.4).
 2450 Binning the dE/dx range, I fit a LanGauss distribution to the corresponding dQ/dx
 2451 distribution to obtain the most probable value. The resulting data points (red bars) are
 2452 shown in Fig. 8.4 (top panel), the horizontal error bars depict the width of the dE/dx
 2453 bin whereas the vertical bars represent the error associated to the most probable value
 2454 estimation. A fit to the logarithmic function in Eq. (8.5) is also shown (black line).
 2455 For this I weighted the data points using the inverse of their relative error, obtaining a
 2456 reduced chi-square value of $\chi^2/ndf = 2.22 \times 10^{-6}$. The best fit parameters I found from
 2457 this fit are $A = 0.883 \pm 0.064$, $B = 5.6 \pm 1.3$ cm/MeV and $C = 4.94 \pm 0.49$ ADC/e. In
 2458 Fig. 8.4 (bottom panel) I show the residuals between the data points and the fit.

2459 The value for the gain I obtained from the fit is in reasonable agreement with our
 2460 expectation. This value is set in GArSoft to 5 ADC/e by default.

2461 One interesting thing to check is what induces this non-linear relation between charge
 2462 and energy. The only effects that modify the amount of electrons reaching the readout
 2463 planes in the simulation are the transverse diffusion and the finite electron lifetime.
 2464 Once the electrons reach the readout chambers, the pad response functions are applied,
 2465 together with an electrons-to-ADC conversion and the ADC saturation limit.

8.1. dE/dx measurement in the TPC




Figure 8.5: Fitted most probable dQ/dx values for each dE/dx bin for three different ADC bit limits, 10 (blue points), 12 (default, yellow points) and 16-bit (red points).

Table 8.1: Calibration parameters obtained from the fit of the ND-GAr simulated stopping proton sample to the calibration function from Eq. (8.5). The fits were performed for the 10, 12, and 16-bit ADC limits.

	χ^2/ndf	Best fit $\pm 1\sigma$		
		A	B (cm/MeV)	C (ADC/e)
10-bit	$1.83 \times 10^{-6}/12$	-9.3 ± 3.9	270 ± 69	27.1 ± 5.4
12-bit	$2.67 \times 10^{-5}/12$	0.883 ± 0.064	5.6 ± 1.3	4.94 ± 0.49
16-bit	$1.44 \times 10^{-5}/12$	0.949 ± 0.024	3.53 ± 0.58	4.52 ± 0.29

By default, GArSot applies a 12-bit ADC limit, which can be changed in the simulation configuration. However, it can only be increased up to 16-bit, as we represent the ADC collection as a `std::vector<short>`. This way, I tried to change the saturation parameter to see how it affects the relation between reconstructed charge and energy. Fig. 8.5 shows a comparison between the most probable dQ/dx for 10, 12 and 16-bit ADC limits. As expected, the lower the limit is the sooner the charge saturates. For higher ADC limits the relation between energy and charge remains linear up to higher dE/dx values, but even for the 16-bit limit the saturation is noticeable for values $\gtrsim 0.5$ MeV/cm.

In Tab. 8.1 I also show the results of fitting the samples with 10 and 16-bits ADC

Chapter 8. Particle ID in GArSoft




Figure 8.6: Top panel: area normalised dE/dx distributions for the true (solid grey) and the reconstructed energy deposits in the stopping proton sample, both after applying the calibration (blue) and the calibration and the normalisation correction (yellow). Also shown is the distribution obtained by applying a correction factor to the dQ/dx values but not the calibration (red). Bottom panel: fractional residuals for the uncorrected (blue), corrected (yellow) and uncalibrated (red) samples.

limits to the calibration function from Eq. (8.5), using the weights based on their relative error as described previously. One interesting feature to notice is how different the best fit points look for the 10-bit ADC saturation when compared to the other two, which are consistent with each other.

At this point we can compare the dE/dx distribution one gets from Geant4, i.e. the true energy loss distribution, and the distribution I found by applying the calibration function to our collection of reconstructed dQ/dx values. Fig. 8.6 (top panel) shows the true (solid grey) and reconstructed (blue, labeled as uncorrected) distributions together. The dashed vertical lines indicate the region of validity of the calibration fit, i.e. the left

8.1. dE/dx measurement in the TPC

and right edges of the first and last dE/dx bin respectively. Notice that these histograms are area-normalised, as the total number of true energy deposits is much higher than the number of reconstructed charge deposits. This is due to a combination of effects, like the finite spatial resolution of the detector, the hit clustering used in the track fitting and the reclustering we have applied here.

The two distributions are significantly different. That can be seen clearly when looking at the fractional residuals, shown in Fig. 8.6 (bottom panel). In particular, the position of the peak is off, which could bias the mean energy loss predictions. It seems like the difference between these may be due to an overall scaling factor. One possibility is to scale the most probable value of the reconstructed distribution to the most probable value predicted by Geant4. I do this by fitting both distributions using a LanGauss function, obtaining $dE/dx_{MPV, true} = 0.1145 \pm 0.0005$ MeV/cm and $dE/dx_{MPV, reco} = 0.0928 \pm 0.0005$ MeV/cm for the true and reconstructed most probable values respectively. These can be translated into a scaling factor $S = 0.579 \pm 0.006$.

The result of applying the scaling correction can be seen in Fig. 8.6 (top panel). The corrected dE/dx distribution (yellow, labeled as corrected) peaks around the same value the true distribution does, as expected. Moreover, the high energy region is also slightly better described. For low ionisations, below the lower limit of the calibration fit, the differences between true and reconstructed are still significant. This low energy excess may be migration of some events from the peak region. The overall effect of the correction can be seen in the fractional residual plot in Fig. 8.6 (bottom panel).

One can also check what happens if instead of applying the logarithmic calibration we simply scale the dQ/dx distribution (post reclustering) to have the same most probable value as the true dE/dx distribution. In this case, following an analogous procedure to the one described earlier, I found the scaling factor $S_{uncalibrated} = 0.414 \pm 0.002$ MeV/ADC⁴. The resulting distribution (red, labeled as uncalibrated) is also shown in in Fig. 8.6 (top panel). The behaviour of the new distribution is similar to the corrected case at low energy losses, around the peak of the true distribution, but it is worse at describing the

⁴Notice that now the scaling factor is not dimensionless, as it acts more like a conversion factor here.

Chapter 8. Particle ID in GArSoft




Figure 8.7: Left panel: fractional residuals between the true and the corrected dE/dx means (blue) and the 60% truncated means (yellow), for each event in the stopping proton sample. Right panel: fractional residuals between the true and the uncorrected (blue), corrected (yellow) and uncalibrated (red) dE/dx 60% truncated means, for each event in the stopping proton sample.

2512 high energy tail. This is expected, it is in the high ionisation regime where saturation
 2513 effects apply and therefore calibration is needed.

2514 8.1.2 Truncated dE/dx mean

2515 Once we have a collection of dE/dx values for each reconstructed track, we can compute
 2516 the corresponding most probable ionisation loss per unit length of the particle. This
 2517 is the value predicted by the Bethe-Bloch or the PAI models, and together with a
 2518 measurement of the momentum it allows for particle identification.

2519 However, estimating the most probable dE/dx value for each reconstructed track
 2520 is not a trivial task. As mentioned before, the dE/dx distributions follow Landau-like
 2521 distributions. Therefore, one should perform e.g. a LanGauss fit to correctly estimate
 2522 the most probable values. Automating this kind of fits is often problematic, as they
 2523 usually incur in convergence problems. Moreover, the reconstructed dE/dx distributions
 2524 we obtain tend to have relatively small statistics, which may also produce poor fits. In
 2525 practice, doing these unsupervised fits may degrade our performance, and a more robust
 2526 method is preferred.

2527 A possibility could be taking the mean of the reconstructed dE/dx distribution for

8.1. dE/dx measurement in the TPC




Figure 8.8: Estimated values of the mean dE/dx bias (left panel) and resolution (right panel) obtained using the corrected data from the stopping proton sample, for different values of the truncation factor.

each particle. The problem with this approach is that the high energy Landau tail, combined with our limited statistics, can induce large fluctuations in the computation of the mean. Imagine you have two protons with the same kinetic energy, but due to reconstruction problems in one case you did not get as many charge deposits reconstructed in its high ionisation loss region. If you do not remove the tails the computed dE/dx means will be significantly different.

In order to avoid those fluctuations, one can compute the mean of a truncated dE/dx distribution instead. By keeping only a given fraction of the lowest energy deposits we obtain an estimate of the mean energy loss that is more resilient to reconstruction inefficiencies and statistical effects. In Fig. 8.7 (left panel) I show a comparison between the $\langle dE/dx \rangle$ computed by taking the mean of the full distribution (blue line) and the 60% lowest energy clusters (yellow line), for the stopping proton sample. The fractional residuals are computed for each proton, taking the corresponding means using their collections of true and reconstructed energy deposits. One can see that using the simple mean translates into a high bias and uncertainty in the $\langle dE/dx \rangle$ estimation, whereas applying the truncation reduces both significantly.

Additionally, I performed a comparison between the 60% truncated mean dE/dx obtained using the different calibration methods discussed earlier, namely the uncorrected

Chapter 8. Particle ID in GArSoft

2546 (blue), corrected (yellow) and uncalibrated (red) distributions. The results are shown
 2547 in Fig. 8.7 (right panel). While the widths of these distributions are similar, the bias
 2548 obtained for the corrected sample, i.e. calibration function and correction factor applied,
 2549 is a factor of ~ 2 lower than in the uncalibrated case and almost three times smaller
 2550 than for the uncorrected sample.

2551 The next step is to optimise the level of truncation we are going to apply to our
 2552 data. To do so, I used different truncation factors, i.e. the percentage of energy-ordered
 2553 reconstructed energy deposits we keep to compute the mean, on the corrected dE/dx
 2554 sample of the stopping protons. Then, following the same procedure of computing the
 2555 fractional residuals as before, I fitted the resulting histograms using a double Gaussian
 2556 function. This is simply the sum of two Gaussian functions of the type:

$$g(x; \mu, \sigma, A) = A e^{-\frac{(x-\mu)^2}{2\sigma^2}}. \quad (8.6)$$

2557 I do not add the classical normalisation factor of the Gaussian, $1/\sqrt{2\pi}\sigma$, therefore
 2558 the amplitude A simply represents the maximum of the function. One of the two
 2559 Gaussian functions describes the core part of the distribution, while the other captures
 2560 the behaviour of the tails.

2561 For each truncation factor, I look at the bias and the resolution I obtain. I define
 2562 these as the weighted means of the corresponding parameters in the fits:

$$\bar{x} = \frac{A_{core} x_{core} + A_{tail} x_{tail}}{A_{core} + A_{tail}}, \quad (8.7)$$

2563 where A_{core} and A_{tail} are the amplitudes of the core and tail distributions respectively
 2564 and x is either the mean μ or the width σ of said distributions.

2565 Fig. 8.8 shows the bias (left panel) and the resolution (right panel) I obtained for
 2566 the stopping proton sample, using different values of the truncation. From these, it
 2567 can be seen that a truncation factor of 50% minimises the bias in the estimation, while
 2568 70% gives the best resolution. That way, I settled on the intermediate value of 60%

8.1. dE/dx measurement in the TPC




Figure 8.9: Examples of the truncated mean dE/dx LanGauss fits for various $\beta\gamma$ bins, from a simulated FHC neutrino sample.

2569 truncation, which yields a $\langle dE/dx \rangle$ resolution of 5.00 ± 0.08 % for stopping protons.

2570 8.1.3 Mean dE/dx parametrisation

2571 Now that we have a way to estimate the mean energy loss of a particle in the HPgTPC,
 2572 we can determine the value of the free parameters in the ALEPH formula, Eq. (8.3).
 2573 For this, I used a sample of 10^5 reconstructed FHC neutrino events inside ND-GAr. In
 2574 this case I cannot use the stopping proton sample, as we need to cover the full kinematic
 2575 range of interest for the neutrino interactions in our detector.

2576 The original data does not contain an estimation of the velocity of the tracks, instead
 2577 the tracks have a value for the reconstructed momentum and the associated PDG code
 2578 of the Geant4-level particle that created the track. Therefore, one can select some of the

Chapter 8. Particle ID in GArSoft




Figure 8.10: Resulting one and two dimensional projections of the posterior probability distributions of the ALEPH $\langle dE/dx \rangle$ parameters obtained by fitting the 60% truncated mean dE/dx values from a FHC neutrino sample in ND-GAr. The vertical dashed lines in the 1D distributions represent the 16th, 50th and 84th percentiles.

2579 particles in the data, in this case I selected electrons, muons, pions and protons, and
 2580 compute β and γ using the reconstructed momentum and their mass. In terms of $\beta\gamma$
 2581 the mean dE/dx does not depend on the particle species, so one can consider all the
 2582 dataset as a whole. For this fit, I will express β in terms of the $\beta\gamma$ product as:

$$\beta = \frac{\beta\gamma}{\sqrt{1 + (\beta\gamma)^2}}, \quad (8.8)$$

2583 which can be easily proven from the definition of γ .

2584 Next, I bin the data in $\beta\gamma$. I chose a fine binning so as to capture the different

8.1. dE/dx measurement in the TPC

2585 features of the ionisation curve. Instead of fixing the bin width, I select them so each one
2586 has approximately the same statistics. Then, for each $\beta\gamma$ slice, I compute the median
2587 and the interquartile range (IQR) of the $\langle dE/dx \rangle$ distribution. Using these, I make a
2588 histogram in the range [median – IQR, median + 5 IQR], which I fit to a LanGauss
2589 function in order to extract the MPV. Using this range accounts for the asymmetric
2590 nature of the distributions, while also helps avoiding a second, lower maximum present
2591 at low $\beta\gamma$, probably a result of reconstruction failures.

2592 A few examples of these fits are shown in Fig. 8.9. The chosen values of $\beta\gamma$ sit in
2593 very distinct points along the $\langle dE/dx \rangle$ curve, going from the high ionisation region at
2594 low velocities (top left panel), to the minimum point (top right panel), the beginning of
2595 the relativistic rise (bottom left panel), and the plateau produced by the density effect
2596 (bottom right panel).

2597 I used the resulting most probable $\langle dE/dx \rangle$ values and the centres of the $\beta\gamma$ bins as
2598 the points to fit to the ALEPH formula. For this particular fit I used the least-squares
2599 method to get a first estimation of the ALEPH parameters. Applying some uniform
2600 priors, I then used these values as the starting point of a 100000 steps MCMC. In
2601 Fig. 8.10 I show the posterior probability distributions I obtain for each parameter.
2602 The reported best fit points are based on the 16th, 50th, and 84th percentiles in the
2603 marginalised distributions.

2604 The resulting fit (black line), compared to the data points (red points) and the
2605 underlying distribution is shown in Fig. 8.11 (top panel). The overall fit is good, with a
2606 reduced chi-squared of $\chi^2/ndf = 1.02$. However, there are some regions where the fit
2607 does not describe the data correctly, like the very low $\beta\gamma$ regime, where the fit severely
2608 underestimates for energy losses $\gtrsim 50$ keV/cm, and the start of the relativistic raise,
2609 where we have a slight overestimation. This is a result of those points having a larger
2610 uncertainty when compared to the ones around the dip or the plateau areas. These
2611 differences can be better seen in the residual plot, Fig. 8.11 (bottom panel).

Chapter 8. Particle ID in GArSoft




Figure 8.11: Truncated mean dE/dx obtained for the FHC neutrino sample as a function of the $\beta\gamma$ product (upper panel). Also shown are the fitted most probable values for each $\beta\gamma$ bin (red points) and the best fit obtained using the ALEPH parametrisation (black line). The residuals resulting from the fit are shown in the lower panel.

2612 8.1.4 Proton identification

2613 8.2 Muon and pion separation in the ECal and MuID

2614 8.2.1 Track-ECal matching

2615 One of the main players in the muon and pion separation is the way we associate clusters
 2616 in the ECal to reconstructed tracks in the TPC. Missing some associations or making
 2617 wrong ones can bias the ECal quantities that we can use for classifying particles. The
 2618 current algorithm in GArSoft provides precise associations, i.e. most of the associations
 2619 that it produces are correct, but it appears to miss an important number of associations
 2620 (at least when using the default configuration).

2621 The current TPC track-ECal cluster association algorithm is divided in four parts.

8.2. Muon and pion separation in the ECal and MuID




Figure 8.12: Estimated values of the mean dE/dx bias (left panel) and resolution (right panel) obtained using the corrected data from the stopping proton sample, for different values of the truncation factor.

- 2622 It first checks whether the track end point fulfils certain conditions to be extrapolated.
 2623 There are two cut values in this step, one for the drift direction and other radial.
 2624 If the point can be extrapolated, the code computes the coordinates of the centre
 2625 of curvature using the Kalman fit estimates at the track end ($y, z, 1/R, \phi, \tan\lambda$). It
 2626 then compares the distance between this and the cluster in the (z, y) plane with R . This
 2627 introduces another cut in the perpendicular direction.
 2628 The next step is different for clusters in the barrel or in one of the end caps. If it
 2629 is a barrel cluster the algorithm extrapolates the track up to the radial distance of the
 2630 cluster. There are three possible outcomes, the extrapolated helix can cut the cylinder
 2631 of radius r_{clus} two, one or zero times. I get the cut point that is closer to the cluster and
 2632 check that it is either in the barrel or the end caps. Computing the difference between
 2633 the x coordinates of the cluster and the extrapolated point, the module checks that this
 2634 is not greater than a certain cut. If the cluster is in an end cap, I propagate the track
 2635 up to the x position of the cluster. Then, the algorithm computes the angle in the (z, y)
 2636 plane between the centre of curvature and the cluster, α , and the centre of curvature
 2637 and the propagated point, α' . A cut is applied to the quantity $(\alpha - \alpha')R$.
 2638 If the cluster contains more than a certain number N of hits, I apply an extra cut to
 2639 the dot product of the direction of the track at the propagated x value and the cluster

Chapter 8. Particle ID in GArSoft




Figure 8.13: Distributions of energy deposits in the ECal for a muon (left) and a charged pion (right) with similar momentum. The energy is projected onto the plane perpendicular to the principal component of the hit clusters, and the positions are relative to the center of the interaction.

2640 direction.

2641 The code makes sure to only associate one end of the track (if any) to a cluster.
 2642 However, it can associate more than one track to the same cluster. This makes sense,
 2643 as different particles can contribute to the same cluster in the ECal, but it makes it
 2644 difficult to quantify the relative contributions of the tracks to a certain cluster.

2645 As a way of comparing the performance of this algorithm, a new, simpler association
 2646 module was written. The goal was to have a simple and robust algorithm, which depends
 2647 on as few parameters as possible and that can produce a one-to-one matching between
 2648 tracks and ECal clusters.

2649 For each reconstructed track, the new algorithms applies the same procedure to the
 2650 forward and the backward fits irrespective of their end point positions. It first gets the
 2651 Kalman fit parameters at the corresponding end point together with the X position, x_0 ,
 2652 (y_0 , z_0 , $1/R$, ϕ_0 , $\tan\lambda$).

2653 For each ECal cluster, I compute the radial distance to the centre of the TPC and
 2654 find the ϕ value in the range $[\phi_0, \phi_0 + \text{sign}(R)\phi_{max}]$ that makes the propagated helix
 2655 intersect with the circle defined with such radius. The (x, y, z) position of the helix for
 2656 the ϕ value found (if any) is then computed. In case there are two intersections, I keep

8.2. Muon and pion separation in the ECal and MuID




Figure 8.14: Left panel: comparison between the precision (blue), sensitivity (yellow) and F_1 score (red) obtained for the default (horizontal lines) and new algorithms, both with the χ^2 -based direction estimator (squares) and cheating the directions (circles), for different values of the χ^2 cut. Right panel: comparison of the performance of the new algorithm when applying the cluster t_0 correction (squares) and when (circles).

2657 the one that minimises the distance between (y, z) and (y_c, z_c) .

2658 I then calculate χ^2 value based on the Euclidean distance between the propagated
2659 point and the cluster:

$$\chi^2/ndf = \frac{\sum_{n=0}^2 (x^{(n)} - x_c^{(n)})^2}{3}. \quad (8.9)$$

2660 If there was no intersection I store a -1 instead. In the end, for each reconstructed track
2661 in the event one ends up with two collections of χ^2 values, one for each ECal cluster
2662 and fit directions.

2663 The current code only supports having ECal clusters associated to one end of each
2664 track. We have two options to decide what track end to keep. The first one tries to
2665 cheat the selection, looking at the distance between the two track ends and the true
2666 start position of the associated MC particle. The second one keeps the track end with
2667 more χ^2 entries below the cut.

2668 This feature of only considering one track end limits the algorithm, making it not
2669 suitable for reconstructing events with particles originating outside the TPC. However,
2670 as for the moment the main concern of the group is the study of neutrino interactions
2671 off the gaseous argon, this is an acceptable assumption.

Chapter 8. Particle ID in GArSoft

2672 In order to associate a cluster to a track, I take all clusters with a χ^2 value in the
2673 range $[0, \chi_{cut}^2]$. If a cluster has been assigned to more than one track we leave it with
2674 the one with the lowest χ^2 .

2675 This default behaviour of the algorithm can be modified to associate more than one
2676 track to each cluster. Not only that, but the χ^2 values can be used to assign relative
2677 weights to the different contributions.

2678 To evaluate the performance of the association method, I use a binary classification
2679 approach. In this case, I check the leading MC Track IDs associated to the reconstructed
2680 tracks and ECal clusters. I count an association as true positive (TP) if both Track
2681 IDs coincide. An association is considered false positive (FP) when the Track IDs are
2682 different. If a cluster has not been associated to any track but it shares the Track ID
2683 with a reconstructed track it is counted as a false negative (FN).

2684 For the testing, I used a sample of 10000 FHC neutrino events inside the HPgTPC.
2685 Fig. 8.14 (left panel) shows the precision (blue line), sensitivity (yellow line), and F_1
2686 score (red line) I obtained for different values of χ_{cut}^2 . For comparison, the same metrics
2687 computed for the default algorithm with the current configuration are also shown (dashed
2688 lines). In the case of the new algorithm, I used both the χ^2 -based method to estimate
2689 the track direction described earlier (square markers) and the cheated direction from the
2690 Geant-level information (circle markers). For either of these we achieve similar values of
2691 the precision compared to the old code, while having a considerably higher sensitivity.
2692 It can be seen that cheating the direction of the tracks only makes a difference at high
2693 χ_{cut}^2 , past the optimal value of the cut around the F_1 score maximum. Therefore, I set
2694 the χ^2 method as the default.

2695 One of the possible weak points of this approach is that it relies on the position along
2696 the drift direction to make the decisions. Within the current ND-GAr design implemented
2697 in GArSoft, the timing information is provided by the ECal. That effectively means
2698 that prior to make the track-ECal associations the reconstructed x positions of the track

8.2. Muon and pion separation in the ECal and MuID




Figure 8.15: Schematics of a possible option to deal with track-ECal associations in non-zero t_0 neutrino interaction events, trying to correct for the drift direction uncertainty in a cluster-by-cluster basis using the cluster time, $t_{cluster}$.

2699 trajectories differ from the simulated ones by an amount:

$$x_{reco}^{(n)} - x_{sim}^{(n)} = v_{drift} t_0, \quad (8.10)$$

2700 where v_{drift} is the mean drift velocity in our medium and the initial time is in the range
 2701 $t_0 \in [0, t_{spill}]$ where t_{spill} is the spill length. For a 10 μs spill this translates into a
 2702 maximum 30 cm uncertainty on the drift direction position.

2703 The current default in GArSoft sets $t_0 = 0$, but the functionality to randomly sample
 2704 this within the spill time is in place. Therefore, we need to understand what is the impact
 2705 of a non-zero t_0 on the associations algorithm and foresee possible ways of minimising a
 2706 loss in performance.

2707 Fig. 8.15 represents a possible option to tackle the association problem when having
 2708 events with a non-zero initial time t_0 . The black and white circles represent the original
 2709 points, whereas the squares indicate the corrected positions. The end points of the track
 2710 and the propagated points up to the cluster radius are indicated using filled and unfilled
 2711 markers respectively. The red square represents the position of the cluster.

2712 Here I try to correct for the drift coordinate position using the time associated to the
 2713 cluster. Assuming that the drift time is much larger than the propagation time, $t_{cluster}$
 2714 could be used as a good estimation of the t_0 . An alternative can be using the earliest
 2715 time associated to a hit in said cluster. Doing this for each cluster before computing
 2716 the χ^2 value could be used as an alternative to knowing the specific value of the t_0 , as

Chapter 8. Particle ID in GArSoft

when the association is correct this will provide the right correction but its impact is small enough to not change the position significantly in the case the cluster does not correspond to a given track.

I tested the effect of this correction again using a sample of 10000 FHC neutrino events. Fig. 8.14 (right panel) shows the precision (blue line), sensitivity (yellow line), and F_1 score (red line) for the case the cluster t_0 correction is applied (square markers) and for the no correction case (circle markers), as a function of $\chi^2_{c}ut$. In this case, the differences are particularly notorious at low values of the cut. It makes sense, as the t_0 effect becomes subdominant when the distance we consider grows large. Overall, the correction increases the sensitivity while keeping the precision almost unchanged. As a result, I apply the t_0 correction to the generated samples as the default.

8.2.2 Feature selection and importance

I am using samples with 100k FHC neutrino interactions, no truth cuts whatsoever. From the Geant4 I am only keeping the true momentum and PDG. From the track reco I keep the momentum (end momentum, both forward and backward fits).

For the ECAL, I compute a series of variables per each reco track. As there can be more than one ECAL cluster associated to each track what I do is collect all associated clusters and compute my variables from the complete collection of hits associated to each track. I can roughly divide the variables in three types: energy-related, geometry-related and statistical.

• Energy-related ECAL

- ECAL total energy (ClusterTotalEnergy): sum of the energy of all the ECAL hits.
- Hit mean energy (HitMeanEnergy): mean of the hits energy distribution.
- Hit std energy (HitStdEnergy): standard deviation of the hits energy distribution.
- Hit max energy (HitMaxEnergy): maximum of the hits energy distribution.

8.2. Muon and pion separation in the ECal and MuID

2743 **Geometry-related ECAL**

- 2744 • Mean distance hit-to-cluster (DistHitClusterMean): mean of the distance distribution
- 2745 between the hits and the corresponding cluster's main axis.
- 2746 • RMS distance hit-to-cluster (DistHitClusterRMS): root mean square of the distance
- 2747 distribution between the hits and the corresponding cluster's main axis.
- 2748 • Maximum distance hit-to-centre (DistHitCenterMax): maximum of the distance
- 2749 distribution between the hits and the centre of the TPC.
- 2750 • Time-of-Flight velocity (TOFVelocity): slope obtained when fitting a straight line
- 2751 to the hit time versus hit distance to the centre (i.e. $d = v \times t$).

2752 **Energy and Geometry ECAL**

- 2753 • Radius 90% energy (Radius90E): distance in the hit-to-cluster distribution for
- 2754 which 90% of the total energy is contained in the hits that are closer to the axis
- 2755 (i.e. radius that contains 90% of the energy).

2756 **Statistical ECAL**

- 2757 • Number of hits (NHits): total number of hits associated to the track.
- 2758 • Number of layers with hits (NLayers): not really a count of all layers with hits but
- 2759 the difference between the last and the first layer with hits.

2760 For the MuID, now that the bug affecting the position of the reco hits has been
2761 solved (07/07/23) I also included some related quantities in the analysis ntuples. They
2762 also fall in the same classification I use for the ECAL variables and are also defined per
2763 reco track as there is also the possibility of having more than one MuID cluster for each
2764 track.

2765 **Energy-related MuID**

- 2766 • MuID total energy (ClusterMuIDTotalEnergy): sum of the energy of all the MuID
- 2767 hits.

Chapter 8. Particle ID in GArSoft

- 2768 • Hit MuID mean energy (HitMuIDMeanEnergy): mean of the MuID hits energy
2769 distribution.

- 2770 • Hit MuID std energy (HitMuIDStdEnergy): standard deviation of the MuID hits
2771 energy distribution.

- 2772 • Hit MuID max energy (HitMuIDMaxEnergy): maximum of the MuID hits energy
2773 distribution.

2774 Geometry-related MuID

- 2775 • Max distance MuID hits (DistHitMuIDMax): maximum distance between pairs of
2776 MuID hits (not sure this is a good variable, distribution looks nuts).

- 2777 • Maximum distance MuID hit-to-centre (DistHitCenterMuIDMax): maximum of
2778 the distance distribution between the MuID hits and the centre of the TPC.

2779 Statistical MuID

- 2780 • Number of hits (NHitsMuID): total number of MuID hits associated to the track.

- 2781 • Number of layers with hits (NLayersMuID): not really a count of all layers with
2782 MuID hits but the difference between the last and the first layer with MuIDhits.

8.3. ECal time-of-flight

2783 **8.2.3 Hyperparameter optimisation**

2784 **8.2.4 Probability calibration**

2785 **8.2.5 Performance**

2786 **8.3 ECal time-of-flight**

2787 **8.3.1 Arrival time estimations**

2788 **8.3.2 Proton and pion separation**

2789 **8.4 Charged pion decay in flight**

2790 **8.4.1 Track breakpoints**

2791 **8.5 Neutral particle identification**

2792 **8.5.1 ECal clustering**

2793 Another important reconstruction item is the clustering algorithm of ECal hits in
2794 GArSoft. The default module features a NN algorithm that treats all hits in the same
2795 way, independently of the layer each hit comes from. However, the current ECal design
2796 of ND-GAr has two very different types of scintillator layers. The inner layers are made
2797 out of tiles, which provide excellent angular and timing resolutions. On the other hand,
2798 the outer layers are cross scintillator strips. That way, an algorithm that treats hits
2799 from both kinds of layers differently may be able to improve the current performance.

2800 Inspired by the reconstruction of T2K’s ND280 downstream ECal [?], the idea was
2801 to put together a clustering module that first builds clusters for the different ECal views
2802 (tiles, strips segmented in the X direction and strips segmented in Y direction), and
2803 then tries to match them together to form the final clusters.

2804 Working on a module-by-module basis, the algorithm first separates the hits depending
2805 on the layer type they come from. Then, it performs a NN clustering for the 3 sets of
2806 hits separately. For the tile hits it clusters together all the hits which are in nearest-

Chapter 8. Particle ID in GArSoft

2807 neighbouring tiles and nearest-neighbouring layers, for strip hits it looks at nearest-
2808 neighbouring strips and next-to-nearest-neighbouring layers (as the layers with strips
2809 along the two directions are alternated). For strip clusters an additional cut in the
2810 direction along the strip length is needed.

2811 After this first clustering I then apply a recursive re-clustering for each collection
2812 of strip clusters based on a PCA method. In each case, we loop over the clusters with
2813 $N_{hits} \geq 2$, computing the centre of mass and three principal components. Propagating
2814 these axes up to the layers of the rest of the clusters, we check if the propagated point
2815 and the centre of mass of the second cluster are within next-to-nearest-neighbouring
2816 strips. An additional cut in the direction along the strip length is also needed. Moreover,
2817 I require that the two closest hits across the two clusters are at most in next-to-nearest-
2818 neighbouring strips. I merge the clusters if these three conditions are satisfied. The
2819 re-clustering is repeated until no more cluster pairs pass the cuts.

2820 The clusters in each strip view are combined if their centres of mass are close enough
2821 and they point in the same direction. An alternative approach for the strip cluster
2822 merging could be to compute the overlap between the ellipsoids defined by the principal
2823 axes of the clusters, and then merge the pair if the overlap exceeds some threshold.
2824 Further study is needed to understand if this change would have an impact in the overall
2825 clustering performance.

2826 To merge the tile clusters to the combined strip clusters I propagate the principal
2827 axis of the strip cluster towards the inner layers, up to the centre of mass layer of the
2828 tile cluster. I merge the clusters if the distance between the propagated point and the
2829 centre of mass is below a certain cut.

2830 The last step is to check if clusters in neighbouring modules should be merged
2831 together, both across two barrel modules, across end cap modules and between barrel
2832 end cap modules. I check the distance between the two closest hits in the pair of clusters
2833 and merge them if it passes this and an additional direction cut.

2834 Fig. ?? presents an example of the clustering steps relevant for strip layer hits, from
2835 the input hits (top left panel) to the NN clustering (top right panel) and re-clustering

8.5. Neutral particle identification




Figure 8.16: Mean values of the F_1 -score marginal distributions for the different free parameters of the new clustering algorithm, with the error bars representing one standard deviation around the mean. The F_1 -score values were computed for the 6561 possible parameter configurations using 1000 ν_μ CC interaction events.

(bottom left panel) for each strip view and the final merging strip clusters (bottom right panel). It shows the hits from a single ECal barrel module in a ν_μ CC interaction event with a neutral pion and a proton in the final state. The two clusters on the left correspond to the photon pair from the π^0 decay and the one on the upper right corner is associated to the proton.

Table 8.2: Summary of parameters and sampled values used in the optimisation of the clustering algorithm.

Name	Units	Sampled values	Description
PrimMinorCut	strips	2, 3, 4	Distance along strip length in NN clustering
RecMinorCut	strips	3, 5, 8	Distance between propagated point and CM along strip length in re-clustering
RecPerpCut	strips	2, 3, 4	Closest hit pair distance in re-clustering
StripDistCut	strips	3, 5, 8	Distance between CMs in strip cluster merging
StripDirCut	cos	0.5, 0.7, 0.9	Main axes direction cut in strip cluster merging
PointDistCut	tiles	3, 5, 8	Distance between propagated point and CM in strip-tile matching
MergePerpCut	cm	10, 20, 30	Closest hit pair distance in module merging
MergeDirCut	cos	0.5, 0.7, 0.9	Main axes direction cut in module merging

Chapter 8. Particle ID in GArSoft

2841 This algorithm has a total number of eight free parameters that need to be optimised.
2842 I used a sample of 1000 ν_μ CC interactions in order to obtain the optimal configuration of
2843 clustering parameters. This sample was generated up to the default ECal hit clustering
2844 level, so then I could run the new clustering algorithm each time with a different
2845 configuration of parameters. As the number of parameters is relatively large, I only
2846 performed a coarse-grained scan of the parameter space. Sampling each of the eight
2847 parameters at three different points each I obtain 6561 different configurations. These
2848 parameters, together with the used values, are summarised in Tab. 8.2.

2849 In order to measure the performance of the clustering, I use a binary classification
2850 approach. For each formed cluster, I identify the Geant4 Track ID of the matching MC
2851 particle and the energy fraction of each hit. Then, I assign to each cluster the Track ID
2852 with the highest total energy fraction. For each of the different Track IDs associated to
2853 the clusters, I select the cluster with the highest energy (only from the hits with the
2854 same Track ID). I identify such a cluster as the main cluster for that Track ID. I count
2855 as true positives (TPs) the hits with the correct Track ID in each main cluster. False
2856 positives (FPs) are the hits with the incorrect Track ID for the cluster they are in, not
2857 only main clusters. The false negatives (FNs) are the hits with the correct Track ID in
2858 clusters other than the main.

2859 Fig. 8.16 shows the computed F_1 -score values for the different cuts. In each case, the
2860 central value represents the mean of the F_1 -score distribution for the specified value of
2861 the corresponding variable and the vertical error bar represents one standard deviation
2862 around the mean. Also shown are the Pearson correlation coefficients of these central
2863 values. We can see that five of the variables have a sizeable effect on the F_1 -score, with
2864 an absolute difference between the last and first values as big as 4%.

2865 The working configuration is obtained as follows. I first select all configurations
2866 with purity $\geq 90\%$. Among those, I choose the combinations that yield the maximum
2867 F_1 -score. If more than one configuration remains I select the one with the highest
2868 sensitivity. Doing so, I end up with a parameter configuration with an efficiency of 88%
2869 and a 90% purity. Compared with the default algorithm, which gives an efficiency of

8.5. Neutral particle identification




Figure 8.17: Left panel: distributions of the number of ECal clusters per photon from π^0 decays for the standard (red) and new (blue) clustering algorithms. Right panel: reconstructed invariant mass distributions for photon pairs from single π^0 events using the standard (red) and new (blue) ECal clustering algorithms.

2870 76% and a purity of 91% for the same sample, I have managed to improve the efficiency
2871 by a factor of 1.16.

2872 **8.5.2 π^0 reconstruction**

2873 One of the potential applications of the new ECal hit clustering is the reconstruction of
2874 neutral particles, in particular pions. Neutral pions decay promptly after being produced,
2875 through the $\pi^0 \rightarrow \gamma\gamma$ channel $(98.823 \pm 0.034)\%$ of the time. The photon pair does
2876 not leave any traces in the HPgTPC (unless one or both of them converts into an
2877 electron-positron pair), but each of them will produce an electromagnetic shower in
2878 the ECal.

2879 To test the potential impact of the new algorithm in π^0 reconstruction, I generated
2880 a MC sample of single, isotropic neutral pions inside the HPgTPC. All pions were
2881 generated with a momentum $p = 500$ MeV and their initial positions were uniformly
2882 sampled inside a $2 \times 2 \times 2$ m box aligned with the centre of the TPC. I ran both the
2883 default and the new clustering algorithms, using for the latter the optimised configuration
2884 discussed above.

2885 The first thing to notice is that the number of clusters produced per photon has
2886 decreased. Fig. 8.17 (left panel) shows these distributions for the default (red) and new

Chapter 8. Particle ID in GArSoft

(blue) algorithms. Using a simple Gaussian fit, we see that the mean number of ECal clusters per photon went from 1.82 ± 0.01 to 1.09 ± 0.03 . This effectively means that with the new algorithm the ECal activity of one true particle is typically reconstructed as a single object. From the reconstruction point of view this can be an advantage. As now most of the photon energy ends up in a single ECal cluster, I can simply use cluster pairs to identify the π^0 decay.

In general, one calculates the invariant mass of the photon pair as:

$$m_{\gamma\gamma} = \sqrt{2E_1E_2(1 - \cos \theta)}, \quad (8.11)$$

where E_i are the energies of the photons and θ the opening angle between them. In this case I can use the energies deposited in the ECal and their incident directions. This quantity is computed for all possible pairs of clusters, using their position together with the true decay point. In a more realistic scenario, e.g. ν_μ CC interaction, one could use the position of the reconstructed primary vertex instead. I also tried to use the principal direction of the clusters, but that approach gave considerably worse results. For each event I only keep the pair with an invariant mass closer to the true π^0 mass value.

Fig. 8.17 (right panel) shows the invariant mass distributions for the photon pairs we get using the default (red) and the new (blue) ECal clustering algorithms. For the fit I used a modified version of the Crystal Ball function [?], obtained by taking the limit where the parameter controlling the power-law tail goes to infinity:

$$f(x; N, \mu, \sigma, \alpha) = N \cdot \begin{cases} e^{\frac{\alpha(2x-2\mu+\alpha\sigma)}{2\sigma}}; & x \leq \mu - \alpha\sigma, \\ e^{-\frac{(x-\mu)^2}{2\sigma^2}}; & x > \mu - \alpha\sigma. \end{cases} \quad (8.12)$$

Comparing the fitted mean and standard deviation values for the Gaussian cores, we see that the distribution for the new algorithm is a 67% narrower and also peaks much closer to the true m_{π^0} value, going from 101.3 ± 0.4 MeV to 130.8 ± 0.6 MeV.

₂₉₀₈ Chapter 9

₂₉₀₉ Conclusions

₂₉₁₀ Appendix A

₂₉₁₁ An appendix

2912 Bibliography

- 2913 [1] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
2914 *Detector Technical Design Report, Volume I Introduction to DUNE*, *JINST* **15**
2915 (2020) T08008 [2002.02967]. 15, 41, 42, 43, 46, 47, 48, 54, 55, 56, 63, 64
- 2916 [2] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
2917 *Detector Technical Design Report, Volume IV: Far Detector Single-phase*
2918 *Technology*, *JINST* **15** (2020) T08010 [2002.03010]. 15, 51
- 2919 [3] J.N. Bahcall, A.M. Serenelli and S. Basu, *New solar opacities, abundances,*
2920 *helioseismology, and neutrino fluxes*, *Astrophys. J. Lett.* **621** (2005) L85
2921 [[astro-ph/0412440](#)]. 21, 111, 113
- 2922 [4] R. Garani and S. Palomares-Ruiz, *Dark matter in the Sun: scattering off electrons*
2923 *vs nucleons*, *JCAP* **05** (2017) 007 [[1702.02768](#)]. 21, 109, 110, 112, 113, 114, 137
- 2924 [5] M. Honda, M. Sajjad Athar, T. Kajita, K. Kasahara and S. Midorikawa,
2925 *Atmospheric neutrino flux calculation using the NRLMSISE-00 atmospheric model*,
2926 *Phys. Rev. D* **92** (2015) 023004 [[1502.03916](#)]. 22, 117, 118
- 2927 [6] M. Colom i Bernadich and C. Pérez de los Heros, *Limits on Kaluza-Klein dark*
2928 *matter annihilation in the Sun from recent IceCube results*, *Eur. Phys. J. C*, **80** 2
2929 (2020) 129 **80** (2019) [[1912.04585](#)]. 22, 122
- 2930 [7] ANTARES collaboration, *Search for Dark Matter in the Sun with the ANTARES*
2931 *Neutrino Telescope in the CMSSM and mUED frameworks*, *Nucl. Instrum. Meth. A*
2932 **725** (2013) 76 [[1204.5290](#)]. 22, 122

BIBLIOGRAPHY

- 2933 [8] N. Deutschmann, T. Flacke and J.S. Kim, *Current LHC Constraints on Minimal*
2934 *Universal Extra Dimensions*, *Phys. Lett. B* **771** (2017) 515 [[1702.00410](#)]. 22, 122,
2935 123
- 2936 [9] ICECUBE collaboration, *Search for GeV-scale dark matter annihilation in the Sun*
2937 *with IceCube DeepCore*, *Phys. Rev. D* **105** (2022) 062004 [[2111.09970](#)]. 24, 134,
2938 135
- 2939 [10] C.-S. Chen, F.-F. Lee, G.-L. Lin and Y.-H. Lin, *Probing Dark Matter*
2940 *Self-Interaction in the Sun with IceCube-PINGU*, *JCAP* **10** (2014) 049 [[1408.5471](#)].
2941 24, 134, 135
- 2942 [11] N.F. Bell, M.J. Dolan and S. Robles, *Searching for dark matter in the Sun using*
2943 *Hyper-Kamiokande*, *JCAP* **11** (2021) 004 [[2107.04216](#)]. 24, 134, 135
- 2944 [12] E. Behnke et al., *Final Results of the PICASSO Dark Matter Search Experiment*,
2945 *Astropart. Phys.* **90** (2017) 85 [[1611.01499](#)]. 24, 134, 135
- 2946 [13] PICO collaboration, *Dark Matter Search Results from the Complete Exposure of*
2947 *the PICO-60 C₃F₈ Bubble Chamber*, *Phys. Rev. D* **100** (2019) 022001
2948 [[1902.04031](#)]. 24, 134, 135
- 2949 [14] DARKSIDE collaboration, *Constraints on Sub-GeV Dark-Matter–Electron*
2950 *Scattering from the DarkSide-50 Experiment*, *Phys. Rev. Lett.* **121** (2018) 111303
2951 [[1802.06998](#)]. 24, 138, 139
- 2952 [15] XENON collaboration, *Light Dark Matter Search with Ionization Signals in*
2953 *XENON1T*, *Phys. Rev. Lett.* **123** (2019) 251801 [[1907.11485](#)]. 24, 138, 139
- 2954 [16] P.F. de Salas, D.V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C.A. Ternes
2955 et al., *2020 global reassessment of the neutrino oscillation picture*, *JHEP* **02** (2021)
2956 071 [[2006.11237](#)]. 29, 38, 43
- 2957 [17] W. Pauli, *Dear radioactive ladies and gentlemen*, *Phys. Today* **31N9** (1978) 27. 33

BIBLIOGRAPHY

- 2958 [18] F. Reines and C.L. Cowan, *Detection of the free neutrino*, *Phys. Rev.* **92** (1953) 830. 33
- 2959
- 2960 [19] ALEPH, DELPHI, L3, OPAL, SLD, LEP ELECTROWEAK WORKING GROUP, SLD ELECTROWEAK GROUP, SLD HEAVY FLAVOUR GROUP collaboration, *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257 [[hep-ex/0509008](#)]. 34
- 2961
- 2962
- 2963
- 2964 [20] SUPER-KAMIOKANDE collaboration, *Evidence for oscillation of atmospheric neutrinos*, *Phys. Rev. Lett.* **81** (1998) 1562 [[hep-ex/9807003](#)]. 34
- 2965
- 2966 [21] B. Pontecorvo, *Inverse beta processes and nonconservation of lepton charge*, *Zh. Eksp. Teor. Fiz.* **34** (1957) 247. 34
- 2967
- 2968 [22] Z. Maki, M. Nakagawa and S. Sakata, *Remarks on the unified model of elementary particles*, *Prog. Theor. Phys.* **28** (1962) 870. 34
- 2969
- 2970 [23] L. Wolfenstein, *Neutrino Oscillations in Matter*, *Phys. Rev. D* **17** (1978) 2369. 36
- 2971
- 2972 [24] B.T. Cleveland, T. Daily, R. Davis, Jr., J.R. Distel, K. Lande, C.K. Lee et al., *Measurement of the solar electron neutrino flux with the Homestake chlorine detector*, *Astrophys. J.* **496** (1998) 505. 37
- 2973
- 2974 [25] F. Kaether, W. Hampel, G. Heusser, J. Kiko and T. Kirsten, *Reanalysis of the GALLEX solar neutrino flux and source experiments*, *Phys. Lett. B* **685** (2010) 47 [[1001.2731](#)]. 37
- 2975
- 2976
- 2977 [26] SAGE collaboration, *Measurement of the solar neutrino capture rate with gallium metal. III: Results for the 2002–2007 data-taking period*, *Phys. Rev. C* **80** (2009) 015807 [[0901.2200](#)]. 37
- 2978
- 2979
- 2980 [27] G. Bellini et al., *Precision measurement of the ^{7}Be solar neutrino interaction rate in Borexino*, *Phys. Rev. Lett.* **107** (2011) 141302 [[1104.1816](#)]. 37
- 2981

BIBLIOGRAPHY

- 2982 [28] SUPER-KAMIOKANDE collaboration, *Solar neutrino measurements in
super-Kamiokande-I*, *Phys. Rev. D* **73** (2006) 112001 [[hep-ex/0508053](#)]. 37
- 2984 [29] SNO collaboration, *Combined Analysis of all Three Phases of Solar Neutrino Data
from the Sudbury Neutrino Observatory*, *Phys. Rev. C* **88** (2013) 025501
[[1109.0763](#)]. 37
- 2987 [30] SUPER-KAMIOKANDE collaboration, *Atmospheric neutrino oscillation analysis with
external constraints in Super-Kamiokande I-IV*, *Phys. Rev. D* **97** (2018) 072001
[[1710.09126](#)]. 37
- 2990 [31] ICECUBE collaboration, *Measurement of Atmospheric Neutrino Oscillations at
6–56 GeV with IceCube DeepCore*, *Phys. Rev. Lett.* **120** (2018) 071801
[[1707.07081](#)]. 37
- 2993 [32] KAMLAND collaboration, *Reactor On-Off Antineutrino Measurement with
KamLAND*, *Phys. Rev. D* **88** (2013) 033001 [[1303.4667](#)]. 37
- 2995 [33] RENO collaboration, *Measurement of Reactor Antineutrino Oscillation Amplitude
and Frequency at RENO*, *Phys. Rev. Lett.* **121** (2018) 201801 [[1806.00248](#)]. 37
- 2997 [34] DAYA BAY collaboration, *Measurement of the Electron Antineutrino Oscillation
with 1958 Days of Operation at Daya Bay*, *Phys. Rev. Lett.* **121** (2018) 241805
[[1809.02261](#)]. 37
- 3000 [35] K.J. Kelly, P.A.N. Machado, S.J. Parke, Y.F. Perez-Gonzalez and R.Z. Funchal,
Neutrino mass ordering in light of recent data, *Phys. Rev. D* **103** (2021) 013004. 38
- 3002 [36] P. Dunne, “Latest Neutrino Oscillation Results from T2K.” Jul, 2020. 38
- 3003 [37] MINOS collaboration, *Combined analysis of ν_μ disappearance and $\nu_\mu \rightarrow \nu_e$
appearance in MINOS using accelerator and atmospheric neutrinos*, *Phys. Rev. Lett.*
112 (2014) 191801 [[1403.0867](#)]. 38

BIBLIOGRAPHY

- 3006 [38] K2K collaboration, *Measurement of Neutrino Oscillation by the K2K Experiment*,
3007 *Phys. Rev. D* **74** (2006) 072003 [[hep-ex/0606032](#)]. 38
- 3008 [39] DUNE collaboration, *Long-baseline neutrino oscillation physics potential of the*
3009 *DUNE experiment*, *Eur. Phys. J. C* **80** (2020) 978 [[2006.16043](#)]. 38
- 3010 [40] HYPER-KAMIOKANDE collaboration, *Physics potential of Hyper-Kamiokande for*
3011 *neutrino oscillation measurements*, *PoS NuFact2019* (2019) 040. 38
- 3012 [41] DUNE collaboration, *Snowmass Neutrino Frontier: DUNE Physics Summary*,
3013 [2203.06100](#). 43, 44
- 3014 [42] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE), Far*
3015 *Detector Technical Design Report, Volume II: DUNE Physics*, [2002.03005](#). 43, 45,
3016 59, 61, 117, 126, 130
- 3017 [43] SUPER-KAMIOKANDE collaboration, *Search for Proton Decay via $p \rightarrow e^+ \pi_0$ and*
3018 *$p \rightarrow \mu^+ \pi_0$ in a Large Water Cherenkov Detector*, *Phys. Rev. Lett.* **102** (2009)
3019 [141801](#) [[0903.0676](#)]. 44
- 3020 [44] S. Raby, *Grand Unified Theories*, in *2nd World Summit: Physics Beyond the*
3021 *Standard Model*, 8, 2006 [[hep-ph/0608183](#)]. 44
- 3022 [45] KAMIOKANDE-II collaboration, *Observation of a Neutrino Burst from the*
3023 *Supernova SN 1987a*, *Phys. Rev. Lett.* **58** (1987) 1490. 45
- 3024 [46] R.M. Bionta et al., *Observation of a Neutrino Burst in Coincidence with Supernova*
3025 *SN 1987a in the Large Magellanic Cloud*, *Phys. Rev. Lett.* **58** (1987) 1494. 45
- 3026 [47] DUNE collaboration, *The DUNE Far Detector Vertical Drift Technology*,
3027 *Technical Design Report*, [2312.03130](#). 49, 50
- 3028 [48] DUNE collaboration, *Deep Underground Neutrino Experiment (DUNE) Near*
3029 *Detector Conceptual Design Report, Instruments* **5** (2021) 31 [[2103.13910](#)]. 53, 54,
3030 57, 61, 141

BIBLIOGRAPHY

- 3031 [49] J. Strait, E. McCluskey, T. Lundin, J. Willhite, T. Hamernik, V. Papadimitriou
3032 et al., *Long-baseline neutrino facility (lbnf) and deep underground neutrino*
3033 *experiment (dune) conceptual design report volume 3: Long-baseline neutrino*
3034 *facility for dune june 24, 2015*, 1601.05823. 58
- 3035 [50] DUNE DAQ, “dtp-firmware.”
3036 <https://gitlab.cern.ch/dune-daq/readout/dtp-firmware>, 2020. 73
- 3037 [51] DUNE DAQ, “dtp-simulation.”
3038 <https://gitlab.cern.ch/dune-daq/readout/dtp-simulation>, 2020. 76
- 3039 [52] DUNE DAQ, “dtpemulator.”
3040 https://github.com/DUNE-DAQ/dtpemulator/tree/fmlopez/filter_ana, 2022.
3041 76
- 3042 [53] J. McClellan and T. Parks, *A personal history of the Parks-McClellan algorithm*,
3043 *IEEE Signal Processing Magazine* **22** (2005) 82. 77
- 3044 [54] G. Turin, *An introduction to matched filters*, *IEEE Transactions on Information*
3045 *Theory* **6** (1960) 311. 80
- 3046 [55] J.W. Goodman, *Statistical Optics*, Wiley (1985). 82
- 3047 [56] B. Dwork, *Detection of a pulse superimposed on fluctuation noise*, *Proceedings of*
3048 *the IRE* **38** (1950) 771. 83
- 3049 [57] L. Wainstein and V. Zubakov, *Extraction of Signals from Noise*, Prentice-Hall
3050 (1962). 83
- 3051 [58] E.D. Church, *Larsoft: A software package for liquid argon time projection drift*
3052 *chambers*, 1311.6774. 86, 87
- 3053 [59] S.V. Stehman, *Selecting and interpreting measures of thematic classification*
3054 *accuracy*, *Remote Sensing of Environment* **62** (1997) 77. 100

BIBLIOGRAPHY

- 3055 [60] A.A. Taha and A. Hanbury, *Metrics for evaluating 3d medical image segmentation: analysis, selection, and tool*, *BMC Medical Imaging* **15** (2015) . 101
- 3056
- 3057 [61] J. Silk, K.A. Olive and M. Srednicki, *The Photino, the Sun and High-Energy Neutrinos*, *Phys. Rev. Lett.* **55** (1985) 257. 107
- 3058
- 3059 [62] M. Srednicki, K.A. Olive and J. Silk, *High-Energy Neutrinos from the Sun and Cold Dark Matter*, *Nucl. Phys. B* **279** (1987) 804. 107
- 3060
- 3061 [63] J.S. Hagelin, K.W. Ng and K.A. Olive, *A High-energy Neutrino Signature From Supersymmetric Relics*, *Phys. Lett. B* **180** (1986) 375. 107
- 3062
- 3063 [64] T.K. Gaisser, G. Steigman and S. Tilav, *Limits on Cold Dark Matter Candidates from Deep Underground Detectors*, *Phys. Rev. D* **34** (1986) 2206. 107
- 3064
- 3065 [65] N. Bernal, J. Martín-Albo and S. Palomares-Ruiz, *A novel way of constraining WIMPs annihilations in the Sun: MeV neutrinos*, *JCAP* **08** (2013) 011
- 3066
- 3067 [1208.0834]. 107, 108, 109, 115
- 3068 [66] C. Rott, J. Siegal-Gaskins and J.F. Beacom, *New Sensitivity to Solar WIMP Annihilation using Low-Energy Neutrinos*, *Phys. Rev. D* **88** (2013) 055005
- 3069
- 3070 [1208.0827]. 107, 114
- 3071 [67] C. Rott, S. In, J. Kumar and D. Yaylali, *Dark Matter Searches for Monoenergetic Neutrinos Arising from Stopped Meson Decay in the Sun*, *JCAP* **11** (2015) 039
- 3072
- 3073 [1510.00170]. 107
- 3074 [68] DUNE collaboration, *Searching for solar KDAR with DUNE*, *JCAP* **10** (2021) 065
- 3075 [2107.09109]. 107
- 3076 [69] G. Busoni, A. De Simone and W.-C. Huang, *On the Minimum Dark Matter Mass Testable by Neutrinos from the Sun*, *JCAP* **07** (2013) 010 [1305.1817]. 108
- 3077
- 3078 [70] V.A. Bednyakov and F. Simkovic, *Nuclear spin structure in dark matter search: The Zero momentum transfer limit*, *Phys. Part. Nucl.* **36** (2005) 131
- 3079
- 3080 [hep-ph/0406218]. 109

BIBLIOGRAPHY

- 3081 [71] A. Gould, *WIMP Distribution in and Evaporation From the Sun*, *Astrophys. J.* **321**
3082 (1987) 560. 110
- 3083 [72] J.N. Bahcall, M.H. Pinsonneault and S. Basu, *Solar models: Current epoch and*
3084 *time dependences, neutrinos, and helioseismological properties*, *Astrophys. J.* **555**
3085 (2001) 990 [[astro-ph/0010346](#)]. 112
- 3086 [73] T. Golan, J.T. Sobczyk and J. Zmuda, *NuWro: the Wroclaw Monte Carlo*
3087 *Generator of Neutrino Interactions*, *Nucl. Phys. B Proc. Suppl.* **229-232** (2012)
3088 499. 116
- 3089 [74] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for*
3090 *likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [[1007.1727](#)].
3091 117
- 3092 [75] T. Kaluza, *Zum Unitätsproblem der Physik, Sitzungsber. Preuss. Akad. Wiss.*
3093 *Berlin (Math. Phys.)* **1921** (1921) 966 [[1803.08616](#)]. 119
- 3094 [76] O. Klein, *Quantum Theory and Five-Dimensional Theory of Relativity. (In*
3095 *German and English)*, *Z. Phys.* **37** (1926) 895. 119
- 3096 [77] T. Appelquist, H.-C. Cheng and B.A. Dobrescu, *Bounds on universal extra*
3097 *dimensions*, *Phys. Rev. D* **64** (2001) 035002 [[hep-ph/0012100](#)]. 119
- 3098 [78] N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *The Hierarchy problem and new*
3099 *dimensions at a millimeter*, *Phys. Lett. B* **429** (1998) 263 [[hep-ph/9803315](#)]. 119
- 3100 [79] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*,
3101 *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)]. 119
- 3102 [80] G. Servant and T.M.P. Tait, *Is the lightest Kaluza-Klein particle a viable dark*
3103 *matter candidate?*, *Nucl. Phys. B* **650** (2003) 391 [[hep-ph/0206071](#)]. 119
- 3104 [81] H.-C. Cheng, K.T. Matchev and M. Schmaltz, *Radiative corrections to*
3105 *Kaluza-Klein masses*, *Phys. Rev. D* **66** (2002) 036005 [[hep-ph/0204342](#)]. 121

BIBLIOGRAPHY

- 3106 [82] M. Blennow, J. Edsjö and T. Ohlsson, *Neutrinos from WIMP annihilations using a*
3107 *full three-flavor Monte Carlo*, *JCAP* **01** (2008) 021 [[0709.3898](#)]. 121, 124
- 3108 [83] J. Edsjö, J. Elevant and C. Niblaeus, *WimpSim Neutrino Monte Carlo*. 121, 124
- 3109 [84] U. Haisch and A. Weiler, *Bound on minimal universal extra dimensions from*
3110 *anti-B → X(s)gamma*, *Phys. Rev. D* **76** (2007) 034014 [[hep-ph/0703064](#)]. 123
- 3111 [85] A. Freitas and U. Haisch, *Anti-B → X(s) gamma in two universal extra*
3112 *dimensions*, *Phys. Rev. D* **77** (2008) 093008 [[0801.4346](#)]. 123
- 3113 [86] C. Rott, D. Jeong, J. Kumar and D. Yaylali, *Neutrino Topology Reconstruction at*
3114 *DUNE and Applications to Searches for Dark Matter Annihilation in the Sun*,
3115 *JCAP* **07** (2019) 006 [[1903.04175](#)]. 126
- 3116 [87] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel et al.,
3117 *Scikit-learn: Machine learning in Python*, *Journal of Machine Learning Research*
3118 **12** (2011) 2825. 132
- 3119 [88] J. Kopp, V. Niro, T. Schwetz and J. Zupan, *DAMA/LIBRA and leptонically*
3120 *interacting Dark Matter*, *Phys. Rev. D* **80** (2009) 083502 [[0907.3159](#)]. 136
- 3121 [89] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2020**
3122 (2020) 083C01. 137
- 3123 [90] M. Beltran, D. Hooper, E.W. Kolb and Z.C. Krusberg, *Deducing the nature of dark*
3124 *matter from direct and indirect detection experiments in the absence of collider*
3125 *signatures of new physics*, *Phys. Rev. D* **80** (2009) 043509 [[0808.3384](#)]. 137
- 3126 [91] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron.*
3127 *Astrophys.* **641** (2020) A6 [[1807.06209](#)]. 138