

Assessment of summer 2018-2019 sea-ice forecasts for the Southern Ocean



Coordinating Seasonal Predictions of Sea Ice
in the Southern Ocean for 2017-2019

F. Massonnet, P. Reid, J. L. Lieser, C. M. Bitz, J. Fyfe, W. Hobbs

with contributions from

*Naval Research Lab, Nico Sun, NASA-GMAO, FIO-ESM, ECMWF, Lamont Sea Ice Group,
Alek Petty, Modified CanSIPS, Met Office, CMCC, Sandra Barreira, UCL*

May 7, 2019

Contact:

`francois.massonnet@uclouvain.be`

1. The Sea Ice Prediction Network South (SIPN South)

Being much thinner than Arctic sea ice and almost entirely seasonal, Antarctic sea ice has long been considered unpredictable beyond weather time scales. However, recent studies have unveiled several mechanisms of sea-ice predictability at seasonal time scales (Holland et al., 2017; Marchi et al., 2018; Holland et al., 2013). The study of sea-ice predictability does not only represent an academic exercise but has also many potential future applications. For example, knowledge of sea-ice presence from weeks to months in advance would be of great interest, since sea ice is one of the many hindrances that face vessels operating in the Antarctic coastal regions. In that context, advance notice of seasonal sea-ice conditions would help reduce costs associated with providing alternative operational logistics.

The Sea Ice Prediction Network South (SIPN South, <http://acecrc.org.au/sipn-south/>) is an international project endorsed by the Year of Polar Prediction (YOPP). One of its main goals is to make an initial assessment of the ability of current systems to predict Antarctic sea ice on hemispheric and regional scales, with a focus on the summer season. SIPN South has the ambition to **lay the foundations for a more systematic and coordinated evaluation of seasonal sea-ice forecasts in the Southern Ocean** in the coming years.

In February 2018, an initial assessment took place (Massonnet et al., 2018). 13 groups contributed 160 forecasts. Forecasts of the total Antarctic sea-ice area appeared consistent with observational verification data, but this agreement was, in fact, hiding regional errors. In particular, in observations, the Ross Sea happened to be almost entirely ice-free in February 2018 due to the passage of a cyclone in late January. All ensemble members of the model forecasts failed to forecast this anomaly, which suggested possible systematic shortcomings in the prediction systems in that sector.

The last milestone of SIPN South is the coordination of a seasonal sea-ice prediction exercise aligned with the YOPP Special Observing Period (YOPP-SOP) in the Southern Hemisphere, which spanned 16 November 2018 to 15 February 2019. The YOPP-SOP is a period of enhanced observational and modeling campaigns aiming at optimizing future observing systems and understanding the impact that selected observations can have on the skill of atmospheric and ocean—sea ice forecasts. Conversely, one of the objectives of SIPN South is to establish if seasonal forecasts can be of use to guide the location and timing of campaigns like those carried out during the SOP. The present document summarizes the main outcomes of this exercise.

2. Summer 2018-2019 in context

SIPN South analyses focus on austral summer, a season of special interest due to the intense marine traffic at this time of the year. In summer, sea ice retreats and exposes Antarctic coastlines to the open ocean, thereby offering possible access to the Antarctic continent, ice sheet or ice shelves.

According to the National Snow and Ice Data Center (NSIDC), the September 2018 sea-ice extent was the second lowest on record. **During austral spring, sea ice retreated anomalously fast** through December and January, setting record lows in early January. The melt slowed down in February.

Fig. 1 shows the evolution of February sea-ice extent since 1979 when satellite observations first became available. According to the NSIDC, **the monthly mean sea-**




Figure 1. February Antarctic sea-ice extent (Fetterer et al., 2017). The star is February 2019. The dashed line is the linear trend and the two shaded intervals show 1 and 2 standard deviations of the residuals around the linear fit, respectively.




Figure 2. Anomalies of sea-ice concentration in February 2019 relative to the 1981-2010 mean (from www.nsidc.org; Fetterer et al., 2017).

ice extent in February 2018 was the seventh lowest value on record (2.66 million km²) in a 41-yr long record time series. Spatially, positive sea-ice concentration anomalies occurred in the King Hakon VII and the East Antarctic sectors (0°E to 60°E). Anomalies were negative in the eastern Ross Sea, positive in the Amundsen Sea and mixed in the Weddell Sea (Fig. 2).

3. Forecasting sea ice for summer 2018-2019

A call for contributions was issued in November 2018 to predict sea-ice conditions during the three-month period from December 1st, 2018 to February 28th, 2019 (thus overlapping the YOPP Special Observing Period by 2 months). **We received a total of 12 submissions (198 forecasts) and would like to thank all contributors for their participation.** Tab. 1 summarizes the contributions received for this exercise.

Table 1. Information about contributors to the summer 2018-2019 coordinated sea ice forecast experiment.

	<i>Contributor name</i>	<i>Short name (in figures)</i>	<i>Forecasting method</i>	<i>Nb. of forecasts</i>	<i>Initialization date</i>	<i>Diagnostics provided</i>
1	Naval Research Lab	nrl	Coupled dynamical model	9	Oct. 31 st , 2018	SIA + rSIA + SIC
2	Nico Sun	Nico-Sun	Statistical model	3	Nov. 30 th , 2018	SIA + SIC
3	NASA-GMAO	nasa-gmao	Coupled dynamical model	10	Nov. 27 th , 2018	SIA + SIC
4	FIO-ESM	FIO-ESM	Coupled dynamical model	1	Nov. 1 st , 2018	SIA
5	ECMWF	ecmwf	Coupled dynamical model	50	Dec. 1 st , 2018	SIA + rSIA
6	Lamont Sea Ice Group	Lamont	Statistical model	1	Oct. 31 st , 2018	SIA + rSIA + SIC (monthly, interp. daily)
7	Alek Petty	Petty-NASA	Statistical model	1	Nov. 30 th , 2018	SIA (monthly, interp. daily)
8	Modified CanSIPS	Modified-CanSIPS	Coupled Dynamical Model	20	Nov. 30 th , 2018	SIA + rSIA
9	Met Office	MetOffice	Coupled Dynamical Model	42	Nov. 25 th , 2018	SIA + rSIA + SIC
10	CMCC	CMCC	Coupled Dynamical Model	50	Nov. 1 st , 2018	SIA
11	UCL	ucl	Ocean—sea ice Dynamical Model	10	July 1 st , 2018	SIA + rSIA + SIC
12	Sandra Barreira	Barreira	Statistical model	1	Dec. 1 st , 2018	SIA + rSIA + SIC (monthly, interp. daily)

Contributors were asked to provide, in order of descending priority, (1) the total Antarctic sea-ice area (denoted "SIA") for each day of December 2018-February 2019, (2) the sea-ice area per 10° longitude band (denoted "rSIA") for each day of December 2018-February 2019, and (3) sea-ice concentration (denoted "SIC") for each day of December 2018-February 2019. All contributors were able to submit (1), two submitted (1) and (2) only, two submitted (1) and (3) only, and five submitted (1), (2) and (3). Three submissions consisted of monthly mean forecasts. These forecasts

were interpolated to daily resolution using a quadratic function passing at the given monthly values on the 15th of each of the three months. Seven groups used fully coupled dynamical models and four groups used a statistical model trained on past data. One group used an ocean—sea ice model forced by atmospheric reanalyses of previous years.

We take note that requesting contributions for the first of the month is not ideal for those groups that produce monthly forecasts initialized at the beginning of each month, and will change our guidelines for subsequent exercises accordingly.

3.1 Circumpolar sea-ice area

Fig. 3 shows the total sea-ice area (SIA) forecast for each day of December 2018–February 2019 as submitted by the 12 contributors. SIA is not a very sensible geophysical diagnostic as it does not reflect regional variations, but it gives a first indication on how the forecasts behaved. In this figure, two observational references are also included to provide a general idea of the importance of observational uncertainty. As seen in Fig. 3, observational uncertainty is small relative to inter-model spread. In the following analyses, we will, therefore, assume that observational




Figure 3. Total (circumpolar) Antarctic sea-ice area of the 12 ensembles of forecasts for each day of the period December 2018–February 2019. The lines are the ensemble means and the shadings are the ensemble ranges. The superscripts in the legend indicate whether the submission is based on a statistical or a dynamical approach and, possibly, if monthly data has been interpolated to daily resolution. The black dashed lines are two observational references (Maslanik and Stroeve, 1999 and Tonboe et al., 2017).

errors are not a major cause for differences between forecasts and observations.

A striking feature in Fig. 3 is the overestimation, already at day 1 of the forecasting period, of the total sea-ice area by several groups. More particularly, this appears to be **a characteristic of several dynamical modeling contributions** (ucl, CMCC, nrl, Modified-CanSIPS). A closer look at dynamical contributions at day one of the forecasts (not shown here) reveals that this overestimation in total area is due, in most cases, to an overestimation of sea-ice concentration in the Ross Sea and, in some cases, to a too northward average position of the ice edge. The presence of biases in sea-ice concentration already at day one of the forecasting period reveals the challenges related to initialization of fully coupled or ocean-sea ice models. By contrast, **forecasts based on statistical models start on average closer to the two observational references.** Through the season, the high initial bias in the sea-ice area is progressively eliminated as the observed melt slows down from late December onwards, a feature not seen in the forecasts. During February, observed Antarctic sea-ice area lies in the full ensemble range. We note also that the full ensemble range of forecasted sea-ice area is similar to the historical range of sea-ice extent (Fig. 1).

We also investigate the ability of the systems to forecast the date of the annual minimum of sea-ice area (Fig. 4). The timing of the minimum of the sea-ice area is a critical parameter from an operational point of view, as it represents the end of the window of opportunity before the oceans start to freeze up and sea ice becomes an




Figure 4. Timing of the 2019 annual minimum of Antarctic sea-ice area from forecasts (colors) and two observational references (Maslanik and Stroeve, 1999 and Tonboe et al., 2017). To filter out the effects of synoptic variability, the minimum was determined from a quadratic fit of the February daily sea-ice area time series. Superscripts in the legend indicate whether the submission is based on a statistical or a dynamical approach and, possibly, if monthly data has been interpolated to daily resolution.

increasing hindrance to the progression of vessels. Last year, the minimum occurred around the 16th of February 2018 and most groups forecasted it to occur later. **This year, the minimum occurred late in the month (27th or 28th of February depending on the observational source) but the systems tended to collectively forecast a too early occurrence.**

3.2 Regional sea-ice area

Because of the strong regional character of Antarctic sea-ice variability, it is of importance to ascertain whether the overall agreement between forecasted and observed sea-ice areas in February (Fig. 3) is obtained for the good reasons or thanks to spatial error compensations. Fig. 5 shows the predicted February mean regional sea-ice area (rSIA), with the data expressed as an anomaly with respect to the 1979-2014 daily climatology estimated from the NASA Team sea-ice concentration dataset




Figure 5. February 2019 mean rSIA anomaly (compared to 1979-2014 NASA Team climatology) by longitude, for each submission, with observed estimates given in black. Solid lines show the ensemble mean for each contribution, with transparent shading indicating the ensemble range.

(Peng et al., 2013). The spread of the ensemble is particularly large in the Ross and Weddell Seas and none of the forecasts seems to capture the regional pattern of anomalies that occurred this year. The regional diagnostics presented in Fig. 5 reveal that **the circumpolar sea-ice area indeed masks strong regional biases and that**

several forecasts have the right total Antarctic sea-ice area for the wrong reasons.

A convenient approach to render the time evolution of regional biases of the sea-ice area is to compute the Integrated Ice Edge Error (IIEE; Goessling et al., 2016). The IIEE is a metric that quantifies the spatial mismatch between two geophysical datasets. It is oriented positively (always positive, with lower values indicating lower errors) and corresponds to the area of all grid cells where a given forecast and a given reference disagree on either one of the two following events: "sea-ice concentration is greater than 15%" or "sea-ice concentration is less than 15%". By design, the IIEE is not prone to cancellation of regional sea-ice area biases as is the total circumpolar area. Calculation of IIEE requires interpolation of the forecast and verification data to a common grid, which was chosen to be a regular $2^\circ \times 2^\circ$ grid.

The IIEE was applied to the seven contributions that provided spatial forecasts of sea-ice concentration. Fig. 6 displays the time evolution, over the forecasting period, of




Figure 6. Integrated Ice Edge Error (Goessling et al., 2016), defined as the area of grid cells where the forecasts and a reference (here, NSIDC-0081; Maslanik and Stroeve, 1999) disagree on concentration being either above or below 15%. The shadings represent ensemble range (IIEE calculated on each member separately) and the thick lines are the mean of all IIEEs for a given forecast system. The superscripts in the legend indicate whether the submission is based on a statistical or a dynamical approach and, possibly, if monthly data has been interpolated to daily resolution. The dark grey line is the IIEE between the other observational product (OSI-401-b; Tonboe et al., 2017) and the NSIDC-0081 reference.

that metric. Again, to gauge the possible role of observational uncertainty in forecast evaluation, the metric was applied to another observational dataset (OSI-401-b). The IIEE of that dataset as compared to the other observational dataset is at least one order of magnitude smaller than that from the forecasts, hence observational error can be assumed small compared to the forecast error.

Consistent with the results of sea-ice area (Fig. 1), the error is already large at day 1 of the forecasting period for most of the forecasts. The error first grows, as initial-condition information is lost progressively throughout the melting season. As discussed in Sec. 2 and seen from Fig. 3, observed sea ice retreated anomalously rapidly in December. From Fig. 6, it is seen that **all forecasting systems struggled to forecast the rapid ice retreat at the regional scale** but that a few systems, likely thanks to error compensations, can simulate that rapid retreat at the circumpolar level (e.g. Nico-Sun; Fig. 3). After the synchronous increase in IIEE during the month of December, the IIEE then decreases, for two reasons. First, observed ice melt slows down in January and February while forecasts keep melting ice at near climatological rates: the biases accumulated in December are progressively eliminated. Second, the surface of the ice to forecast evolves towards its minimum: with less ice to predict, there is less room for error. When normalizing the IIEE by the observed area (not shown here), forecast errors reach a plateau after one month (1st of January 2019) before slightly decreasing until the end of February. In any case, in all contributions, **a rapid error growth occurs during the first month of the forecast**, indicative of loss of predictive skill regardless of the prediction approach.

One forecast deserves particular attention as it outperforms the other ones (in the sense of the IIEE) through the entire period: Nico-Sun. This contribution is the only statistical one that provided daily information (the other statistical contributions were only available monthly and were interpolated to daily), so it is not possible at this stage to determine if statistical methods are generally superior to dynamical ones. The Nico-Sun method assumes that past day-to-day sea-ice concentration changes are representative of the conditions that may prevail for the coming forecast period. Starting from the latest NSIDC estimates, sea-ice concentration is updated day after day by adding increments estimated from past years. There is another state variable in the model (sea-ice thickness), that is also updated based on sea-ice melt estimated from the locally varying albedo due to sea-ice concentration changes. Despite its simplicity, the method appears to provide the most accurate forecast for this year. It is worth reminding that, for the exercise of last year, it is a dynamical contribution (NASA-GMAO) that was found to be the most consistent with observed data.

Drawing conclusions on which type of approach (statistical or dynamical) is superior to the other is therefore premature at this stage.

3.3 Spatial information

We finally display in Fig. 7 the probability of sea-ice presence for the 15th of February 2019. Green pixels are those where sea ice was forecast to be unlikely present, while red ones are those where sea ice was forecast to be likely present. The three statistical




Figure 7. Probability of ice presence for the 15th of February 2019, as forecasted by the seven groups that submitted daily sea-ice concentration information. The white lines are the actual ice edges from the verification datasets on that day. The probability of presence corresponds to the fraction of ensemble members that simulate sea-ice concentration larger than 15% in a given grid cell, for that day. A dynamic animation of that figure for all 28 days of February is available [here](#)

contributions (Lamont, Nico-Sun, Barreira) display sharp transitions between areas of certain ice presence and certain ice absence. By contrast, dynamical contributions suggest that, in some regions like the Ross Sea, sea ice presence was very uncertain. The dynamical model ensembles are designed to sample weather variability and results from Fig. 7 indicate that weather variability can imprint sea ice variability in key sectors like the Ross Sea, a region that was already very difficult to forecast last year. Whether those ensemble forecasts are correctly calibrated will be investigated once more retrospective forecasts will be available.

4. Conclusions

We warmly thank all 12 contributors to this second exercise of coordinated forecasts of sea ice in the Southern Ocean. The great enthusiasm for SIPN is much appreciated and we are looking forward to continuing our activities. Indeed, more hindcasts are necessary to ensure the robustness of the results. Still, this analysis has already revealed several elements:

- When viewed as a group, the range of multi-model forecast of total February Antarctic sea-ice area includes the two observational verification datasets. However, errors can be large for individual submissions. Observational uncertainty alone cannot explain the forecast-data mismatch.
- The timing of the minimum of Antarctic sea-ice area is not well predicted by the ensemble. The date of the minimum is in part driven by the change in insolation (which is predictable) and can be modulated by a few days by the passage of synoptic weather systems. Models, regardless of their nature, should capture weather uncertainty but it appears that ensemble spread is generally too narrow, i.e. that the systems are under-dispersive.
- At the regional level, the range of forecasts includes the observations in most of the sectors but individual forecasts show errors that tend to compensate when zonal averages are performed. Thus, the total area is not a suitable diagnostic for evaluating SIPN South forecasts.
- The only statistical contribution that provided daily information outperformed other contributions. Several dynamical models have difficulties in representing sea-ice concentration fields already on the first day of the forecasting period.
- At this stage, the SIPN South data set is not mature yet for practical use in applications like field trip planning or maritime route forecasting. Long records of retrospective forecasts are lacking in order to properly identify the origin of systematic forecast errors.

Data availability

The analyses presented in this report can be reproduced bit-wise by cloning the SIPN South Github project at <https://github.com/fmassonn/sipn-south-public> (commit d4b3feb4df84eabc107e34d6dab10be3be93e91f). Instructions to retrieve the data and process the analyses are given in the README.md file of this repository.

Citing this report

F. Massonnet, P. Reid, J. L. Lieser, C. M. Bitz, J. Fyfe, W. Hobbs (2019). Assessment of summer 2018-2019 sea-ice forecasts for the Southern Ocean. Technical Note, Université catholique de Louvain (2018), available at <http://acecrc.org.au/sipn-south/>

References

Fetterer, F., K. Knowles, W. Meier, M. Savoie, and A. K. Windnagel, 2017, updated daily. Sea Ice Index, Version 3 (G02135). Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/N5K072F8>. [Accessed May 2018].

Goessling, H. F., Tietsche, S., Day, J. J., Hawkins, E., & Jung, T. (2016). Predictability of the Arctic sea ice edge. *Geophysical Research Letters*, 43(4), 1642–1650. <https://doi.org/10.1002/2015gl067232>

Holland, M. M., Blanchard-Wrigglesworth, E., Kay, J., & Vavrus, S. (2013). Initial-value predictability of Antarctic sea ice in the Community Climate System Model 3. *Geophysical Research Letters*, 40(10), 2121–2124. <https://doi.org/10.1002/grl.50410>

Holland, M. M., Landrum, L., Raphael, M., & Stammerjohn, S. (2017). Springtime winds drive Ross Sea ice variability and change in the following autumn. *Nature Communications*, 8(1). <https://doi.org/10.1038/s41467-017-00820-0>

Marchi, S., Fichet, T., Goosse, H., Zunz, V., Tietsche, S., Day, J. J., & Hawkins, E. (2018). Reemergence of Antarctic sea ice predictability and its link to deep ocean mixing in global climate models. *Climate Dynamics*, 52(5–6), 2775–2797. <https://doi.org/10.1007/s00382-018-4292-2>

Maslanik, J. and J. Stroeve, 1999, updated daily. Near-Real-Time DMSP SSMIS Daily Polar Gridded Sea Ice Concentrations, Version 1. [NSIDC-0081]. Boulder, Colorado USA. NASA National Snow

Massonnet, F., P. Reid, J. L. Lieser, C. M. Bitz, J. Fyfe, W. Hobbs (2018). Assessment of February 2018 sea-ice forecasts for the Southern Ocean. <https://eprints.utas.edu.au/27184/>

Peng, G., W. Meier, D. Scott, and M. Savoie, 2013. A long-term and reproducible passive microwave sea ice concentration data record for climate studies and monitoring, *Earth Syst. Sci. Data*. 5. 311–318. <http://dx.doi.org/10.5194/essd-5-311-2013> and Ice Data Center Distributed Active Archive Center. doi: <http://dx.doi.org/10.5067/U8C09DWVX9LM>. [Accessed January 30th, 2018].

Tonboe, R., J. Lavelle, R. H. Pfeiffer and E. Howe, 2017. Product User Manual for OSI SAF Global Sea Ice Concentration (Product OSI-401-b). http://osisaf.met.no/docs/osisaf_cdop3_ss2_pum_ice-conc_v1p6.pdf [Accessed May 30th, 2018]