

Lecture 5

Ambiguity for the max plus semiring

Hierarchy of classes for weighted automata

- The inclusions are strict
for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$
and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Weighted automata (WA)

⊂

Polynomially ambiguous WA

⊂

Finitely ambiguous WA

⊂

Unambiguous WA

⊂

Deterministic WA

Hierarchy of classes for weighted automata

- The inclusions are strict
for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$
and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$
- Not over 1-letter alphabet
(last lecture)

Weighted automata (WA)

⊂

Polynomially ambiguous WA

⊂

Finitely ambiguous WA

⊂

Unambiguous WA

⊂

Deterministic WA

Hierarchy of classes for weighted automata

- The inclusions are strict
for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$
and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Not over 1-letter alphabet
(last lecture)

- We focus on $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$
unambiguous, finitely ambiguous and polynomially ambiguous

Weighted automata (WA)

⊂

Polynomially ambiguous WA

⊂

Finitely ambiguous WA

⊂

Unambiguous WA

⊂

Deterministic WA

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough

Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough

Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$

- By a case analysis $y \in a^*$, $y \in a^* b^*$ or $y \in b^*$ contradiction

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough
Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$
- By a case analysis $y \in a^*$, $y \in a^* b^*$ or $y \in b^*$ contradiction
- Pumping lemmas for weighted automata?

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough

Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$

- By a case analysis $y \in a^*$, $y \in a^* b^*$ or $y \in b^*$ contradiction

- Pumping lemmas for weighted automata?

- Fix $u \cdot v \cdot w \in \Sigma^*$

We say that $\hat{u} \cdot \hat{v} \cdot \hat{w} \in \Sigma^*$, is a *refinement* of $u \cdot v \cdot w$ if

1. $u \cdot v \cdot w = \hat{u} \cdot \hat{v} \cdot \hat{w}$

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough

Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$

- By a case analysis $y \in a^*$, $y \in a^* b^*$ or $y \in b^*$ contradiction

- Pumping lemmas for weighted automata?

- Fix $u \cdot v \cdot w \in \Sigma^*$

We say that $\hat{u} \cdot \hat{v} \cdot \hat{w} \in \Sigma^*$, is a *refinement* of $u \cdot v \cdot w$ if

1. $u \cdot v \cdot w = \hat{u} \cdot \hat{v} \cdot \hat{w}$

2. there exist u', w' such that $u \cdot u' = \hat{u}$, $w' \cdot w = \hat{w}$, $u' \cdot \hat{v} \cdot w' = v$, and $\hat{v} \neq \epsilon$.

Pumping lemmas

How to prove $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular?

- Pumping argument: let $w \in L$ and $|w|$ big enough

Then $w = xyz$ such that $xy^i z \in L$ for some $y \neq \epsilon$ and all $i \in \mathbb{N}$

- By a case analysis $y \in a^*$, $y \in a^* b^*$ or $y \in b^*$ contradiction

- Pumping lemmas for weighted automata?

- Fix $u \cdot v \cdot w \in \Sigma^*$

We say that $\hat{u} \cdot \hat{v} \cdot \hat{w} \in \Sigma^*$, is a *refinement* of $u \cdot v \cdot w$ if

1. $u \cdot v \cdot w = \hat{u} \cdot \hat{v} \cdot \hat{w}$

2. there exist u', w' such that $u \cdot u' = \hat{u}$, $w' \cdot w = \hat{w}$, $u' \cdot \hat{v} \cdot w' = v$, and $\hat{v} \neq \epsilon$.

Example: $aab \cdot \underline{bb} \cdot ba$ refines $aa \cdot bbbb \cdot a$

Unambiguous automata (1)

Fix the semiring $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Unambiguous automata (1)

Fix the semiring $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Theorem (M. and Riveros 2018)

Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$ be definable by unambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$. There exists N such that for all words of the form $u \cdot v \cdot w \in \Sigma^*$ with $|v| \geq N$, $v \neq \epsilon$, there exists a refinement $\hat{u} \cdot \hat{v} \cdot \hat{w}$ of $u \cdot \underline{v} \cdot w$ such that one of the conditions holds:

- (1) $f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- (2) $f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Unambiguous automata (1)

Fix the semiring $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Theorem (M. and Riveros 2018)

Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$ be definable by unambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$. There exists N such that for all words of the form $u \cdot v \cdot w \in \Sigma^*$ with $|v| \geq N$, $v \neq \epsilon$, there exists a refinement $\hat{u} \cdot \hat{v} \cdot \hat{w}$ of $u \cdot v \cdot w$ such that one of the conditions holds:

- (1) $f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- (2) $f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

- Example: $f(w) = \max(\#_a(w), \#_b(w))$, fix N from the lemma

Unambiguous automata (1)

Fix the semiring $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Theorem (M. and Riveros 2018)

Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$ be definable by unambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$. There exists N such that for all words of the form $u \cdot v \cdot w \in \Sigma^*$ with $|v| \geq N$, $v \neq \epsilon$, there exists a refinement $\hat{u} \cdot \hat{v} \cdot \hat{w}$ of $u \cdot v \cdot w$ such that one of the conditions holds:

$$(1) f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w}) \text{ for every } i \geq N.$$

$$(2) f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w}) \text{ for every } i \geq N.$$

- Example: $f(w) = \max(\#_a(w), \#_b(w))$, fix N from the lemma

$$f(a^{(N+1)^2} \cdot b^N \cdot \epsilon) = (N+1)^2, \text{ refining } a^{(N+1)^2} \cdot b^N \text{ we get } a^{(N+1)^2} b^n \cdot \underline{b}^m \cdot b^l$$

Unambiguous automata (1)

Fix the semiring $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Theorem (M. and Riveros 2018)

Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$ be definable by unambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$. There exists N such that for all words of the form $u \cdot v \cdot w \in \Sigma^*$ with $|v| \geq N$, $v \neq \epsilon$, there exists a refinement $\hat{u} \cdot \hat{v} \cdot \hat{w}$ of $u \cdot v \cdot w$ such that one of the conditions holds:

$$(1) f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w}) \text{ for every } i \geq N.$$

$$(2) f(\hat{u} \cdot \hat{v}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{v}^{i+1} \cdot \hat{w}) \text{ for every } i \geq N.$$

- Example: $f(w) = \max(\#_a(w), \#_b(w))$, fix N from the lemma

$$f(a^{(N+1)^2} \cdot b^N \cdot \epsilon) = (N+1)^2, \text{ refining } a^{(N+1)^2} \cdot b^N \text{ we get } a^{(N+1)^2} b^n \cdot \underline{b}^m \cdot b^l$$

- Since $n + mN + l < (N+1)^2$ then (1). But for i big enough (2)

Unambiguous automata (2)

Corollary

Unambiguous WA \subsetneq finitely ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Unambiguous automata (2)

Corollary

Unambiguous WA \subsetneq finitely ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Proof.

Let \mathcal{A} unambiguous automaton defining f . And let uvw with $v \geq N \gg 2^{|Q|}$

Unambiguous automata (2)

Corollary

Unambiguous WA \subsetneq finitely ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Proof.

Let \mathcal{A} unambiguous automaton defining f . And let uvw with $v \geq N \gg 2^{|Q|}$

- if $f(uvw) > -\infty$ then there is a unique accepting run on uvw

Let $q_0, \dots, q_{|v|}$ be the set of states on v

Unambiguous automata (2)

Corollary

Unambiguous WA \subsetneq finitely ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Proof.

Let \mathcal{A} unambiguous automaton defining f . And let uvw with $v \geq N \gg 2^{|Q|}$

- if $f(uvw) > -\infty$ then there is a unique accepting run on uvw

Let $q_0, \dots, q_{|v|}$ be the set of states on v

There is a cycle $\rho = q_i, \dots, q_j$ for $i < j$

If the value on ρ is 0 then (1) otherwise (2)

Unambiguous automata (2)

Corollary

Unambiguous WA \subsetneq finitely ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Proof.

Let \mathcal{A} unambiguous automaton defining f . And let uvw with $v \geq N \gg 2^{|Q|}$

- if $f(uvw) > -\infty$ then there is a unique accepting run on uvw

Let $q_0, \dots, q_{|v|}$ be the set of states on v

There is a cycle $\rho = q_i, \dots, q_j$ for $i < j$

If the value on ρ is 0 then (1) otherwise (2)

- if $f(uvw) = -\infty$ then since there is at most $n = \exp(|Q|)$ runs on uvw (\mathcal{A} is unambiguous)

Unambiguous automata (3)

We can present all runs on v as sequences

$$(q_{0,1}, \dots, q_{0,n}), \dots, (q_{|v|,1}, \dots, q_{|v|,n})$$

Unambiguous automata (3)

We can present all runs on v as sequences

$$(q_{0,1}, \dots, q_{0,n}), \dots, (q_{|v|,1}, \dots, q_{|v|,n})$$

- For $|v|$ big enough there is $i < j$ such that

$$(q_{i,1}, \dots, q_{i,n}) = (q_{j,1}, \dots, q_{j,n})$$

Unambiguous automata (3)

We can present all runs on v as sequences

$$(q_{0,1}, \dots, q_{0,n}), \dots, (q_{|v|,1}, \dots, q_{|v|,n})$$

- For $|v|$ big enough there is $i < j$ such that

$$(q_{i,1}, \dots, q_{i,n}) = (q_{j,1}, \dots, q_{j,n})$$

- Then i to j is a cycle for all old runs

the output for $uv^i w$ for all runs $1, \dots, n$ remain $-\infty$

Unambiguous automata (3)

We can present all runs on v as sequences

$$(q_{0,1}, \dots, q_{0,n}), \dots, (q_{|v|,1}, \dots, q_{|v|,n})$$

- For $|v|$ big enough there is $i < j$ such that

$$(q_{i,1}, \dots, q_{i,n}) = (q_{j,1}, \dots, q_{j,n})$$

- Then i to j is a cycle for all old runs

the output for $uv^i w$ for all runs $1, \dots, n$ remain $-\infty$

- If new runs occur then the number of runs for $uv^i w$ would be at least $n + i - 1$
(contradiction with finite ambiguity)



Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if $v_k = x_k \cdot y_k \cdot z_k$, $u'_k = z_k \cdot u_k \cdot x_{k+1}$; where $z_0 = x_{n+1} = \epsilon$.

and $y_k \neq \epsilon$ for every k

Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if $v_k = x_k \cdot y_k \cdot z_k$, $u'_k = z_k \cdot u_k \cdot x_{k+1}$; where $z_0 = x_{n+1} = \epsilon$.

and $y_k \neq \epsilon$ for every k

- Let $S \subseteq \{1, \dots, n\}$

Then $\underline{y_k}(S, i)$ is y_k^i if $k \in S$ and y_k otherwise

Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if $v_k = x_k \cdot y_k \cdot z_k$, $u'_k = z_k \cdot u_k \cdot x_{k+1}$; where $z_0 = x_{n+1} = \epsilon$.

and $y_k \neq \epsilon$ for every k

- Let $S \subseteq \{1, \dots, n\}$

Then $\underline{y_k}(S, i)$ is y_k^i if $k \in S$ and y_k otherwise

By $w(S, i)$ we denote the word

$$w = u'_0 \cdot \underline{y_1}(S, i) \cdot u'_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n}(S, i) \cdot u'_n.$$

Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if $v_k = x_k \cdot y_k \cdot z_k$, $u'_k = z_k \cdot u_k \cdot x_{k+1}$; where $z_0 = x_{n+1} = \epsilon$.

and $y_k \neq \epsilon$ for every k

- Let $S \subseteq \{1, \dots, n\}$

Then $\underline{y_k}(S, i)$ is y_k^i if $k \in S$ and y_k otherwise

By $w(S, i)$ we denote the word

$$w = u'_0 \cdot \underline{y_1}(S, i) \cdot u'_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n}(S, i) \cdot u'_n.$$

- Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$. A refinement is linear if

$f(w(S, i + i)) = K + f(w(S, i))$ for all i big enough

For linear refinements we denote $\Delta(S) = K$

Finitely ambiguous automata (1)

Let $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if $v_k = x_k \cdot y_k \cdot z_k$, $u'_k = z_k \cdot u_k \cdot x_{k+1}$; where $z_0 = x_{n+1} = \epsilon$.

and $y_k \neq \epsilon$ for every k

- Let $S \subseteq \{1, \dots, n\}$

Then $\underline{y_k}(S, i)$ is y_k^i if $k \in S$ and y_k otherwise

By $w(S, i)$ we denote the word

$$w = u'_0 \cdot \underline{y_1}(S, i) \cdot u'_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n}(S, i) \cdot u'_n.$$

- Let $f : \Sigma^* \rightarrow \mathbb{N} \cup \{-\infty\}$. A refinement is linear if

$f(w(S, i + i)) = K + f(w(S, i))$ for all i big enough

For linear refinements we denote $\Delta(S) = K$

S is decomposable if $\Delta(S) = \sum_{j \in S} \Delta(\{j\})$

Finitely ambiguous automata (2)

Theorem

Let f definable by finitely ambiguous automaton over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$.

There exists $N \in \mathbb{N}$ such that for every $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot \underline{v_n} \cdot u_n$, where $n \geq N$ and $|\underline{v_i}| \geq N$ for all i , there exists a linear refinement

$$w = x_0 \cdot \underline{y_1} \cdot x_1 \cdot \underline{y_2} \cdot \dots \cdot \underline{y_n} \cdot x_n$$

such that for every sequence of pairwise different, non-empty sets

$S_1, S_2, \dots, S_k \subseteq \{1, \dots, n\}$ with $k \geq N$, one of the following holds:

(1) exists j s.t. S_j is not decomposable

(2) exist j_1 and j_2 s.t. $\{l_1, l_2\}$ is decomposable for every $l_1 \in S_{j_1}$ and $l_2 \in S_{j_2}$.

Finitely ambiguous automata (2)

Theorem

Let f definable by finitely ambiguous automaton over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$.

There exists $N \in \mathbb{N}$ such that for every $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot \underline{v_n} \cdot u_n$, where $n \geq N$ and $|\underline{v_i}| \geq N$ for all i , there exists a linear refinement

$$w = x_0 \cdot \underline{y_1} \cdot x_1 \cdot \underline{y_2} \cdot \dots \cdot \underline{y_n} \cdot x_n$$

such that for every sequence of pairwise different, non-empty sets

$S_1, S_2, \dots, S_k \subseteq \{1, \dots, n\}$ with $k \geq N$, one of the following holds:

(1) exists j s.t. S_j is not decomposable

(2) exist j_1 and j_2 s.t. $\{l_1, l_2\}$ is decomposable for every $l_1 \in S_{j_1}$ and $l_2 \in S_{j_2}$.

- Example : f longest block of b 's. Let N from the lemma

Finitely ambiguous automata (2)

Theorem

Let f definable by finitely ambiguous automaton over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$.

There exists $N \in \mathbb{N}$ such that for every $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot \underline{v_n} \cdot u_n$, where $n \geq N$ and $|v_i| \geq N$ for all i , there exists a linear refinement

$$w = x_0 \cdot \underline{y_1} \cdot x_1 \cdot \underline{y_2} \cdot \dots \cdot \underline{y_n} \cdot x_n$$

such that for every sequence of pairwise different, non-empty sets

$S_1, S_2, \dots, S_k \subseteq \{1, \dots, n\}$ with $k \geq N$, one of the following holds:

(1) exists j s.t. S_j is not decomposable

(2) exist j_1 and j_2 s.t. $\{l_1, l_2\}$ is decomposable for every $l_1 \in S_{j_1}$ and $l_2 \in S_{j_2}$.

- Example : f longest block of b 's. Let N from the lemma

Let $(\underline{b^{N+1}a})^{N+1}$ and define $S_j = \{j\}$

Finitely ambiguous automata (2)

Theorem

Let f definable by finitely ambiguous automaton over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$.

There exists $N \in \mathbb{N}$ such that for every $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot \underline{v_n} \cdot u_n$, where $n \geq N$ and $|\underline{v_i}| \geq N$ for all i , there exists a linear refinement

$$w = x_0 \cdot \underline{y_1} \cdot x_1 \cdot \underline{y_2} \cdot \dots \cdot \underline{y_n} \cdot x_n$$

such that for every sequence of pairwise different, non-empty sets

$S_1, S_2, \dots, S_k \subseteq \{1, \dots, n\}$ with $k \geq N$, one of the following holds:

(1) exists j s.t. S_j is not decomposable

(2) exist j_1 and j_2 s.t. $\{l_1, l_2\}$ is decomposable for every $l_1 \in S_{j_1}$ and $l_2 \in S_{j_2}$.

- Example : f longest block of b 's. Let N from the lemma

Let $(\underline{b^{N+1}}a)^{N+1}$ and define $S_j = \{j\}$

Every S_j is decomposable but any $\{j_1, j_2\}$ is not decomposable

Finitely ambiguous automata (3)

Corollary

Finitely ambiguous WA \subsetneq poly-ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Finitely ambiguous automata (3)

Corollary

Finitely ambiguous WA \subsetneq poly-ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- **Proof.**

Suppose f is recognised by \mathcal{A} which is m -ambiguous, $|Q| = r$ and $f(w) > -\infty$

Finitely ambiguous automata (3)

Corollary

Finitely ambiguous WA \subsetneq poly-ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- **Proof.**

Suppose f is recognised by \mathcal{A} which is m -ambiguous, $|Q| = r$ and $f(w) > -\infty$

- Set $N = \max\{r^m, m\} + 1$.

The refinement is defined in such a way that every y_i is a cycle on all accepting runs

Finitely ambiguous automata (3)

Corollary

Finitely ambiguous WA \subsetneq poly-ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- **Proof.**

Suppose f is recognised by \mathcal{A} which is m -ambiguous, $|Q| = r$ and $f(w) > -\infty$

- Set $N = \max\{r^m, m\} + 1$.

The refinement is defined in such a way that every y_i is a cycle on all accepting runs

- Such a refinement is linear
and $\Delta(S)$ is determined by maximal cycle weights in blocks in S

Finitely ambiguous automata (3)

Corollary

Finitely ambiguous WA \subsetneq poly-ambiguous WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- **Proof.**

Suppose f is recognised by \mathcal{A} which is m -ambiguous, $|Q| = r$ and $f(w) > -\infty$

- Set $N = \max\{r^m, m\} + 1$.

The refinement is defined in such a way that every y_i is a cycle on all accepting runs

- Such a refinement is linear
and $\Delta(S)$ is determined by maximal cycle weights in blocks in S
- We will assume there are only accepting runs (to simplify technicalities)

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

- Notice that by pumping y_i the number of runs cannot increase (otherwise a contradiction with finitely ambiguous)

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

- Notice that by pumping y_i the number of runs cannot increase (otherwise a contradiction with finitely ambiguous)
- So for all S_j the number of runs in $w(S_j, i)$ is m and they come from ρ_1, \dots, ρ_m

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

- Notice that by pumping y_i the number of runs cannot increase (otherwise a contradiction with finitely ambiguous)
- So for all S_j the number of runs in $w(S_j, i)$ is m and they come from ρ_1, \dots, ρ_m
- For every $j \in \{1, \dots, n\}$ and $l \in \{1, \dots, m\}$ let $\rho_l[j]$ be the corresponding cycle
Let $wt(\rho_l[j])$ be its weight

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

- Notice that by pumping y_i the number of runs cannot increase (otherwise a contradiction with finitely ambiguous)
- So for all S_j the number of runs in $w(S_j, i)$ is m and they come from ρ_1, \dots, ρ_m
- For every $j \in \{1, \dots, n\}$ and $l \in \{1, \dots, m\}$ let $\rho_l[j]$ be the corresponding cycle
Let $wt(\rho_l[j])$ be its weight
- A cycle $\rho_l[j]$ is dominant if $wt(\rho_l[j]) \geq wt(\rho_{l'}[j])$ for all $l' \in \{1, \dots, m\}$

Finitely ambiguous automata (4)

Denote runs by ρ_1, \dots, ρ_m

- Notice that by pumping y_i the number of runs cannot increase (otherwise a contradiction with finitely ambiguous)
- So for all S_j the number of runs in $w(S_j, i)$ is m and they come from ρ_1, \dots, ρ_m
- For every $j \in \{1, \dots, n\}$ and $l \in \{1, \dots, m\}$ let $\rho_l[j]$ be the corresponding cycle
Let $wt(\rho_l[j])$ be its weight
- A cycle $\rho_l[j]$ is dominant if $wt(\rho_l[j]) \geq wt(\rho_{l'}[j])$ for all $l' \in \{1, \dots, m\}$
- If a cycle $\rho_l[j]$ is dominant then $\Delta(\{j\}) = wt(\rho_l[j])$

Finitely ambiguous automata (5)

Lemma

Let $S \subseteq \{1, \dots, n\}$ a linear refinement. Then S is decomposable iff for one of the runs ρ the cycle $\rho[j]$ is dominant for all $j \in S$.

Finitely ambiguous automata (5)

Lemma

Let $S \subseteq \{1, \dots, n\}$ a linear refinement. Then S is decomposable iff for one of the runs ρ the cycle $\rho[j]$ is dominant for all $j \in S$.

Proof. (\implies)

Let ρ be such that $\sum_{j \in S} wt(\rho[j])$ is maximal

Finitely ambiguous automata (5)

Lemma

Let $S \subseteq \{1, \dots, n\}$ a linear refinement. Then S is decomposable iff for one of the runs ρ the cycle $\rho[j]$ is dominant for all $j \in S$.

Proof. (\implies)

Let ρ be such that $\sum_{j \in S} wt(\rho[j])$ is maximal

- We claim that ρ satisfies the lemma

By definition $\Delta(S) = \sum_{j \in S} wt(\rho[j])$

Finitely ambiguous automata (5)

Lemma

Let $S \subseteq \{1, \dots, n\}$ a linear refinement. Then S is decomposable iff for one of the runs ρ the cycle $\rho[j]$ is dominant for all $j \in S$.

Proof. (\implies)

Let ρ be such that $\sum_{j \in S} wt(\rho[j])$ is maximal

- We claim that ρ satisfies the lemma

By definition $\Delta(S) = \sum_{j \in S} wt(\rho[j])$

- For a contradiction let j^* s.t. $wt(\rho[j^*]) < \Delta(\{j^*\})$

Then $\Delta(S) = \sum_{j \in S} wt(\rho[j]) < \sum_{j \in S} \Delta(\{j\})$

Finitely ambiguous automata (5)

Lemma

Let $S \subseteq \{1, \dots, n\}$ a linear refinement. Then S is decomposable iff for one of the runs ρ the cycle $\rho[j]$ is dominant for all $j \in S$.

Proof. (\implies)

Let ρ be such that $\sum_{j \in S} wt(\rho[j])$ is maximal

- We claim that ρ satisfies the lemma

By definition $\Delta(S) = \sum_{j \in S} wt(\rho[j])$

- For a contradiction let j^* s.t. $wt(\rho[j^*]) < \Delta(\{j^*\})$

Then $\Delta(S) = \sum_{j \in S} wt(\rho[j]) < \sum_{j \in S} \Delta(\{j\})$

- A contradiction with S being linear

Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

- Then $\sum_{j \in S} wt(\rho[j]) \geq \sum_{j \in S} wt(\rho'[j])$ for any other run ρ'

Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

- Then $\sum_{j \in S} wt(\rho[j]) \geq \sum_{j \in S} wt(\rho'[j])$ for any other run ρ'
- So when pumping $w(S, i)$ the value increases by $\sum_{j \in S} wt(\rho[j])$
Which is equal to $\sum_{j \in S} \Delta(\{j\})$ since cycles are dominant



Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

- Then $\sum_{j \in S} wt(\rho[j]) \geq \sum_{j \in S} wt(\rho'[j])$ for any other run ρ'
- So when pumping $w(S, i)$ the value increases by $\sum_{j \in S} wt(\rho[j])$
Which is equal to $\sum_{j \in S} \Delta(\{j\})$ since cycles are dominant



- To prove the theorem suppose all S_j are decomposable
By lemma for all S_j there is ρ_{I_j} in which all cycles are dominant

Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

- Then $\sum_{j \in S} wt(\rho[j]) \geq \sum_{j \in S} wt(\rho'[j])$ for any other run ρ'
- So when pumping $w(S, i)$ the value increases by $\sum_{j \in S} wt(\rho[j])$
Which is equal to $\sum_{j \in S} \Delta(\{j\})$ since cycles are dominant



- To prove the theorem suppose all S_j are decomposable

By lemma for all S_j there is ρ_{l_j} in which all cycles are dominant

- By choice of N there are more sets than runs so $l_{j_1} = l_{j_2}$ for some $j_1 \neq j_2$

Finitely ambiguous automata (6)

(\Leftarrow) Let ρ be s.t. $\rho[j]$ is dominant for all $j \in S$

- Then $\sum_{j \in S} wt(\rho[j]) \geq \sum_{j \in S} wt(\rho'[j])$ for any other run ρ'
- So when pumping $w(S, i)$ the value increases by $\sum_{j \in S} wt(\rho[j])$
Which is equal to $\sum_{j \in S} \Delta(\{j\})$ since cycles are dominant



- To prove the theorem suppose all S_j are decomposable

By lemma for all S_j there is ρ_{l_j} in which all cycles are dominant

- By choice of N there are more sets than runs so $l_{j_1} = l_{j_2}$ for some $j_1 \neq j_2$
- By lemma $\{k_1, k_2\}$ is decomposable for every $k_1 \in S_{j_1}$ and $k_2 \in S_{j_2}$



Polynomially ambiguous automata (1)

Let S_1, \dots, S_m over $\{1, \dots, n\}$ a *partition* (S_i nonempty, pairwise disjoint)

We say that $S \subseteq \{1, \dots, n\}$ is a *selection set* if $|S \cap S_i| = 1$ for every i .

Polynomially ambiguous automata (1)

Let S_1, \dots, S_m over $\{1, \dots, n\}$ a *partition* (S_i nonempty, pairwise disjoint)

We say that $S \subseteq \{1, \dots, n\}$ is a *selection set* if $|S \cap S_i| = 1$ for every i .

Theorem

Let f poly-ambiguous over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$. There exist N and a function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$, where $|v_i| \geq N$ for every $1 \leq i \leq n$, there exists a linear refinement

$$w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n,$$


such that for every partition $\pi = S_1, S_2, \dots, S_m$ of $\{1, \dots, n\}$ with $m \geq \varphi(\max_j(|S_j|))$ one of the following holds:

- (1) there exists j such that S_j is decomposable;
- (2) there exists a selection set S for π such that S is not decomposable.

Polynomially ambiguous automata (2)

Example $w_0 \# w_1 \# \dots \# w_n$ with $w_i \in \{a, b\}^*$


$$f(w) = \sum_{i=0}^n \max\{|w_i|_a, |w_i|_b\}$$



Polynomially ambiguous automata (2)

Example $w_0 \# w_1 \# \dots \# w_n$ with $w_i \in \{a, b\}^*$

$$f(w) = \sum_{i=0}^n \max\{|w_i|_a, |w_i|_b\}$$




- Fix N and φ from the theorem. Let $m \geq \varphi(2)$
consider refinements of $(\underline{a^N} \underline{b^N} \#)^m$

Polynomially ambiguous automata (2)

Example $w_0 \# w_1 \# \dots \# w_n$ with $w_i \in \{a, b\}^*$

$$f(w) = \sum_{i=0}^n \max\{|w_i|_a, |w_i|_b\}$$




- Fix N and φ from the theorem. Let $m \geq \varphi(2)$
consider refinements of $(\underline{a^N} \underline{b^N} \#)^m$
denote by $j, j' \in \{1, \dots, m\}$ blocks of a and b $S_j = \{j, j'\}$ (not decomposable)

Polynomially ambiguous automata (2)

Example $w_0 \# w_1 \# \dots \# w_n$ with $w_i \in \{a, b\}^*$

$$f(w) = \sum_{i=0}^n \max\{|w_i|_a, |w_i|_b\}$$



- Fix N and φ from the theorem. Let $m \geq \varphi(2)$
consider refinements of $(\underline{a^N} \underline{b^N} \#)^m$
denote by $j, j' \in \{1, \dots, m\}$ blocks of a and b $S_j = \{j, j'\}$ (not decomposable)
But every selection set is decomposable

Polynomially ambiguous automata (3)

Corollary

Poly-ambiguous WA \subsetneq WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Polynomially ambiguous automata (3)

Corollary

Poly-ambiguous WA \subsetneq WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Decision problems for weighted automata

- Is a given automaton \mathcal{A} finitely ambiguous, polynomially ambiguous?
in NLOGSPACE (tutorials)

Polynomially ambiguous automata (3)

Corollary

Poly-ambiguous WA \subsetneq WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Decision problems for weighted automata

- Is a given automaton \mathcal{A} finitely ambiguous, polynomially ambiguous?
in NLOGSPACE (tutorials)
- Given an automaton \mathcal{A} is there an equivalent deterministic automaton \mathcal{B} ?
Long-standing open problem for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Polynomially ambiguous automata (3)

Corollary

Poly-ambiguous WA \subsetneq WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Decision problems for weighted automata

- Is a given automaton \mathcal{A} finitely ambiguous, polynomially ambiguous?
in NLOGSPACE (tutorials)
- Given an automaton \mathcal{A} is there an equivalent deterministic automaton \mathcal{B} ?
Long-standing open problem for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$
(less popular) open problem for $(\mathbb{Q}, +, \cdot, 0, 1)$

Polynomially ambiguous automata (3)

Corollary

Poly-ambiguous WA \subsetneq WA over $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

Decision problems for weighted automata

- Is a given automaton \mathcal{A} finitely ambiguous, polynomially ambiguous?
in NLOGSPACE (tutorials)
- Given an automaton \mathcal{A} is there an equivalent deterministic automaton \mathcal{B} ?
Long-standing open problem for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$
(less popular) open problem for $(\mathbb{Q}, +, \cdot, 0, 1)$
- Partial results: decidable for $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ and $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$
If we assume that \mathcal{A} is unambiguous, finitely ambiguous or poly-ambiguous

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

- **Emptiness:** Given \mathcal{A} is there a word w such that:

$$\llbracket \mathcal{A} \rrbracket (w) = c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \geq c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \leq c$$

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

- **Emptiness:** Given \mathcal{A} is there a word w such that:

$$\llbracket \mathcal{A} \rrbracket (w) = c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \geq c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \leq c$$

Usually undecidable or trivially decidable

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

- **Emptiness:** Given \mathcal{A} is there a word w such that:

$$\llbracket \mathcal{A} \rrbracket (w) = c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \geq c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \leq c$$

Usually undecidable or trivially decidable

$\llbracket \mathcal{A} \rrbracket (w) = 0$ undecidable for $(\mathbb{Q}, +, \cdot, 0, 1)$ (next slide)

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

- **Emptiness:** Given \mathcal{A} is there a word w such that:

$$\llbracket \mathcal{A} \rrbracket (w) = c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \geq c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \leq c$$

Usually undecidable or trivially decidable

$\llbracket \mathcal{A} \rrbracket (w) = 0$ undecidable for $(\mathbb{Q}, +, \cdot, 0, 1)$ (next slide)

But assuming 1-letter alphabet this is equivalent to the problem:

Given a linear recursive sequence u_0, u_1, \dots is there $n \in \mathbb{N}$ s.t. $u_n = 0$

Decision problems for weighted automata (2)

Variants of the classical emptiness problems for finite automata

- **Emptiness:** Given \mathcal{A} is there a word w such that:

$$\llbracket \mathcal{A} \rrbracket (w) = c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \geq c, \quad \text{or } \llbracket \mathcal{A} \rrbracket (w) \leq c$$

Usually undecidable or trivially decidable

$\llbracket \mathcal{A} \rrbracket (w) = 0$ undecidable for $(\mathbb{Q}, +, \cdot, 0, 1)$ (next slide)

But assuming 1-letter alphabet this is equivalent to the problem:

Given a linear recursive sequence u_0, u_1, \dots is there $n \in \mathbb{N}$ s.t. $u_n = 0$

This is known as the Skolem problem (open for many years)

Next two weeks there will be a result related to this problem

Decision problems for weighted automata (3)

Theorem (Bertoni 1974)

Fix the semiring $(\mathbb{Q}, +, \cdot, 0, 1)$. The problem if given \mathcal{A} is there a word w such that $\llbracket \mathcal{A} \rrbracket (w) = 0$ is undecidable.

Decision problems for weighted automata (3)

Theorem (Bertoni 1974)

Fix the semiring $(\mathbb{Q}, +, \cdot, 0, 1)$. The problem if given \mathcal{A} is there a word w such that $\llbracket \mathcal{A} \rrbracket (w) = 0$ is undecidable.

Proof.

We reduce from the Post correspondence problem

Given two morphisms $\varphi_1, \varphi_2 : \Sigma^* \rightarrow \{0, 1\}^*$ is there a word w s.t. $\varphi_1(w) = \varphi_2(w)$

Decision problems for weighted automata (3)

Theorem (Bertoni 1974)

Fix the semiring $(\mathbb{Q}, +, \cdot, 0, 1)$. The problem if given \mathcal{A} is there a word w such that $\llbracket \mathcal{A} \rrbracket(w) = 0$ is undecidable.

Proof.

We reduce from the Post correspondence problem

Given two morphisms $\varphi_1, \varphi_2 : \Sigma^* \rightarrow \{0, 1\}^*$ is there a word w
s.t. $\varphi_1(w) = \varphi_2(w)$

For example $\varphi_1(a) = 0, \varphi_1(b) = 10011, \varphi_2(a) = 001, \varphi_2(b) = 1$

Decision problems for weighted automata (3)

Theorem (Bertoni 1974)

Fix the semiring $(\mathbb{Q}, +, \cdot, 0, 1)$. The problem if given \mathcal{A} is there a word w such that $\llbracket \mathcal{A} \rrbracket(w) = 0$ is undecidable.

Proof.

We reduce from the Post correspondence problem

Given two morphisms $\varphi_1, \varphi_2 : \Sigma^* \rightarrow \{0, 1\}^*$ is there a word w

s.t. $\varphi_1(w) = \varphi_2(w)$

For example $\varphi_1(a) = 0$, $\varphi_1(b) = 10011$, $\varphi_2(a) = 001$, $\varphi_2(b) = 1$

Then $\varphi_1(aab) = \varphi_2(aab) = 0010011$

Decision problems for weighted automata (3)

Theorem (Bertoni 1974)

Fix the semiring $(\mathbb{Q}, +, \cdot, 0, 1)$. The problem if given \mathcal{A} is there a word w such that $\llbracket \mathcal{A} \rrbracket(w) = 0$ is undecidable.

Proof.

We reduce from the Post correspondence problem


Given two morphisms $\varphi_1, \varphi_2 : \Sigma^* \rightarrow \{0, 1\}^*$ is there a word w
s.t. $\varphi_1(w) = \varphi_2(w)$

For example $\varphi_1(a) = 0, \varphi_1(b) = 10011, \varphi_2(a) = 001, \varphi_2(b) = 1$

Then $\varphi_1(aab) = \varphi_2(aab) = 0010011$


- One can assume that if such a w exists then the last letter of w is 1

Decision problems for weighted automata (4)




Decision problems for weighted automata (4)

- $\mathcal{A}(a_1 a_2 \dots a_n) = \text{bin}(a_1 \dots a_n)$
 $= \sum_{i=1}^n \frac{a_i}{2^i}$



Decision problems for weighted automata (4)


- $\mathcal{A}(a_1 a_2 \dots a_n) = \text{bin}(a_1 \dots a_n)$
 $= \sum_{i=1}^n \frac{a_i}{2^i}$



- Notice that (almost) every w has a unique value $\llbracket \mathcal{A} \rrbracket (w)$
 (assuming w ends with 1)


Decision problems for weighted automata (4)

- $\mathcal{A}(a_1 a_2 \dots a_n) = \text{bin}(a_1 \dots a_n)$
 $= \sum_{i=1}^n \frac{a_i}{2^i}$




- Notice that (almost) every w has a unique value $\llbracket \mathcal{A} \rrbracket (w)$
 (assuming w ends with 1)
- Given a morphism $\varphi : \Sigma^* \rightarrow \{0, 1\}^*$ we define \mathcal{A}_φ s.t.
 $\llbracket \mathcal{A}_\varphi \rrbracket (w) = \text{bin}(\varphi(w))$

Decision problems for weighted automata (5)




Decision problems for weighted automata (5)

- Say $\varphi(a) = 01$




Decision problems for weighted automata (5)

- Say $\varphi(a) = 01$



Decision problems for weighted automata (5)


- Say $\varphi(a) = 01$



- Define \mathcal{A}_{φ_1} and \mathcal{A}_{φ_2}

Decision problems for weighted automata (5)

- Say $\varphi(a) = 01$



- Define \mathcal{A}_{φ_1} and \mathcal{A}_{φ_2}
- Then \mathcal{A} defined as $\mathcal{A}_{\varphi_1} - \mathcal{A}_{\varphi_2}$ has the property that $\mathcal{A}(w) = 0$ iff the Post correspondence instance is valid

