

Pumping lemmas for weighted automata

Filip Mazowiecki¹ and Cristian Riveros²

¹University of Bordeaux

²Pontificia Universidad Católica de Chile

Pumping lemmas for weighted automata


Filip Mazowiecki¹ and Cristian Riveros²

¹University of Bordeaux


²Pontificia Universidad Católica de Chile

ULB 2018

Automata




Automata



$$f : \Sigma^* \rightarrow \{0, 1\}$$

Automata




$$f : \Sigma^* \rightarrow \{0, 1\}$$

Weighted automata

$$f : \Sigma^* \rightarrow \text{"some numbers"}$$

Automata



$$f : \Sigma^* \rightarrow \{0, 1\}$$

Weighted automata


$$f : \Sigma^* \rightarrow \text{"some numbers"}? \quad \mathbb{N}?$$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$


Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Weighted automata (WA)


Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




Consider $w = bbab$

b	b	a	b
1	1	0	0

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




Consider $w = bbab$

b b a b

$$1 + 1 + 0 + 0 = 2$$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 & + & 1 & + & 0 & + & 0 & = & 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 & 0 & 0 & 1 \end{array}$$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$


$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$

Output: $\min\{2, 1\} = 1$

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$


$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$

Output: $\min\{2, 1\} = 1$

In general: \odot transitions, \oplus accepting runs

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$


Output: $\min\{2, 1\} = 1$

In general: \odot transitions, \oplus accepting runs

\emptyset if there is no accepting run

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$

Output: $\min\{2, 1\} = 1$


In general: \odot transitions, \oplus accepting runs

\emptyset if there is no accepting run

“smallest block of b’s”

Weighted automata (WA)

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



Consider $w = bbab$

$$\begin{array}{cccc} b & b & a & b \\ 1 + 1 + 0 + 0 = 2 \end{array}$$

$$\begin{array}{cccc} b & b & a & b \\ 0 + 0 + 0 + 1 = 1 \end{array}$$

Output: $\min\{2, 1\} = 1$

In general: \odot transitions, \oplus accepting runs

\emptyset if there is no accepting run

“smallest block of b’s”

(∞ if there is no b)

WA subclasses


Number of accepting runs?

WA subclasses

Number of accepting runs?

- could be exponential

accepting runs: 2^n (for a^n)



WA subclasses


Number of accepting runs?

- could be exponential

accepting runs: 2^n (for a^n)

- “smallest block of b ’s”

accepting runs: blocks of b ’s (linear)



WA subclasses

Number of accepting runs?


- could be exponential

accepting runs: 2^n (for a^n)

- “smallest block of b 's”

accepting runs: blocks of b 's (linear)

- $\min_{a \in \Sigma} \{ \text{number of } a\text{'s} \}$?




WA subclasses

Number of accepting runs?

- could be exponential

accepting runs: 2^n (for a^n)



- “smallest block of b 's”

accepting runs: blocks of b 's (linear)

- $\min_{a \in \Sigma} \{ \text{number of } a\text{'s} \}?$

accepting runs: $|\Sigma|$ (constant)

$a / 1, b / 0$



$a / 0, b / 1$




WA subclasses

Number of accepting runs?

- could be exponential

accepting runs: 2^n (for a^n)




- “smallest block of b 's”

accepting runs: blocks of b 's (linear)


- $\min_{a \in \Sigma} \{ \text{number of } a\text{'s} \}$?

accepting runs: $|\Sigma|$ (constant)



- f longest suffix of a 's; $f(abaa) = 2$

accepting runs: 1




Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$

Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$


State of art



Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$


State of art



Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$


State of art



Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$


State of art



Hierarchy of subclasses

Fix $N_\infty(\min, +, \infty, 0)$

State of art




- Strictness shown by examples

Hierarchy of subclasses

Fix $\mathbb{N}_\infty(\min, +, \infty, 0)$

State of art



- Strictness shown by examples
- Papers are about determinization

Meanwhile other formalisms

Boolean world

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Solution: pumping lemma

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Solution: pumping lemma

Take $w \in L$ big enough

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Solution: pumping lemma

Take $w \in L$ big enough

exists a decomposition $w = xyz$, $|y| > 0$

s.t. $xy^n z \in L$ for all n

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Solution: pumping lemma

Take $w \in L$ big enough

exists a decomposition $w = xyz$, $|y| > 0$

s.t. $xy^n z \in L$ for all n

quick case analysis

Meanwhile other formalisms

Boolean world

- Finite automata

Show that $L = \{a^n b^n \mid n \in \mathbb{N}\}$ is not regular.

Solution: pumping lemma

Take $w \in L$ big enough

exists a decomposition $w = xyz$, $|y| > 0$

s.t. $xy^n z \in L$ for all n

quick case analysis

- Context-free languages – pumping lemmas
- First order logic – Ehrenfeucht-Fraïssé games

On which fragments we work?

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$

On which fragments we work?

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$

WA

\cup

polynomially ambiguous WA

\cup


finitely ambiguous WA

\cup

unambiguous WA

On which fragments we work?

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$




On which fragments we work?

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



On which fragments we work?

Semiring: $\mathbb{N}_\infty(\min, +, \infty, 0)$



The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot v \cdot w \in \Sigma^*$

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

The no \min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}

e.g. $aaab\underline{bbbbb}aa$ is refined by $aaab\underline{bbbbb}aa$

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}
e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1)

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}
e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1)

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$;
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}
e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1)

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$;
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Example: f – longest suffix of a 's

Take any $u \cdot v \cdot w$, $N = 1$, and trivial refinement

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}

e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1)

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$;
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$. ← $v, w \in a^*$

Example: f – longest suffix of a 's

Take any $u \cdot v \cdot w$, $N = 1$, and trivial refinement

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}

e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1)

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$; ← otherwise
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$. ← $v, w \in a^*$

Example: f – longest suffix of a 's

Take any $u \cdot v \cdot w$, $N = 1$, and trivial refinement

The no min fragment

Unambiguous WA (U-WA) over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Let $u \cdot \underline{v} \cdot w \in \Sigma^*$

A refinement is $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w} = u \cdot \underline{v} \cdot w$ s.t. $\hat{\underline{v}} \neq \epsilon$ is a fragment of \underline{v}

e.g. $aaabbbbaa$ is refined by $aaabbbbaa$

Theorem (Pumping Lemma 1) works for a broader class

Let f recognizable by an U-WA over $(\min, +)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$; otherwise
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$. $v, w \in a^*$

Example: f – longest suffix of a 's

Take any $u \cdot v \cdot w$, $N = 1$, and trivial refinement

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Let $u \cdot v \cdot w = a^{(N+1)^2} \underline{b^N}$, $f(u \cdot v \cdot w) = N$

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Let $u \cdot v \cdot w = a^{(N+1)^2} \underline{b^N}$, $f(u \cdot v \cdot w) = N$

Any refinement $\hat{u} \cdot \hat{v} \cdot \hat{w} = a^{(N+1)^2} \cdot b^n \underline{b^m} b^l$, $1 \leq m \leq N$

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Let $u \cdot v \cdot w = a^{(N+1)^2} \underline{b^N}$, $f(u \cdot v \cdot w) = N$

Any refinement $\hat{u} \cdot \hat{v} \cdot \hat{w} = a^{(N+1)^2} \cdot b^n \underline{b^m} b^l$, $1 \leq m \leq N$

$f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = (N+1)^2$ for i big enough

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Let $u \cdot v \cdot w = a^{(N+1)^2} \underline{b^N}$, $f(u \cdot v \cdot w) = N$

Any refinement $\hat{u} \cdot \hat{v} \cdot \hat{w} = a^{(N+1)^2} \cdot b^n \underline{b^m} b^l$, $1 \leq m \leq N$

$f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = (N+1)^2$ for i big enough

but take $i = N$, then $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) \leq N + mN \leq N + N^2 < (N+1)^2$

Theorem (Pumping lemma 1)

Let f be a WA over $\mathbb{N}_\infty(\min, +, \infty, 0)$

there exists N s.t. for every $u \cdot \underline{v} \cdot w$ with $|v| \geq N$

there is a refinement $\hat{u} \cdot \hat{\underline{v}} \cdot \hat{w}$ and either:

- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.
- $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) < f(\hat{u} \cdot \hat{\underline{v}}^{i+1} \cdot \hat{w})$ for every $i \geq N$.

Let $f(w) = \min(\#_a(w), \#_b(w))$ ($f \in \text{FA-WA}$)

Let $u \cdot v \cdot w = a^{(N+1)^2} \underline{b^N}$, $f(u \cdot v \cdot w) = N$

Any refinement $\hat{u} \cdot \hat{v} \cdot \hat{w} = a^{(N+1)^2} \cdot b^n \underline{b^m} b^l$, $1 \leq m \leq N$

$f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) = (N+1)^2$ for i big enough

but take $i = N$, then $f(\hat{u} \cdot \hat{\underline{v}}^i \cdot \hat{w}) \leq N + mN \leq N + N^2 < (N+1)^2$

Corollary: $\text{U-WA} \subsetneq \text{FA-WA}$ over $(\min, +)$

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if y_i refine v_i

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if y_i refine v_i

Let $S \subseteq \{1, \dots, n\}$, $\underline{y_k}(S, i) = y_k^i$ if $k \in S$ and $\underline{y_k}(S, i) = y_k$ otherwise.

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if y_i refine v_i

Let $S \subseteq \{1, \dots, n\}$, $\underline{y_k}(S, i) = y_k^i$ if $k \in S$ and $\underline{y_k}(S, i) = y_k$ otherwise.

$w(S, i) = u_0 \cdot \underline{y_1}(S, i) \cdot u_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u_{n-1} \cdot \underline{y_n}(S, i) \cdot u_n$

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if y_i refine v_i

Let $S \subseteq \{1, \dots, n\}$, $\underline{y_k}(S, i) = y_k^i$ if $k \in S$ and $\underline{y_k}(S, i) = y_k$ otherwise.

$w(S, i) = u_0 \cdot \underline{y_1}(S, i) \cdot u_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u_{n-1} \cdot \underline{y_n}(S, i) \cdot u_n$

Example, a $(3, 2)$ -representation

$w = ab^3a\underline{ab}^2\underline{ab}^2aa$

Preparing for the next lemma

Word n -representation: $w = u_0 \cdot \underline{v_1} \cdot u_1 \cdot \underline{v_2} \cdot \dots \cdot u_{n-1} \cdot \underline{v_n} \cdot u_n$

(n, N) -representation: $|v_i| \geq N$ for all i

A refinement is $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

if y_i refine v_i

Let $S \subseteq \{1, \dots, n\}$, $\underline{y_k}(S, i) = y_k^i$ if $k \in S$ and $\underline{y_k}(S, i) = y_k$ otherwise.

$w(S, i) = u_0 \cdot \underline{y_1}(S, i) \cdot u_1 \cdot \underline{y_2}(S, i) \cdot \dots \cdot u_{n-1} \cdot \underline{y_n}(S, i) \cdot u_n$

Example, a $(3, 2)$ -representation

$w = ab^3a\underline{ab^2}a\underline{ab^2}aa$

$w(\{1, 3\}, 3) = a\underline{b^9}a\underline{ab^2}a\underline{b^6}aa$

Theorem (Pumping Lemma 2)

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations $(n \geq N)$

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$

there exists N s.t. for all (n, N) -representations $(n \geq N)$

exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of b 's”

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of $b's$ ”

Let $w = (\underline{b^N} a^N)^N$ ($n = N$)

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of $b's$ ”

Let $w = (\underline{b^N} a^N)^N$ ($n = N$)

Let $S_j = \{1, \dots, N\} \setminus \{j\}$, $f(w(S_j, i)) = N$ for all i, j

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of $b's$ ”

Let $w = (\underline{b^N} a^N)^N$ ($n = N$)

Let $S_j = \{1, \dots, N\} \setminus \{j\}$, $f(w(S_j, i)) = N$ for all i, j

But $S_{j_1} \cup S_{j_2} = \{1, \dots, N\}$ for $j_1 \neq j_2$

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of $b's$ ”

Let $w = (\underline{b^N} a^N)^N$ ($n = N$)

Let $S_j = \{1, \dots, N\} \setminus \{j\}$, $f(w(S_j, i)) = N$ for all i, j

But $S_{j_1} \cup S_{j_2} = \{1, \dots, N\}$ for $j_1 \neq j_2$

Hence $f(w(S_{j_1} \cup S_{j_2}, i)) < f(w(S_{j_1} \cup S_{j_2}, i + 1))$

Theorem (Pumping Lemma 2)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by FA-WA over $(\min, +)$
there exists N s.t. for all (n, N) -representations ($n \geq N$)
exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. given $S_1, \dots, S_N \subseteq \{1 \dots n\}$ (nonempty, different) either:

- exists j s.t. $f(w(S_j, i)) < f(w(S_j, i + 1))$;
- exists $j_1 \neq j_2$ s.t. $f(w(S_{j_1} \cup S_{j_2}, i)) = f(w(S_{j_1} \cup S_{j_2}, i + 1))$.

Example: f – “smallest block of $b's$ ”

Let $w = (\underline{b^N} a^N)^N$ ($n = N$)

Let $S_j = \{1, \dots, N\} \setminus \{j\}$, $f(w(S_j, i)) = N$ for all i, j

But $S_{j_1} \cup S_{j_2} = \{1, \dots, N\}$ for $j_1 \neq j_2$

Hence $f(w(S_{j_1} \cup S_{j_2}, i)) < f(w(S_{j_1} \cup S_{j_2}, i + 1))$

Corollary: FA-WA \subsetneq PA-WA over $(\min, +)$

Polynomially-ambiguous over $(\min, +)$

over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Polynomially-ambiguous over $(\min, +)$

over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Example: “smallest block of b 's”

Polynomially-ambiguous over $(\min, +)$

over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Example: “smallest block of b 's”


Negative example: let $w = w_0 \# w_1 \# \dots \# w_m$, where $w_i \in \{a, b\}^*$
$$f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$$

Polynomially-ambiguous over $(\min, +)$

over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Example: “smallest block of b 's”

Negative example: let $w = w_0 \# w_1 \# \dots \# w_m$, where $w_i \in \{a, b\}^*$
$$f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$$




Polynomially-ambiguous over $(\min, +)$

over $\mathbb{N}_\infty(\min, +, \infty, 0)$

Example: “smallest block of b 's”

Negative example: let $w = w_0 \# w_1 \# \dots \# w_m$, where $w_i \in \{a, b\}^*$
$$f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$$



Number of runs: 2^k

Pumping Lemma 3

First, some notation

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Theorem (Pumping lemma 3)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by PA-WA over $\mathbb{N}_\infty(\min, +)$
there exists N s.t. for all (n, N) -representations

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Theorem (Pumping lemma 3)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by PA-WA over $\mathbb{N}_\infty(\min, +)$

there exists N s.t. for all (n, N) -representations

there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Theorem (Pumping lemma 3)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by PA-WA over $\mathbb{N}_\infty(\min, +)$

there exists N s.t. for all (n, N) -representations

there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Theorem (Pumping lemma 3)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by PA-WA over $\mathbb{N}_\infty(\min, +)$

there exists N s.t. for all (n, N) -representations

there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$

Pumping Lemma 3

First, some notation

Let $S_1, \dots, S_m \subseteq \{1, \dots, n\}$

A partition if $\bigcup_{j=1}^m S_j = \{1, \dots, n\}$, S_j nonempty, $S_{j_1} \cap S_{j_2} = \emptyset$

$S \subseteq \{1, \dots, n\}$ is a selector if $|S \cap S_j| = 1$ for all j

Theorem (Pumping lemma 3)

Let $f : \Sigma^* \rightarrow \mathbb{N}_\infty$ be recognizable by PA-WA over $\mathbb{N}_\infty(\min, +)$

there exists N s.t. for all (n, N) -representations

there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$

s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Example: $f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Example: $f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$

$$w = (\underline{a^N} \underline{b^N} \#)^m$$

$$\{1 \dots n\} = \{(1, 1), (2, 1) \dots (1, m), (2, m)\}$$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Example: $f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$

$$w = (\underline{a^N} \underline{b^N} \#)^m$$

$$\{1 \dots n\} = \{(1, 1), (2, 1) \dots (1, m), (2, m)\}$$

$$S_j = \{(1, j), (2, j)\}, \quad f(w(S_j, i)) < f(w(S_j, i + 1))$$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Example: $f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$

$$w = (\underline{a^N} \underline{b^N} \#)^m$$

$$\{1 \dots n\} = \{(1, 1), (2, 1) \dots (1, m), (2, m)\}$$

$$S_j = \{(1, j), (2, j)\}, \quad f(w(S_j, i)) < f(w(S_j, i + 1))$$

for every selector $f(w(S, i)) = f(w(S, i + 1))$

Theorem (Pumping lemma 3)

(...) there exists N s.t. for all (n, N) -representations
there exists refinement $w = u'_0 \cdot \underline{y_1} \cdot u'_1 \cdot \underline{y_2} \cdot \dots \cdot u'_{n-1} \cdot \underline{y_n} \cdot u'_n$
s.t. for every partition S_1, \dots, S_m of $\{1, \dots, n\}$ either:

- there exists j s.t. $f(w(S_j, i)) = f(w(S_j, i + 1))$
- there exists a selector S s.t. $f(w(S, i)) < f(w(S, i + 1))$

Example: $f(w) = \sum_{i=0}^k \min(\#_a(w_i), \#_b(w_i))$

$$w = (\underline{a^N} \underline{b^N} \#)^m$$

$$\{1 \dots n\} = \{(1, 1), (2, 1) \dots (1, m), (2, m)\}$$

$$S_j = \{(1, j), (2, j)\}, \quad f(w(S_j, i)) < f(w(S_j, i + 1))$$

for every selector $f(w(S, i)) = f(w(S, i + 1))$

Corollary: $\text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$

Conclusions

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)
- Can we hope for characterizations?

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)
- Can we hope for characterizations?

Not for free. For every $f : \Sigma^* \rightarrow \mathbb{N}$ consider $g(w) = f(w) + |w|$

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)
- Can we hope for characterizations?

Not for free. For every $f : \Sigma^* \rightarrow \mathbb{N}$ consider $g(w) = f(w) + |w|$
 f and g should be in the same class

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)
- Can we hope for characterizations?

Not for free. For every $f : \Sigma^* \rightarrow \mathbb{N}$ consider $g(w) = f(w) + |w|$
 f and g should be in the same class
but g satisfies all lemmas.

Conclusions

- Corollary: $\text{U-WA} \subsetneq \text{FA-WA} \subsetneq \text{PA-WA} \subsetneq \text{WA}$ over $(\min, +)$
(only the last one is new)
- Pumping Lemmas 1 and 2 hold for more (WA over $\mathbb{N}(+, \cdot, 0, 1)$)

- Can we hope for characterizations?

Not for free. For every $f : \Sigma^* \rightarrow \mathbb{N}$ consider $g(w) = f(w) + |w|$

f and g should be in the same class

but g satisfies all lemmas.

- Beyond weighted automata

Pumping lemmas for weighted logic and cost-register automata?