

Lecture 2

Linear recursive sequences

Recursive sequences

- **Geometric sequences**

$$q_0 = a, \quad q_{n+1} = q_n \cdot c$$

Recursive sequences

- Geometric sequences

$$q_0 = a, \quad q_{n+1} = q_n \cdot c$$

- Arithmetic sequences

$$u_0 = a, \quad u_{n+1} = u_n + b$$

Recursive sequences

- Geometric sequences

$$q_0 = a, \quad q_{n+1} = q_n \cdot c$$

- Arithmetic sequences

$$u_0 = a, \quad u_{n+1} = u_n + b$$

- Fibonacci sequence F_n

$$f_0 = 0, \quad f_1 = 1, \quad f_n = f_{n-1} + f_{n-2} \quad \text{for } n \geq 2$$

f_0	f_1	f_2	f_3	f_4	f_5	f_6	f_7	f_8	f_9	f_{10}	f_{11}
0	1	1	2	3	5	8	13	21	34	55	89

Linear recursive sequences (LRS)

Fix a semiring $\mathbb{S}(\oplus, \odot, 0, 1)$ (I often omit \odot)

$$u_{n+k} = a_k u_{n+k-1} \oplus a_{k-1} u_{n+k-2} \oplus \dots \oplus a_1 u_n$$

fixing $a_1, \dots, a_k \in \mathbb{S}$ and $u_0, \dots, u_{k-1} \in \mathbb{S}$

Linear recursive sequences (LRS)

Fix a semiring $\mathbb{S}(\oplus, \odot, 0, 1)$ (I often omit \odot)

$$u_{n+k} = a_k u_{n+k-1} \oplus a_{k-1} u_{n+k-2} \oplus \dots \oplus a_1 u_n$$

$$\text{fixing } a_1, \dots, a_k \in \mathbb{S} \quad \text{and } u_0, \dots, u_{k-1} \in \mathbb{S}$$

- Examples for $\mathbb{S}(\oplus, \odot, 0, 1) = (\mathbb{Q}, +, \cdot, 0, 1)$:

Fibonacci sequence F_n $k = 2$, $a_1 = a_2 = 1$, $F_0 = 0$, $F_1 = 1$

Linear recursive sequences (LRS)

Fix a semiring $\mathbb{S}(\oplus, \odot, 0, 1)$ (I often omit \odot)

$$u_{n+k} = a_k u_{n+k-1} \oplus a_{k-1} u_{n+k-2} \oplus \dots \oplus a_1 u_n$$

$$\text{fixing } a_1, \dots, a_k \in \mathbb{S} \quad \text{and } u_0, \dots, u_{k-1} \in \mathbb{S}$$

- Examples for $\mathbb{S}(\oplus, \odot, 0, 1) = (\mathbb{Q}, +, \cdot, 0, 1)$:

Fibonacci sequence F_n $k = 2$, $a_1 = a_2 = 1$, $F_0 = 0$, $F_1 = 1$

Geometric sequences q_n $k = 1$, $a_1 = c$, $q_0 = a$

Linear recursive sequences (LRS)

Fix a semiring $\mathbb{S}(\oplus, \odot, 0, 1)$ (I often omit \odot)

$$u_{n+k} = a_k u_{n+k-1} \oplus a_{k-1} u_{n+k-2} \oplus \dots \oplus a_1 u_n$$

$$\text{fixing } a_1, \dots, a_k \in \mathbb{S} \quad \text{and } u_0, \dots, u_{k-1} \in \mathbb{S}$$

- Examples for $\mathbb{S}(\oplus, \odot, 0, 1) = (\mathbb{Q}, +, \cdot, 0, 1)$:

Fibonacci sequence F_n $k = 2$, $a_1 = a_2 = 1$, $F_0 = 0$, $F_1 = 1$

Geometric sequences q_n $k = 1$, $a_1 = c$, $q_0 = a$

Definition

u_n is (homogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

Linear recursive sequences (LRS)

Fix a semiring $\mathbb{S}(\oplus, \odot, 0, 1)$ (I often omit \odot)

$$u_{n+k} = a_k u_{n+k-1} \oplus a_{k-1} u_{n+k-2} \oplus \dots \oplus a_1 u_n$$

$$\text{fixing } a_1, \dots, a_k \in \mathbb{S} \quad \text{and } u_0, \dots, u_{k-1} \in \mathbb{S}$$

- Examples for $\mathbb{S}(\oplus, \odot, 0, 1) = (\mathbb{Q}, +, \cdot, 0, 1)$:

Fibonacci sequence F_n $k = 2$, $a_1 = a_2 = 1$, $F_0 = 0$, $F_1 = 1$

Geometric sequences q_n $k = 1$, $a_1 = c$, $q_0 = a$

Definition

u_n is (homogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

Fibonacci: $L(x_1, x_2) = x_1 + x_2$

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1x_1 \oplus \dots \oplus a_kx_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

- Example for $(\mathbb{Q}, +, \cdot, 0, 1)$

Arithmetic sequences u_n $k = 1, a_1 = 1, a_0 = b, u_0 = a$

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1x_1 \oplus \dots \oplus a_kx_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

- Example for $(\mathbb{Q}, +, \cdot, 0, 1)$

Arithmetic sequences u_n $k = 1, a_1 = 1, a_0 = b, u_0 = a$

- But notice that $u_{n+1} - u_n = b$ for all n

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

- Example for $(\mathbb{Q}, +, \cdot, 0, 1)$

Arithmetic sequences u_n $k = 1, a_1 = 1, a_0 = b, u_0 = a$

- But notice that $u_{n+1} - u_n = b$ for all n

So $u_{n+2} - u_{n+1} = u_{n+1} - u_n$

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

- Example for $(\mathbb{Q}, +, \cdot, 0, 1)$

Arithmetic sequences u_n $k = 1, a_1 = 1, a_0 = b, u_0 = a$

- But notice that $u_{n+1} - u_n = b$ for all n

So $u_{n+2} - u_{n+1} = u_{n+1} - u_n$

This gives us a homogeneous definition for $k = 2$

$$u_{n+2} = 2u_{n+1} - u_n$$

Nonhomogeneous linear recursive sequences

Definition

u_n is (nonhomogeneous) linear recursive if there is

$$L(x_1, x_2, \dots, x_k) = a_1 x_1 \oplus \dots \oplus a_k x_k \oplus a_0, \quad a_i \in \mathbb{S}$$

s.t. $u_{n+k} = L(u_n, \dots, u_{n+k-1})$ for all n

- Example for $(\mathbb{Q}, +, \cdot, 0, 1)$

Arithmetic sequences u_n $k = 1, a_1 = 1, a_0 = b, u_0 = a$

- But notice that $u_{n+1} - u_n = b$ for all n

So $u_{n+2} - u_{n+1} = u_{n+1} - u_n$

This gives us a homogeneous definition for $k = 2$

$$u_{n+2} = 2u_{n+1} - u_n$$

it's not a coincidence

Linear recursive sequences recursion depth

Number of previous elements referred to (k)

Linear recursive sequences recursion depth

Number of previous elements referred to (k)

Examples: Fibonacci $k = 2$, geometric $k = 1$,
arithmetic $k = 2$ (homogeneous) or $k = 1$ (nonhomogeneous)

Linear recursive sequences recursion depth

Number of previous elements referred to (k)

Examples: Fibonacci $k = 2$, geometric $k = 1$,
arithmetic $k = 2$ (homogeneous) or $k = 1$ (nonhomogeneous)

- Can we restrict to recursion depth 1?

Linear recursive sequences recursion depth

Number of previous elements referred to (k)

Examples: Fibonacci $k = 2$, geometric $k = 1$,
arithmetic $k = 2$ (homogeneous) or $k = 1$ (nonhomogeneous)

- Can we restrict to recursion depth 1?

Not for homogeneous: $u_{n+1} = c \cdot u_n$ are geometric sequences

For nonhomogeneous also no (see tutorials)

Linear recursive sequences recursion depth

Number of previous elements referred to (k)

Examples: Fibonacci $k = 2$, geometric $k = 1$,
arithmetic $k = 2$ (homogeneous) or $k = 1$ (nonhomogeneous)

- Can we restrict to recursion depth 1?

Not for homogeneous: $u_{n+1} = c \cdot u_n$ are geometric sequences

For nonhomogeneous also no (see tutorials)

- Idea: system of linear sequences

Example: Fibonacci F_n with an extra sequence G_n ($= F_{n+1}$)

$$\begin{cases} F_0 = 0 \\ G_0 = 1 \end{cases} \quad \begin{cases} F_{n+1} = G_n \\ G_{n+1} = F_n + G_n \end{cases}$$

System of linear sequences

Definition

A sequence is defined by a system of linear sequences if $u_n = u_n^1$ and

$$\left\{ \begin{array}{l} u_0^1 = c_1 \\ u_0^2 = c_2 \\ \vdots \\ u_0^k = c_k \end{array} \right. \quad \left\{ \begin{array}{l} u_{n+1}^1 = L_1(u_n^1, u_n^2, \dots, u_n^k) \\ u_{n+1}^2 = L_2(u_n^1, u_n^2, \dots, u_n^k) \\ \vdots \\ u_{n+1}^k = L_k(u_n^1, u_n^2, \dots, u_n^k) \end{array} \right.$$

where L_i are linear, $c_i \in \mathbb{S}$

System of linear sequences

Definition

A sequence is defined by a system of linear sequences if $u_n = u_n^1$ and

$$\begin{cases} u_0^1 = c_1 \\ u_0^2 = c_2 \\ \vdots \\ u_0^k = c_k \end{cases} \quad \begin{cases} u_{n+1}^1 = L_1(u_n^1, u_n^2, \dots, u_n^k) \\ u_{n+1}^2 = L_2(u_n^1, u_n^2, \dots, u_n^k) \\ \vdots \\ u_{n+1}^k = L_k(u_n^1, u_n^2, \dots, u_n^k) \end{cases}$$

where L_i are linear, $c_i \in \mathbb{S}$

- This is equivalent to

We have k dimensional vector of sequences: $\vec{u}_n^T = (u_n^1, \dots, u_n^k)$

With a k dimensional vector of initial values $I^T = (c_1, \dots, c_k)$

And a matrix M of dimension $k \times k$ s.t. $\vec{u}_n^T = I^T M^n$

$(M[i, \bullet] = L_i)$

System of linear sequences

Definition

A sequence is defined by a system of linear sequences if $u_n = u_n^1$ and

$$\begin{cases} u_0^1 = c_1 \\ u_0^2 = c_2 \\ \vdots \\ u_0^k = c_k \end{cases} \quad \begin{cases} u_{n+1}^1 = L_1(u_n^1, u_n^2, \dots, u_n^k) \\ u_{n+1}^2 = L_2(u_n^1, u_n^2, \dots, u_n^k) \\ \vdots \\ u_{n+1}^k = L_k(u_n^1, u_n^2, \dots, u_n^k) \end{cases}$$

where L_i are linear, $c_i \in \mathbb{S}$

$$\begin{cases} F_0 = 0 \\ G_0 = 1 \end{cases} \quad \begin{cases} F_{n+1} = G_n \\ G_{n+1} = F_n + G_n \end{cases}$$

$$(F_n, G_n) = (0, 1) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^n$$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example

Consider $a_n = n^2$, recall $(n + 1)^2 = n^2 + 2n + 1$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example

Consider $a_n = n^2$, recall $(n+1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example

Consider $a_n = n^2$, recall $(n+1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example

Consider $a_n = n^2$, recall $(n+1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$

$$(a_n, b_n, c_n) = I^T M^n$$

Example definable by a system of linear equations

Over $(\mathbb{Q}, +, \cdot, 0, 1)$

Example

Consider $a_n = n^2$, recall $(n+1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$

$$(a_n, b_n, c_n) = I^T M^n$$

Can you define it as a linear sequence?

Equivalence of the definitions

Theorem

If \mathbb{S} is a commutative **ring** then

- (1) a_n is homogeneous linear recursive iff
- (2) a_n is nonhomogeneous linear recursive iff
- (3) a_n is definable as a system of linear equations

Equivalence of the definitions

Theorem

If \mathbb{S} is a commutative ring then

- (1) a_n is homogeneous linear recursive iff
- (2) a_n is nonhomogeneous linear recursive iff
- (3) a_n is definable as a system of linear equations

Proof.

- We will prove
 - (1) \implies (2) obvious

Equivalence of the definitions

Theorem

If \mathbb{S} is a commutative **ring** then

- (1) a_n is homogeneous linear recursive iff
- (2) a_n is nonhomogeneous linear recursive iff
- (3) a_n is definable as a system of linear equations

Proof.

- We will prove
 - (1) \implies (2) obvious
 - (2) \implies (3) next slide

Equivalence of the definitions

Theorem

If \mathbb{S} is a commutative **ring** then

- (1) a_n is homogeneous linear recursive iff
- (2) a_n is nonhomogeneous linear recursive iff
- (3) a_n is definable as a system of linear equations

Proof.

- We will prove
 - (1) \implies (2) obvious
 - (2) \implies (3) next slide
 - (3) \implies (1) the hard part

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

- We use k extra sequences $u_n^0, u_n^1, \dots, u_n^k$, where $u_n = u_n^1$

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

- We use k extra sequences $u_n^0, u_n^1, \dots, u_n^k$, where $u_n = u_n^1$
- Define $u_0^0 = a_0$ and $u_{n+1}^0 = u_n^1$

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

- We use k extra sequences $u_n^0, u_n^1, \dots, u_n^k$, where $u_n = u_n^1$
- Define $u_0^0 = a_0$ and $u_{n+1}^0 = u_n^1$
- And $u_0^i = u_{i-1}$ for all $0 < i \leq k$ and $u_{n+1}^i = u_n^{i+1}$ for $0 < i < k$

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

- We use k extra sequences $u_n^0, u_n^1, \dots, u_n^k$, where $u_n = u_n^1$
- Define $u_0^0 = a_0$ and $u_{n+1}^0 = u_n^1$
- And $u_0^i = u_{i-1}$ for all $0 < i \leq k$ and $u_{n+1}^i = u_n^{i+1}$ for $0 < i < k$
- Finally $u_{n+1}^k = a_1u_n^1 + \dots + a_ku_n^k + u_n^0$

From nonhomogeneous to a system

Let $L(x_1, x_2, \dots, x_k) = a_1x_1 + \dots + a_kx_k + a_0$ (semiring $+$)

Such that $u_{n+k} = L(u_n, \dots, u_{n+k-1})$

- We use k extra sequences $u_n^0, u_n^1, \dots, u_n^k$, where $u_n = u_n^1$
- Define $u_0^0 = a_0$ and $u_{n+1}^0 = u_n^1$
- And $u_0^i = u_{i-1}$ for all $0 < i \leq k$ and $u_{n+1}^i = u_n^{i+1}$ for $0 < i < k$
- Finally $u_{n+1}^k = a_1u_n^1 + \dots + a_ku_n^k + u_n^0$

• Example

$$u_{n+3} = 3u_{n+2} - 2u_{n+1} + 4u_n - 5 \rightsquigarrow \begin{cases} x_0 = -5 \\ u_0 = u_0 \\ b_0 = u_1 \\ c_0 = u_2 \\ d_0 = u_3 \end{cases} \begin{cases} x_{n+1} = x_n \\ u_{n+1} = b_n \\ b_{n+1} = c_n \\ c_{n+1} = d_n \\ d_{n+1} = 3u_n - 2b_n + 4c_n + x_n \end{cases}$$

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

and $u_n = u_n^1$

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

$$\text{and } u_n = u_n^1$$

Notice that $u_n = I^\top M^n F$ for $F^\top = (1, 0, \dots, 0)$

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

$$\text{and } u_n = u_n^1$$

Notice that $u_n = I^\top M^n F$ for $F^\top = (1, 0, \dots, 0)$

- We fix the first k elements as u_0^1, \dots, u_{k-1}^1

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

$$\text{and } u_n = u_n^1$$

Notice that $u_n = I^\top M^n F$ for $F^\top = (1, 0, \dots, 0)$

- We fix the first k elements as u_0^1, \dots, u_{k-1}^1
- Let $\det(\lambda \text{Id} - M) = p(\lambda) = a_k \lambda^k + \dots + a_1 \lambda^1 + a_0$

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

$$\text{and } u_n = u_n^1$$

Notice that $u_n = I^\top M^n F$ for $F^\top = (1, 0, \dots, 0)$

- We fix the first k elements as u_0^1, \dots, u_{k-1}^1
- Let $\det(\lambda \text{Id} - M) = p(\lambda) = a_k \lambda^k + \dots + a_1 \lambda^1 + a_0$

$p(\lambda)$ is the characteristic polynomial of M so:

$$a_k = 1$$

From systems to homogeneous linear recursive sequences

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

$$\text{and } u_n = u_n^1$$

Notice that $u_n = I^\top M^n F$ for $F^\top = (1, 0, \dots, 0)$

- We fix the first k elements as u_0^1, \dots, u_{k-1}^1
- Let $\det(\lambda \text{Id} - M) = p(\lambda) = a_k \lambda^k + \dots + a_1 \lambda^1 + a_0$

$p(\lambda)$ is the characteristic polynomial of M so:

$$a_k = 1$$

$$p(A) = 0 \text{ (Cayley–Hamilton theorem)}$$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

From systems to homogeneous linear recursive sequences

- $u_n = I^\top M^n F$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

$$a_k = 1$$

From systems to homogeneous linear recursive sequences

- $u_n = I^T M^n F$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

$$a_k = 1$$

- $M^{n+k} = \sum_{i=0}^{k-1} -a_i M^{n+i}$

From systems to homogeneous linear recursive sequences

- $u_n = I^T M^n F$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

$$a_k = 1$$

- $M^{n+k} = \sum_{i=0}^{k-1} -a_i M^{n+i}$

$$u_{n+k} = I^T M^{n+k} F = I^T \left(\sum_{i=0}^{k-1} -a_i M^{n+i} \right) F$$

From systems to homogeneous linear recursive sequences

- $u_n = I^T M^n F$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

$$a_k = 1$$

- $M^{n+k} = \sum_{i=0}^{k-1} -a_i M^{n+i}$

$$u_{n+k} = I^T M^{n+k} F = I^T \left(\sum_{i=0}^{k-1} -a_i M^{n+i} \right) F$$

By linearity:

$$u_{n+k} = \sum_{i=0}^{k-1} -a_i (I^T M^{n+i} F) = \sum_{i=0}^{k-1} -a_i u_{n+i}$$

From systems to homogeneous linear recursive sequences

- $u_n = I^T M^n F$

$$a_k M^k + \dots + a_1 M^1 + a_0 I = 0$$

$$a_k = 1$$

- $M^{n+k} = \sum_{i=0}^{k-1} -a_i M^{n+i}$

$$u_{n+k} = I^T M^{n+k} F = I^T \left(\sum_{i=0}^{k-1} -a_i M^{n+i} \right) F$$

By linearity:

$$u_{n+k} = \sum_{i=0}^{k-1} -a_i (I^T M^{n+i} F) = \sum_{i=0}^{k-1} -a_i u_{n+i}$$

This defines u_n as a homogeneous linear recursive sequence



Example

Example

Consider $a_n = n^2$, recall $(n + 1)^2 = n^2 + 2n + 1$

$$\left\{ \begin{array}{l} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{array} \right. \quad \left\{ \begin{array}{l} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{array} \right.$$

Example

Example

Consider $a_n = n^2$, recall $(n + 1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix} \quad \det(\lambda \text{Id} - M) = \lambda^3 - 3\lambda^2 - 3\lambda - 1$$

Example

Example

Consider $a_n = n^2$, recall $(n + 1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix} \quad \det(\lambda \text{Id} - M) = \lambda^3 - 3\lambda^2 - 3\lambda - 1$$

Then $a_{n+3} = 3a_{n+2} - 3a_{n+1} + a_n$

Example

Example

Consider $a_n = n^2$, recall $(n + 1)^2 = n^2 + 2n + 1$

$$\begin{cases} a_0 = 0 \\ b_0 = 0 \\ c_0 = 1 \end{cases} \quad \begin{cases} a_{n+1} = a_n + 2b_n + c_n \\ b_{n+1} = b_n + c_n \\ c_{n+1} = c_n \end{cases}$$

$$I = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad M = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix} \quad \det(\lambda \text{Id} - M) = \lambda^3 - 3\lambda^2 - 3\lambda - 1$$

Then $a_{n+3} = 3a_{n+2} - 3a_{n+1} + a_n$

Verify that $(n + 3)^2 = 3(n + 2)^2 - 3(n + 1)^2 + n^2$

Comments

- These worked for any semiring:

$$(1) \implies (2)$$

$$(2) \implies (3)$$

Comments

- These worked for any semiring:
 $(1) \implies (2)$
 $(2) \implies (3)$
- $(3) \implies (1)$ this required the Cayley–Hamilton theorem
 which is true for commutative semirings

Comments

- These worked for any semiring:

$$(1) \implies (2)$$

$$(2) \implies (3)$$

- $(3) \implies (1)$ this required the Cayley–Hamilton theorem
which is true for commutative semirings

- The most general class:

The class of sequences defined by systems of linear sequences

When I refer to linear sequences I will refer to sequences definable by a system

Comments

- These worked for any semiring:

$$(1) \implies (2)$$

$$(2) \implies (3)$$

- $(3) \implies (1)$ this required the Cayley–Hamilton theorem which is true for commutative semirings

- The most general class:

The class of sequences defined by systems of linear sequences

When I refer to linear sequences I will refer to sequences definable by a system

- On tutorials: for some semirings that are not rings the inclusions:

$$(1) \subseteq (2) \subseteq (3) \text{ are strict}$$

Connections with weighted automata

Recall: $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \mathbb{S}$

$$\llbracket \mathcal{A} \rrbracket (a_1 a_2 \dots a_n) = I^\top M_{a_1} M_{a_2} \dots M_{a_n} F$$

Connections with weighted automata

Recall: $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \mathbb{S}$

$$\llbracket \mathcal{A} \rrbracket (a_1 a_2 \dots a_n) = I^\top M_{a_1} M_{a_2} \dots M_{a_n} F$$

- Assume that $\Sigma = \{a\}$, i.e. $|\Sigma| = 1$

Then $\Sigma^* = \{\epsilon, a, a^2, \dots\} \equiv \mathbb{N}$

(identify every word with its length)

Connections with weighted automata

Recall: $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \mathbb{S}$

$$\llbracket \mathcal{A} \rrbracket (a_1 a_2 \dots a_n) = I^\top M_{a_1} M_{a_2} \dots M_{a_n} F$$

- Assume that $\Sigma = \{a\}$, i.e. $|\Sigma| = 1$

Then $\Sigma^* = \{\epsilon, a, a^2, \dots\} \equiv \mathbb{N}$

(identify every word with its length)

$\llbracket \mathcal{A} \rrbracket : \mathbb{N} \rightarrow \mathbb{S}$, they define sequences

$$\llbracket \mathcal{A} \rrbracket (n) = I^\top M^n F$$

Connections with weighted automata

Recall: $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \mathbb{S}$

$$\llbracket \mathcal{A} \rrbracket (a_1 a_2 \dots a_n) = I^\top M_{a_1} M_{a_2} \dots M_{a_n} F$$

- Assume that $\Sigma = \{a\}$, i.e. $|\Sigma| = 1$

Then $\Sigma^* = \{\epsilon, a, a^2, \dots\} \equiv \mathbb{N}$

(identify every word with its length)

$\llbracket \mathcal{A} \rrbracket : \mathbb{N} \rightarrow \mathbb{S}$, they define sequences

$$\llbracket \mathcal{A} \rrbracket (n) = I^\top M^n F$$

- For $F = (1, 0, \dots, 0)$ these are sequences definable by systems
On tutorials you'll see that for any F we are still in the same class

From sequences to weighted automata

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

and $u_n = u_n^1$

From sequences to weighted automata

- Suppose $\vec{u}_n^\top = I^\top M^n$
where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional
 $I^\top = (u_0^1, \dots, u_0^k)$
and $u_n = u_n^1$
- Set $\mathcal{A} = (Q, \Sigma, T, I, F)$
 $Q = \{1, \dots, k\}$, $I^\top = (u_0^1, \dots, u_0^k)$, $F^\top = (1, 0, \dots, 0)$
 $T = \{(p, a, s, q) \mid M_a[p, q] = s\}$

From sequences to weighted automata

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

and $u_n = u_n^1$

- Set $\mathcal{A} = (Q, \Sigma, T, I, F)$


$$Q = \{1, \dots, k\}, I^\top = (u_0^1, \dots, u_0^k), F^\top = (1, 0, \dots, 0)$$

$$T = \{(p, a, s, q) \mid M_a[p, q] = s\}$$

- Example

Fibonacci over $(\mathbb{Q}, +, \cdot, 0, 1)$

$$(F_n, G_n) = (0, 1) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^n$$



From sequences to weighted automata

- Suppose $\vec{u}_n^\top = I^\top M^n$

where $\vec{u}_n^\top = (u_n^1, \dots, u_n^k)$ and M is $k \times k$ dimensional

$$I^\top = (u_0^1, \dots, u_0^k)$$

and $u_n = u_n^1$

- Set $\mathcal{A} = (Q, \Sigma, T, I, F)$


$$Q = \{1, \dots, k\}, I^\top = (u_0^1, \dots, u_0^k), F^\top = (1, 0, \dots, 0)$$

$$T = \{(p, a, s, q) \mid M_a[p, q] = s\}$$

- Example

Fibonacci over $(\mathbb{Q}, +, \cdot, 0, 1)$

$$(F_n, G_n) = (0, 1) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^n$$



- When weights are 1 then it is equivalent to counting the accepting runs

Closed form

Fix $(\mathbb{Q}, +, \cdot, 0, 1)$

- $u_n = I^T M^n F$ characteristic polynomial $p(x) = (x - \lambda_1)^{m_1} \dots (x - \lambda_s)^{m_s}$

Closed form

Fix $(\mathbb{Q}, +, \cdot, 0, 1)$

- $u_n = I^T M^n F$ characteristic polynomial $p(x) = (x - \lambda_1)^{m_1} \dots (x - \lambda_s)^{m_s}$

$$M^n = B J^n B^{-1}, \quad J^n = \begin{bmatrix} J_{m_1}^n(\lambda_1) & 0 & 0 & \dots & 0 \\ 0 & J_{m_2}^n(\lambda_2) & 0 & \dots & 0 \\ \vdots & \dots & \ddots & \dots & \vdots \\ 0 & \dots & 0 & J_{m_{s-1}}^n(\lambda_{s-1}) & 0 \\ 0 & \dots & \dots & 0 & J_{m_s}^n(\lambda_s) \end{bmatrix}$$

$$J_{m_i}^n(\lambda_i) = \begin{bmatrix} \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} & \binom{n}{2} \lambda_i^{n-2} & \dots & \binom{n}{m_i-1} \lambda_i^{n-m_i+1} \\ 0 & \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} & \dots & \binom{n}{m_i-2} \lambda_i^{n-m_i+2} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} \\ 0 & 0 & \dots & 0 & \lambda_i^n \end{bmatrix},$$

Closed form

Fix $(\mathbb{Q}, +, \cdot, 0, 1)$

- $u_n = I^\top M^n F$ characteristic polynomial $p(x) = (x - \lambda_1)^{m_1} \dots (x - \lambda_s)^{m_s}$

$$M^n = BJ^nB^{-1}, \quad J^n = \begin{bmatrix} J_{m_1}^n(\lambda_1) & 0 & 0 & \dots & 0 \\ 0 & J_{m_2}^n(\lambda_2) & 0 & \dots & 0 \\ \vdots & \dots & \ddots & \dots & \vdots \\ 0 & \dots & 0 & J_{m_{s-1}}^n(\lambda_{s-1}) & 0 \\ 0 & \dots & \dots & 0 & J_{m_s}^n(\lambda_s) \end{bmatrix}$$

$$J_{m_i}^n(\lambda_i) = \begin{bmatrix} \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} & \binom{n}{2} \lambda_i^{n-2} & \dots & \binom{n}{m_i-1} \lambda_i^{n-m_i+1} \\ 0 & \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} & \dots & \binom{n}{m_i-2} \lambda_i^{n-m_i+2} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_i^n & \binom{n}{1} \lambda_i^{n-1} \\ 0 & 0 & \dots & 0 & \lambda_i^n \end{bmatrix},$$

Note $\lambda_i \in \mathbb{C}$

Closed form continued

- $u_n = I^T M^n F = \sum_{i=1}^s p_i(n) \lambda_i^n$

Where $\lambda_i \in \mathbb{C}$ roots of the characteristic polynomial

p_i are polynomials over \mathbb{C}

Closed form continued

- $u_n = I^T M^n F = \sum_{i=1}^s p_i(n) \lambda_i^n$

Where $\lambda_i \in \mathbb{C}$ roots of the characteristic polynomial

p_i are polynomials over \mathbb{C}

- For example for Fibonacci

Characteristic polynomial is $\det \left(x \text{Id} - \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \right) = x^2 - x + 1$

$$= \left(x - \frac{1+\sqrt{5}}{2} \right) \left(x - \frac{1-\sqrt{5}}{2} \right)$$

Closed form continued

- $u_n = I^T M^n F = \sum_{i=1}^s p_i(n) \lambda_i^n$

Where $\lambda_i \in \mathbb{C}$ roots of the characteristic polynomial

p_i are polynomials over \mathbb{C}

- For example for Fibonacci

Characteristic polynomial is $\det \left(x \text{Id} - \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \right) = x^2 - x + 1$

$$= \left(x - \frac{1+\sqrt{5}}{2} \right) \left(x - \frac{1-\sqrt{5}}{2} \right)$$

- One can verify that $F_n = \frac{1}{\sqrt{5}} \frac{1+\sqrt{5}}{2}^n - \frac{1}{\sqrt{5}} \frac{1-\sqrt{5}}{2}^n$

Asymptotics

Lemma

If u_n is a linear recursive sequence over $(\mathbb{Q}, +, \cdot, 0, 1)$

then $|u_n| \leq c^n$ for some $c \in \mathbb{Q}$

Asymptotics

Lemma

If u_n is a linear recursive sequence over $(\mathbb{Q}, +, \cdot, 0, 1)$
then $|u_n| \leq c^n$ for some $c \in \mathbb{Q}$

Corollary

The sequence $u_n = n!$ cannot be defined by weighted automata over $(\mathbb{Q}, +, \cdot, 0, 1)$

Asymptotics

Lemma

If u_n is a linear recursive sequence over $(\mathbb{Q}, +, \cdot, 0, 1)$
then $|u_n| \leq c^n$ for some $c \in \mathbb{Q}$

Corollary

The sequence $u_n = n!$ cannot be defined by weighted automata over $(\mathbb{Q}, +, \cdot, 0, 1)$

Proof.

$n! > \left(\frac{n}{e}\right)^n$ for n big enough (by Stirling)

Asymptotics

Lemma

If u_n is a linear recursive sequence over $(\mathbb{Q}, +, \cdot, 0, 1)$
then $|u_n| \leq c^n$ for some $c \in \mathbb{Q}$

Corollary

The sequence $u_n = n!$ cannot be defined by weighted automata over $(\mathbb{Q}, +, \cdot, 0, 1)$

Proof.

$n! > \left(\frac{n}{e}\right)^n$ for n big enough (by Stirling)

$$\left(\frac{n}{e}\right)^n = 2^{n(\log(n) - \log(e))}$$

