

When are Emptiness and Containment Decidable for Probabilistic Automata?

Laure Daviaud¹ Marcin Jurdziński² Ranko Lazić²
Filip Mazowiecki³ Guillermo A. Pérez⁴ James Worrell⁵

¹City University of London

²University of Warwick

³Université de Bordeaux

⁴University of Antwerp

⁵University of Oxford

Kaiserslautern May 2019

When are Emptiness and Containment Decidable for Probabilistic Automata?

Laure Daviaud¹ Marcin Jurdziński² Ranko Lazić²
Filip Mazowiecki³ Guillermo A. Pérez⁴ James Worrell⁵

¹City University of London

²University of Warwick

³Université de Bordeaux

⁴University of Antwerp


⁵University of Oxford

Kaiserslautern May 2019

Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}




Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}


\mathcal{A} accepts $\Sigma^*1\Sigma^*$



Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}




\mathcal{A} accepts $\Sigma^*1\Sigma^*$

In general $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{\text{yes}, \text{no}\}$

Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}



\mathcal{A} accepts $\Sigma^*1\Sigma^*$

In general $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{\text{yes}, \text{no}\}$


Probabilistic automata

$\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \text{probabilities}$

Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}



\mathcal{A} accepts $\Sigma^*1\Sigma^*$

In general $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{\text{yes}, \text{no}\}$


Probabilistic automata

$\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \text{probabilities}$, probabilities = $\mathbb{Q} \cap [0, 1]$

Automata

$$\Sigma = \{0, 1\}$$

\mathcal{A}



\mathcal{A} accepts $\Sigma^*1\Sigma^*$

In general $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{\text{yes}, \text{no}\}$

Probabilistic automata


$\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \text{probabilities}$, probabilities = $\mathbb{Q} \cap [0, 1]$

artificial intelligence, verification of probabilistic systems. . .

Probabilistic automata (PA)

$$\Sigma = \{0, 1\}$$


\mathcal{A}



Probabilistic automata (PA)


$$\Sigma = \{0, 1\}$$

\mathcal{A}



Probabilistic automata (PA)

$$\Sigma = \{0, 1\}$$




Given state (●) and letter (0)

transitions induce a distribution $(\frac{1}{2}, \frac{1}{2})$

Probabilistic automata (PA)

$$\Sigma = \{0, 1\}$$




Given state (●) and letter (0)
transitions induce a distribution $(\frac{1}{2}, \frac{1}{2})$

Semantics: for every accepting run multiply probabilities
and sum over all accepting runs

Probabilistic automata (PA)

$$\Sigma = \{0, 1\}$$



Given state (●) and letter (0)


transitions induce a distribution $(\frac{1}{2}, \frac{1}{2})$

Semantics: for every accepting run multiply probabilities
and sum over all accepting runs

E.g. $\llbracket \mathcal{A} \rrbracket(1010) = 2^{-1} + 2^{-3}$

Probabilistic automata (PA)

$$\Sigma = \{0, 1\}$$



Given state (●) and letter (0)
 transitions induce a distribution $(\frac{1}{2}, \frac{1}{2})$

Semantics: for every accepting run multiply probabilities
 and sum over all accepting runs

E.g. $\llbracket \mathcal{A} \rrbracket(1010) = 2^{-1} + 2^{-3}$, $\llbracket \mathcal{A} \rrbracket(w) = \text{bin}(0.w) = \sum_{i|w(i)=1} 2^{-i}$

Decision problems

What do we study?

Decision problems

What do we study?

Emptiness problem

GIVEN: a PA automaton \mathcal{A}

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \frac{1}{2}$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ in short)

Decision problems

What do we study?

ϵ -gap emptiness problem

GIVEN: ϵ , a PA automaton \mathcal{A} s.t. either $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ or $\llbracket \mathcal{A} \rrbracket(w) > \frac{1}{2} + \epsilon$

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \frac{1}{2}$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ and ϵ - $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ in short)

Decision problems

What do we study?

ϵ -gap emptiness problem

GIVEN: ϵ , a PA automaton \mathcal{A} s.t. either $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ or $\llbracket \mathcal{A} \rrbracket(w) > \frac{1}{2} + \epsilon$

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \frac{1}{2}$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ and ϵ - $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ in short)

Containment problem

GIVEN: two PA automata \mathcal{A}, \mathcal{B}

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \llbracket \mathcal{B} \rrbracket(w)$ hold for all w ?

Decision problems

What do we study?

ϵ -gap emptiness problem

GIVEN: ϵ , a PA automaton \mathcal{A} s.t. either $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ or $\llbracket \mathcal{A} \rrbracket(w) > \frac{1}{2} + \epsilon$

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \frac{1}{2}$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ and ϵ - $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ in short)

Containment problem

GIVEN: two PA automata \mathcal{A}, \mathcal{B}

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \llbracket \mathcal{B} \rrbracket(w)$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \llbracket \mathcal{B} \rrbracket$ in short)

Decision problems

What do we study?

ϵ -gap emptiness problem

GIVEN: ϵ , a PA automaton \mathcal{A} s.t. either $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ or $\llbracket \mathcal{A} \rrbracket(w) > \frac{1}{2} + \epsilon$

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \frac{1}{2}$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ and ϵ - $\llbracket \mathcal{A} \rrbracket \leq \frac{1}{2}$ in short)

Containment problem

GIVEN: two PA automata \mathcal{A}, \mathcal{B}

DECIDE: does $\llbracket \mathcal{A} \rrbracket(w) \leq \llbracket \mathcal{B} \rrbracket(w)$ hold for all w ?

($\llbracket \mathcal{A} \rrbracket \leq \llbracket \mathcal{B} \rrbracket$ in short)

Remark

Emptiness reduces to containment (define $\llbracket \mathcal{B} \rrbracket = \frac{1}{2}$)

State of art

tl;dr: it's all almost always undecidable

State of art

tl;dr: it's all almost always undecidable

See recent SIGLOG news article:

“Undecidability results for probabilistic automata” [Fijalkow, 2017]

State of art

tl;dr: it's all almost always undecidable

See recent SIGLOG news article:

“Undecidability results for probabilistic automata” [Fijalkow, 2017]

- Containment and emptiness are undecidable [Paz, 1971]

State of art

tl;dr: it's all almost always undecidable

See recent SIGLOG news article:

“Undecidability results for probabilistic automata” [Fijalkow, 2017]

- Containment and emptiness are undecidable [Paz, 1971]
- ϵ -gap emptiness is undecidable [Condon and Lipton, 1989]

State of art

tl;dr: it's all almost always undecidable

See recent SIGLOG news article:

“Undecidability results for probabilistic automata” [Fijalkow, 2017]

- Containment and emptiness are undecidable [Paz, 1971]
- ϵ -gap emptiness is undecidable [Condon and Lipton, 1989]
- But equivalence is in PTIME [Schützenberger, 1961]

State of art

tl;dr: it's all almost always undecidable

See recent SIGLOG news article:

“Undecidability results for probabilistic automata” [Fijalkow, 2017]

- Containment and emptiness are undecidable [Paz, 1971]
- ϵ -gap emptiness is undecidable [Condon and Lipton, 1989]
- But equivalence is in PTIME [Schützenberger, 1961]
- Many subclasses with some decidability results:
hierarchical, leaktight, bounded ambiguity

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)

- polynomial in $|w|$ (the first example – number of 1's in w)

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)

- polynomial in $|w|$ (the first example – number of 1's in w)

polynomially ambiguous PA

linearly ambiguous PA

...

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)


- polynomial in $|w|$ (the first example – number of 1's in w)

polynomially ambiguous PA

linearly ambiguous PA

...

- constant



$$\llbracket \mathcal{B} \rrbracket (0^i 1 w) = 2^{-i}$$

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)

- polynomial in $|w|$ (the first example – number of 1's in w)

polynomially ambiguous PA

linearly ambiguous PA


...

- constant

finitely ambiguous

k -ambiguous (unambiguous if $k = 1$)

\mathcal{B}



$$\llbracket \mathcal{B} \rrbracket(0^i 1 w) = 2^{-i}$$

Bounded ambiguity

Number of accepting runs in $|w|$? (possibly exponential)

- polynomial in $|w|$ (the first example – number of 1's in w)

polynomially ambiguous PA

linearly ambiguous PA


...

- constant

finitely ambiguous

k -ambiguous (unambiguous if $k = 1$)

\mathcal{B}



$$\llbracket \mathcal{B} \rrbracket(0^i 1 w) = 2^{-i}$$

Remark

Nothing between finitely ambiguous and linearly ambiguous PA

Summary for bounded ambiguous PA

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
1. Emptiness undecidable for quadratically ambiguous PA
(hence containment too)

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA
(hence containment too)
 2. Emptiness decidable for finitely ambiguous PA

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA (hence containment too)
 2. Emptiness decidable for finitely ambiguous PA
- Our results:

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA (hence containment too)
 2. Emptiness decidable for finitely ambiguous PA
- Our results:
 1. Emptiness and containment undecidable for linearly ambiguous PA

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA
(hence containment too)
 2. Emptiness decidable for finitely ambiguous PA
- Our results:
 1. Emptiness and containment undecidable for linearly ambiguous PA
 2. Containment decidable for finitely ambiguous PA
(provided one automaton is unambiguous + Schanuel's conjecture)

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA
(hence containment too)
 2. Emptiness decidable for finitely ambiguous PA
- Our results:
 1. Emptiness and containment undecidable for linearly ambiguous PA
 2. Containment decidable for finitely ambiguous PA
(provided one automaton is unambiguous + Schanuel's conjecture)
 3. ϵ -gap emptiness decidable for polynomially ambiguous PA
(only in the submitted journal version)

Summary for bounded ambiguous PA

- Introduced in [Fijalkow, Riveros and Worrell; 2017]
 1. Emptiness undecidable for quadratically ambiguous PA
(hence containment too)
 2. Emptiness decidable for finitely ambiguous PA
- Our results:
 1. Emptiness and containment undecidable for linearly ambiguous PA
 2. Containment decidable for finitely ambiguous PA
(provided one automaton is unambiguous + Schanuel's conjecture)
 3. ϵ -gap emptiness decidable for polynomially ambiguous PA
(only in the submitted journal version)

rest of the talk

ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

(for finitely ambiguous even emptiness is decidable)


ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

(for finitely ambiguous even emptiness is decidable)

Example:




ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

(for finitely ambiguous even emptiness is decidable)

Example:




ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

(for finitely ambiguous even emptiness is decidable)

Example:




ϵ -gap emptiness problem for polynomially ambiguous PA

INPUT: ϵ , polynomially ambiguous PA \mathcal{A}

Reduce \mathcal{A} to a finitely ambiguous \mathcal{A}'

(for finitely ambiguous even emptiness is decidable)

Example:



The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

We need m extra copies so that $\mathcal{A}(w) \leq \mathcal{A}'(w) + \epsilon$

The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

We need m extra copies so that $\mathcal{A}(w) \leq \mathcal{A}'(w) + \epsilon$

Probability of an ignored run is at most α^n ($0 < \alpha < 1$, $n > m$)

The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

We need m extra copies so that $\mathcal{A}(w) \leq \mathcal{A}'(w) + \epsilon$

Probability of an ignored run is at most α^n ($0 < \alpha < 1$, $n > m$)

in total $\sum_{n>m} P(n) \cdot \alpha^n$

The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

We need m extra copies so that $\mathcal{A}(w) \leq \mathcal{A}'(w) + \epsilon$

Probability of an ignored run is at most α^n ($0 < \alpha < 1$, $n > m$)

in total $\sum_{n>m} P(n) \cdot \alpha^n$

if P is polynomial $\implies \sum_{n>m} P(n) \cdot \alpha^n < \epsilon$ (for m big enough)

The reduction

Clearly $\mathcal{A}'(w) \leq \mathcal{A}(w)$

We need m extra copies so that $\mathcal{A}(w) \leq \mathcal{A}'(w) + \epsilon$

Probability of an ignored run is at most α^n ($0 < \alpha < 1$, $n > m$)

in total $\sum_{n>m} P(n) \cdot \alpha^n$

if P is polynomial $\implies \sum_{n>m} P(n) \cdot \alpha^n < \epsilon$ (for m big enough)

P is polynomial because of polynomial ambiguity


Containment of finitely ambiguous PA

$$\mathcal{A} \leq \mathcal{B} \quad (\mathcal{A}(w) \leq \mathcal{B}(w) \text{ for all } w)$$


Containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$? ($\mathcal{A}(w) \leq \mathcal{B}(w)$ for all w)

Example for $0 < p < 1$




\mathcal{B}




Containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$? ($\mathcal{A}(w) \leq \mathcal{B}(w)$ for all w)

Example for $0 < p < 1$



\mathcal{B}




- Only words of the form a^*ba^* have non-zero value


Containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$? ($\mathcal{A}(w) \leq \mathcal{B}(w)$ for all w)

Example for $0 < p < 1$



\mathcal{B}




- Only words of the form a^*ba^* have non-zero value
- Suffices to solve: does for all $x_1, x_2 \in \mathbb{N}$ hold

$$\left(\frac{1}{6}\right)^{x_1} \left(\frac{1}{6}\right)^{x_2} \leq p \left(\frac{1}{12}\right)^{x_1} \left(\frac{1}{2}\right)^{x_2} + (1 - p) \left(\frac{1}{3}\right)^{x_1} \left(\frac{1}{18}\right)^{x_2}$$


Containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$? ($\mathcal{A}(w) \leq \mathcal{B}(w)$ for all w)

Example for $0 < p < 1$



\mathcal{B}



- Only words of the form a^*ba^* have non-zero value
- Suffices to solve: does for all $x_1, x_2 \in \mathbb{N}$ hold

$$\left(\frac{1}{6}\right)^{x_1} \left(\frac{1}{6}\right)^{x_2} \leq p \left(\frac{1}{12}\right)^{x_1} \left(\frac{1}{2}\right)^{x_2} + (1 - p) \left(\frac{1}{3}\right)^{x_1} \left(\frac{1}{18}\right)^{x_2}$$

$$1 \leq p \left(\frac{1}{2}\right)^{x_1} (3)^{x_2} + (1 - p) (2)^{x_1} \left(\frac{1}{3}\right)^{x_2}$$

Reformulating containment of finitely ambiguous PA

$$\mathcal{A} \leq \mathcal{B}?$$

Reformulating containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$?

Exponential inequalities

GIVEN: $k, l, n > 0$, and vectors $\mathbf{p} \in \mathbb{Q}_{>0}^k$ and $\mathbf{q}_1, \dots, \mathbf{q}_k \in \mathbb{Q}_{>0}^n$

$\mathbf{r} \in \mathbb{Q}_{>0}^l$ and $\mathbf{s}_1, \dots, \mathbf{s}_l \in \mathbb{Q}_{>0}^n$

DECIDE: for every $x_1, \dots, x_n \in \mathbb{N}$ does it hold

$$\sum_{i=1}^k p_i \cdot q_{i,1}^{x_1} \cdot \dots \cdot q_{i,n}^{x_n} \leq \sum_{i=1}^l r_i \cdot s_{i,1}^{x_1} \cdot \dots \cdot s_{i,n}^{x_n}$$

Reformulating containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$?

Exponential inequalities

GIVEN: $k, l, n > 0$, and vectors $\mathbf{p} \in \mathbb{Q}_{>0}^k$ and $\mathbf{q}_1, \dots, \mathbf{q}_k \in \mathbb{Q}_{>0}^n$
 $\mathbf{r} \in \mathbb{Q}_{>0}^l$ and $\mathbf{s}_1, \dots, \mathbf{s}_l \in \mathbb{Q}_{>0}^n$

DECIDE: for every $x_1, \dots, x_n \in \mathbb{N}$ does it hold

$$\sum_{i=1}^k p_i \cdot q_{i,1}^{x_1} \cdot \dots \cdot q_{i,n}^{x_n} \leq \sum_{i=1}^l r_i \cdot s_{i,1}^{x_1} \cdot \dots \cdot s_{i,n}^{x_n}$$

Proposition

Containment of finitely ambiguous PA and exponential inequalities
are effectively equi-decidable

Reformulating containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$?

Exponential inequalities

GIVEN: $k, l, n > 0$, and vectors $\mathbf{p} \in \mathbb{Q}_{>0}^k$ and $\mathbf{q}_1, \dots, \mathbf{q}_k \in \mathbb{Q}_{>0}^n$
 $\mathbf{r} \in \mathbb{Q}_{>0}^l$ and $\mathbf{s}_1, \dots, \mathbf{s}_l \in \mathbb{Q}_{>0}^n$

DECIDE: for every $x_1, \dots, x_n \in \mathbb{N}$ does it hold

$$\sum_{i=1}^k p_i \cdot q_{i,1}^{x_1} \cdot \dots \cdot q_{i,n}^{x_n} \leq \sum_{i=1}^l r_i \cdot s_{i,1}^{x_1} \cdot \dots \cdot s_{i,n}^{x_n}$$

Proposition

Containment of finitely ambiguous PA and exponential inequalities are effectively equi-decidable

- k, l ambiguity of \mathcal{A} and \mathcal{B}

Reformulating containment of finitely ambiguous PA

$\mathcal{A} \leq \mathcal{B}$?

Exponential inequalities

GIVEN: $k, l, n > 0$, and vectors $\mathbf{p} \in \mathbb{Q}_{>0}^k$ and $\mathbf{q}_1, \dots, \mathbf{q}_k \in \mathbb{Q}_{>0}^n$
 $\mathbf{r} \in \mathbb{Q}_{>0}^l$ and $\mathbf{s}_1, \dots, \mathbf{s}_l \in \mathbb{Q}_{>0}^n$

DECIDE: for every $x_1, \dots, x_n \in \mathbb{N}$ does it hold

$$\sum_{i=1}^k p_i \cdot q_{i,1}^{x_1} \cdot \dots \cdot q_{i,n}^{x_n} \leq \sum_{i=1}^l r_i \cdot s_{i,1}^{x_1} \cdot \dots \cdot s_{i,n}^{x_n}$$

Proposition

Containment of finitely ambiguous PA and exponential inequalities are effectively equi-decidable

- k, l ambiguity of \mathcal{A} and \mathcal{B}
- simple cycle decomposition

Exponential inequalities

for every $x_1, \dots, x_n \in \mathbb{N}$ does it hold

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} \leq \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \cdots s_{i,n}^{x_n}$$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \cdots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$

Two cases:

- there exists i, j such that $q_{i,j} > 1$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$

Two cases:

- there exists i, j such that $q_{i,j} > 1$

take x_j very big and $x_s = 0$ for $s \neq j$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$

Two cases:

- there exists i, j such that $q_{i,j} > 1$

take x_j very big and $x_s = 0$ for $s \neq j$

- all $q_{i,j} \leq 1$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$

Two cases:

- there exists i, j such that $q_{i,j} > 1$

take x_j very big and $x_s = 0$ for $s \neq j$

- all $q_{i,j} \leq 1$

best chance for $x_1 = \dots = x_n = 0$

Exponential inequalities

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > \sum_{i=1}^l r_i s_{i,1}^{x_1} \dots s_{i,n}^{x_n}$$

We consider only for $k = 1$ or $l = 1$

If $l = 1$ it boils down to

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} > 1$$


Two cases:

- there exists i, j such that $q_{i,j} > 1$

take x_j very big and $x_s = 0$ for $s \neq j$

- all $q_{i,j} \leq 1$

best chance for $x_1 = \dots = x_n = 0$



decidable
(without
Shanuel)

Seriously though

What if $k = 1$?

Seriously though

What if $k = 1$?

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} < 1$$

Seriously though

What if $k = 1$?

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} < 1$$

Example for $k = n = 2$ and fixed $0 < p < 1$

Seriously though

What if $k = 1$?

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} < 1$$

Example for $k = n = 2$ and fixed $0 < p < 1$

Do there exist $x_1, x_2 \in \mathbb{N}$ such that

$$p \cdot \left(\frac{1}{2}\right)^{x_1} \cdot 3^{x_2} + (1 - p) \cdot 2^{x_1} \cdot \left(\frac{1}{3}\right)^{x_2} < 1$$

Seriously though

What if $k = 1$?

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} < 1$$

Example for $k = n = 2$ and fixed $0 < p < 1$

Do there exist $x_1, x_2 \in \mathbb{N}$ such that

$$p \cdot \left(\frac{1}{2}\right)^{x_1} \cdot 3^{x_2} + (1 - p) \cdot 2^{x_1} \cdot \left(\frac{1}{3}\right)^{x_2} < 1$$

Why is this hard? Let's do some logarithms

Seriously though

What if $k = 1$?

do there exists $x_1, \dots, x_n \in \mathbb{N}$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \dots q_{i,n}^{x_n} < 1$$

Example for $k = n = 2$ and fixed $0 < p < 1$

Do there exist $x_1, x_2 \in \mathbb{N}$ such that

$$p \cdot \left(\frac{1}{2}\right)^{x_1} \cdot 3^{x_2} + (1 - p) \cdot 2^{x_1} \cdot \left(\frac{1}{3}\right)^{x_2} < 1$$

Why is this hard? Let's do some logarithms

$$\begin{aligned} & \exp(\log(p) - x_1 \log(2) + x_2 \log(3)) \\ & + \exp(\log(1 - p) + x_1 \log(2) - x_2 \log(3)) < 1 \end{aligned}$$

Geometric presentation

$$\begin{aligned} & \exp(\log(p) - x_1 \log(2) + x_2 \log(3)) \\ & + \exp(\log(1 - p) + x_1 \log(2) - x_2 \log(3)) \end{aligned} < 1$$


Geometric presentation

$$\overbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}^x + \underbrace{\exp(\log(1 - p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$

Geometric presentation

$$\underbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}_x + \underbrace{\exp(\log(1-p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$

$$V = \{(x, y) \in \mathbb{R}^2 \mid e^x + e^y < 1\} \quad (\log = \log_e)$$




Geometric presentation

$$\overbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}^x + \underbrace{\exp(\log(1 - p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$

$$V = \{(x, y) \in \mathbb{R}^2 \mid e^x + e^y < 1\} \quad (\log = \log_e)$$

$$b = (\log(p), \log(1 - p)) \notin V$$

(for $x_1 = x_2 = 0$)



Geometric presentation

$$\underbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}_x + \underbrace{\exp(\log(1-p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$


$$V = \{(x, y) \in \mathbb{R}^2 \mid e^x + e^y < 1\} \quad (\log = \log_e)$$

$$b = (\log(p), \log(1-p)) \notin V$$

(for $x_1 = x_2 = 0$)

$$u = (-\log(2), \log(2)),$$

$$v = (\log(3), -\log(3))$$



Geometric presentation

$$\underbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}_x + \underbrace{\exp(\log(1-p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$

$$V = \{(x, y) \in \mathbb{R}^2 \mid e^x + e^y < 1\} \quad (\log = \log_e)$$


$$b = (\log(p), \log(1-p)) \notin V$$

(for $x_1 = x_2 = 0$)

$$u = (-\log(2), \log(2)),$$

$$v = (\log(3), -\log(3))$$

$$b + x_1 u + x_2 v \in V?$$



Geometric presentation

$$\underbrace{\exp(\log(p) - x_1 \log(2) + x_2 \log(3))}_x + \underbrace{\exp(\log(1-p) + x_1 \log(2) - x_2 \log(3))}_y < 1$$

$$V = \{(x, y) \in \mathbb{R}^2 \mid e^x + e^y < 1\} \quad (\log = \log_e)$$


$$b = (\log(p), \log(1-p)) \notin V$$

(for $x_1 = x_2 = 0$)

$$u = (-\log(2), \log(2)),$$

$$v = (\log(3), -\log(3))$$

$$b + x_1 u + x_2 v \in V? \quad \text{yes iff } p \neq \frac{1}{2}$$



Back to the problem

Is there $\mathbf{x} \in \mathbb{N}^n$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Back to the problem

Is there $\mathbf{x} \in \mathbb{N}^n$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(1) This is semi-decidable (enumerate all \mathbf{x})

Back to the problem

Is there $\mathbf{x} \in \mathbb{N}^n$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(1) This is semi-decidable (enumerate all \mathbf{x})

The complement problem: for all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Back to the problem

Is there $\mathbf{x} \in \mathbb{N}^n$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(1) This is semi-decidable (enumerate all \mathbf{x})

The complement problem: for all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(2) Is this also semi-decidable?

Back to the problem

Is there $\mathbf{x} \in \mathbb{N}^n$ such that

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(1) This is semi-decidable (enumerate all \mathbf{x})

The complement problem: for all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

(2) Is this also semi-decidable?

Corollary

(1) + (2) give decidability

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Allow $\mathbf{x} \in \mathbb{R}^n$, denote by X all solutions (in \mathbb{R}^n)

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Allow $\mathbf{x} \in \mathbb{R}^n$, denote by X all solutions (in \mathbb{R}^n)

Key Lemma (I'm not proving it)

If $X \cap \mathbb{Z}^n = \emptyset$ then exist $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$ s. t. $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Allow $\mathbf{x} \in \mathbb{R}^n$, denote by X all solutions (in \mathbb{R}^n)

Key Lemma (I'm not proving it)

If $X \cap \mathbb{Z}^n = \emptyset$ then exist $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$ s. t. $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Allow $\mathbf{x} \in \mathbb{R}^n$, denote by X all solutions (in \mathbb{R}^n)

Key Lemma (I'm not proving it)

If $X \cap \mathbb{Z}^n = \emptyset$ then exist $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$ s. t. $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Given \mathbf{d} , a , b expressible in first-order logic over $(\mathbb{R}, +, \times, \exp)$

Solving the complement problem

For all $\mathbf{x} \in \mathbb{N}^n$ this is not true

$$\sum_{i=1}^k p_i q_{i,1}^{x_1} \cdots q_{i,n}^{x_n} < 1$$

Allow $\mathbf{x} \in \mathbb{R}^n$, denote by X all solutions (in \mathbb{R}^n)

Key Lemma (I'm not proving it)

If $X \cap \mathbb{Z}^n = \emptyset$ then exist $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$ s. t. $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Given \mathbf{d} , a , b expressible in first-order logic over $(\mathbb{R}, +, \times, \exp)$

Satisfiability decidable assuming Schanuel [Macintyre, Wilkie; 1996]

Solving the complement problem (continued)

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

Solving the complement problem (continued)

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

- Step 2. For all $i \in [a, b]$ ($i \in \mathbb{Z}$)

consider $\mathbf{d}^\top \mathbf{x} = d_1 x_1 + \dots + d_n x_n = i$

Solving the complement problem (continued)

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

- Step 2. For all $i \in [a, b]$ ($i \in \mathbb{Z}$)

consider $\mathbf{d}^\top \mathbf{x} = d_1 x_1 + \dots + d_n x_n = i$

If there is a solution:

recursively start over with dimension reduced by 1 (or output NO)

Solving the complement problem (continued)

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

- Step 2. For all $i \in [a, b]$ ($i \in \mathbb{Z}$)

consider $\mathbf{d}^\top \mathbf{x} = d_1 x_1 + \dots + d_n x_n = i$

If there is a solution:

recursively start over with dimension reduced by 1 (or output NO)

If there is no solution: try another i

If you tried all i then go up in the recursion (or output YES)

Solving the complement problem (continued)

Procedure:

- Step 1. Find $\mathbf{d} \in \mathbb{Z}^n$, $a, b \in \mathbb{Z}$

such that $\{\mathbf{d}^\top \mathbf{x} \mid \mathbf{x} \in X\} \subseteq [a, b]$

- Step 2. For all $i \in [a, b]$ ($i \in \mathbb{Z}$)

consider $\mathbf{d}^\top \mathbf{x} = d_1 x_1 + \dots + d_n x_n = i$

If there is a solution:

recursively start over with dimension reduced by 1 (or output NO)

If there is no solution: try another i

If you tried all i then go up in the recursion (or output YES)

Previous lemma guarantees semi-decidability (\mathbb{Z} to \mathbb{N} is a technicality)

Conclusion

Conclusion

Containment for finitely ambiguous PA:

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

- Getting rid of Schanuel

Show some small solution property

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

- Getting rid of Schanuel

Show some small solution property

- Getting rid of the unambiguity assumption

We don't know if the Key Lemma holds

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

- Getting rid of Schanuel

Show some small solution property

- Getting rid of the unambiguity assumption

We don't know if the Key Lemma holds

Recall $V = \{(x_1, \dots, x_k) \mid e^{x_1} + \dots + e^{x_k} < 1\}$

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

- Getting rid of Schanuel

Show some small solution property

- Getting rid of the unambiguity assumption

We don't know if the Key Lemma holds

Recall $V = \{(x_1, \dots, x_k) \mid e^{x_1} + \dots + e^{x_k} < 1\}$

Nice, convex etc.

Conclusion

Containment for finitely ambiguous PA:

- We solved it assuming:

Schanuel and that one automaton is unambiguous

- Getting rid of Schanuel

Show some small solution property

- Getting rid of the unambiguity assumption

We don't know if the Key Lemma holds

Recall $V = \{(x_1, \dots, x_k) \mid e^{x_1} + \dots + e^{x_k} < 1\}$

Nice, convex etc.

$V' = \{(x_1, \dots, x_k, y_1, \dots, y_l) \mid e^{x_1} + \dots + e^{x_k} < e^{y_1} + \dots + e^{y_l}\} ???$