

# Lecture 1

Weighted automata basic definitions

# Finite automata

## Definition

A finite automaton is  $\mathcal{A} = (Q, \Sigma, T, I, F)$ , where:

- $Q$  is a finite set of states
- $\Sigma$  is a finite alphabet
- $T \subseteq Q \times \Sigma \times Q$  is a finite set of transitions
- $I, F \subseteq Q$  are the sets of initial and final states


# Finite automata

## Definition

A finite automaton is  $\mathcal{A} = (Q, \Sigma, T, I, F)$ , where:

- $Q$  is a finite set of states
- $\Sigma$  is a finite alphabet
- $T \subseteq Q \times \Sigma \times Q$  is a finite set of transitions
- $I, F \subseteq Q$  are the sets of initial and final states

Example:



- $Q = \{p, q\}$
- $\Sigma = \{a, b\}$
- $T = \{(p, a, p), (p, b, p), (p, b, q), (q, b, q)\}$
- $I = \{p\}, F = \{q\}$

## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

It is an accepting run if

- $p_0 \in I, q_n \in F$

## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

It is an accepting run if

- $p_0 \in I, q_n \in F$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

It is an accepting run if

- $p_0 \in I, q_n \in F$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For every  $\rho \in R_w$ :  $val(\rho) = true$  if  $\rho$  is accepting, otherwise  $val(\rho_w) = false$

## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

It is an accepting run if

- $p_0 \in I, q_n \in F$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For every  $\rho \in R_w$ :  $val(\rho) = true$  if  $\rho$  is accepting, otherwise  $val(\rho_w) = false$

- $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{false, true\}$

$$\llbracket \mathcal{A} \rrbracket (w) = \bigvee_{\rho \in R_w} val(\rho)$$



## Finite automata runs

Let  $\mathcal{A} = (Q, \Sigma, T, I, F)$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

It is an accepting run if

- $p_0 \in I, q_n \in F$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For every  $\rho \in R_w$ :  $val(\rho) = true$  if  $\rho$  is accepting, otherwise  $val(\rho_w) = false$

- $\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \{false, true\}$

$$\llbracket \mathcal{A} \rrbracket (w) = \bigvee_{\rho \in R_w} val(\rho)$$

$$L(\mathcal{A}) = \{w \mid \llbracket \mathcal{A}(w) \rrbracket = true\}$$

## Automata counting things

What about  $\mathcal{A} : \Sigma^* \rightarrow \text{numbers}, \mathbb{N}?, \mathbb{Q}?$

How many  $a$ 's are there in the word?

What is the probability of acceptance?


## Automata counting things

What about  $\mathcal{A} : \Sigma^* \rightarrow \text{numbers}, \mathbb{N}?, \mathbb{Q}?$

How many  $a$ 's are there in the word?

What is the probability of acceptance?

Add values on transitions




## Automata counting things

What about  $\mathcal{A} : \Sigma^* \rightarrow \text{numbers}, \mathbb{N}?, \mathbb{Q}?$

How many  $a$ 's are there in the word?

What is the probability of acceptance?

Add values on transitions




## Automata counting things

What about  $\mathcal{A} : \Sigma^* \rightarrow \text{numbers}, \mathbb{N}?, \mathbb{Q}?$

How many  $a$ 's are there in the word?

What is the probability of acceptance?

Add values on transitions



To discuss what numbers first we will describe the semiring structure in the following slides

## Commutative semirings

$\mathbb{S}(\oplus, \odot, 0, 1)$  a set  $\mathbb{S}$  with two operations and axioms

# Commutative semirings

$\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  a set  $\mathbb{S}$  with two operations and axioms

1.  $(\mathbb{S}, \oplus)$  is a commutative monoid with identity  $\mathbb{0}$

- $(a \oplus b) \oplus c = a \oplus (b \oplus c)$
- $\mathbb{0} \oplus a = a \oplus \mathbb{0} = a$
- $a \oplus b = b \oplus a$

2.  $(\mathbb{S}, \odot)$  is a commutative monoid with identity  $\mathbb{1}$

- $(a \odot b) \odot c = a \odot (b \odot c)$
- $\mathbb{1} \odot a = a \odot \mathbb{1} = a$
- $a \odot b = b \odot a$

3. Distributivity

- $a \odot (b \oplus c) = a \odot b \oplus a \odot c$

4. Annihilation

- $\mathbb{0} \odot a = a \odot \mathbb{0} = \mathbb{0}$

## (Commutative) semirings examples

- Rings like  $(\mathbb{Q}, +, \cdot, 0, 1)$
- Natural numbers  $(\mathbb{N}, +, \cdot, 0, 1)$



## (Commutative) semirings examples

- Rings like  $(\mathbb{Q}, +, \cdot, 0, 1)$
- Natural numbers  $(\mathbb{N}, +, \cdot, 0, 1)$

### Remark

$(\mathbb{N}, +, \cdot, 0, 1)$  is not a ring because  $-1 \notin \mathbb{N}$ .

## (Commutative) semirings examples

- Rings like  $(\mathbb{Q}, +, \cdot, 0, 1)$
- Natural numbers  $(\mathbb{N}, +, \cdot, 0, 1)$

### Remark

$(\mathbb{N}, +, \cdot, 0, 1)$  is not a ring because  $-1 \notin \mathbb{N}$ .

### Tropical semirings

- $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ , where  $\mathbb{N}_{+\infty} = \mathbb{N} \cup \{+\infty\}$
- $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$ , where  $\mathbb{N}_{-\infty} = \mathbb{N} \cup \{-\infty\}$

## (Commutative) semirings examples

- Rings like  $(\mathbb{Q}, +, \cdot, 0, 1)$
- Natural numbers  $(\mathbb{N}, +, \cdot, 0, 1)$

### Remark

$(\mathbb{N}, +, \cdot, 0, 1)$  is not a ring because  $-1 \notin \mathbb{N}$ .

### Tropical semirings

- $(\mathbb{N}_{+\infty}, \min, +, \infty, 0)$ , where  $\mathbb{N}_{+\infty} = \mathbb{N} \cup \{+\infty\}$
- $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$ , where  $\mathbb{N}_{+\infty} = \mathbb{N} \cup \{-\infty\}$

Note:  $\oplus = \min$ ,  $\odot = +$ ,  $\mathbb{0} = +\infty$ ,  $\mathbb{1} = 0$

Axioms work:

$$n \oplus \mathbb{0} = n \quad \text{becomes} \quad \min(n, +\infty) = n$$

$$n \odot \mathbb{1} = n \quad \text{becomes} \quad n + 0 = n$$

# Weighted automata

## Definition

A weighted automaton over a semiring  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $\mathcal{A} = (Q, \Sigma, T, I, F)$ :

- $Q$  is a finite set of states
- $\Sigma$  is a finite alphabet
- $T \subseteq Q \times \Sigma \times \mathbb{S} \times Q$  is a finite set of transitions
- $I, F : Q \rightarrow \mathbb{S}$  are the initial and final functions

# Weighted automata

## Definition

A weighted automaton over a semiring  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $\mathcal{A} = (Q, \Sigma, T, I, F)$ :

- $Q$  is a finite set of states
- $\Sigma$  is a finite alphabet
- $T \subseteq Q \times \Sigma \times \mathbb{S} \times Q$  is a finite set of transitions
- $I, F : Q \rightarrow \mathbb{S}$  are the initial and final functions

## Remark

For every  $p, q \in Q$  and  $a \in \Sigma$  we assume there is at most one  $(p, a, s, q) \in T$

# Weighted automata

## Definition

A weighted automaton over a semiring  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $\mathcal{A} = (Q, \Sigma, T, I, F)$ :

- $Q$  is a finite set of states
- $\Sigma$  is a finite alphabet
- $T \subseteq Q \times \Sigma \times \mathbb{S} \times Q$  is a finite set of transitions
- $I, F : Q \rightarrow \mathbb{S}$  are the initial and final functions

## Remark

For every  $p, q \in Q$  and  $a \in \Sigma$  we assume there is at most one  $(p, a, s, q) \in T$

So ignoring  $\mathbb{S}$  in  $T$

and identifying  $I$  (and  $F$ ) with  $I' = \{q \mid I(q) \neq \mathbb{0}\}$

we get a finite automaton

## Weighted automata runs and output

Given  $\mathcal{A} = (Q, \Sigma, T, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, s_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

## Weighted automata runs and output

Given  $\mathcal{A} = (Q, \Sigma, T, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, s_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For  $t = (p, a, s, q) \in T$  we write  $val(t) = s$ .



## Weighted automata runs and output

Given  $\mathcal{A} = (Q, \Sigma, T, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, s_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For  $t = (p, a, s, q) \in T$  we write  $val(t) = s$ .

For every  $\rho = t_1 \dots t_n \in R_w$ :  $val(\rho) = I(p_0) \odot \bigodot_{i=1}^n val(t_i) \odot F(q_n)$

## Weighted automata runs and output

Given  $\mathcal{A} = (Q, \Sigma, T, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$

A run of  $\mathcal{A}$  on  $w = a_1 \dots a_n \in \Sigma^*$  is  $\rho = t_1 \dots t_n$ , where:

- $t_i = (p_i, a_i, s_i, q_i) \in T$
- $q_{i-1} = p_i$  for all  $i = 2, \dots, n$

The set of all run of  $\mathcal{A}$  on  $w$  is  $R_w$

For  $t = (p, a, s, q) \in T$  we write  $val(t) = s$ .

For every  $\rho = t_1 \dots t_n \in R_w$ :  $val(\rho) = I(p_0) \odot \bigodot_{i=1}^n val(t_i) \odot F(q_n)$

Then  $\llbracket \mathcal{A} \rrbracket (w) = \bigoplus_{\rho \in R_w} val(\rho)$        $\llbracket \mathcal{A} \rrbracket (\epsilon) = \bigoplus_{q \in Q} I(q) \odot F(q)$


## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$


- Longest block of  $b$ 's



## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Longest block of  $b$ 's




$$T = \{(p, a, 0, p), (p, b, 0, p), (p, b, 1, q), (q, b, 1, q), (q, a, 0, r), \dots\}$$

## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Longest block of  $b$ 's



$$T = \{(p, a, 0, p), (p, b, 0, p), (p, b, 1, q), (q, b, 1, q), (q, a, 0, r), \dots\}$$


### Remark

Usually  $I, F : Q \rightarrow \{0, 1\} = \{+\infty, 0\}$ . Then initial state means the value of  $I$  is  $1$  and  $0$  otherwise. Here,  $I(p) = 0$ ,  $I(q) = +\infty$  and  $I(r) = +\infty$ . Similarly with accepting states and  $F$ .

## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$


- Longest block of  $b$ 's



## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Longest block of  $b$ 's



- Let  $w = bbab$

All runs starting in  $q$  or  $r$  have value  $-\infty + \dots + = -\infty$


All runs ending in  $p$  have value  $\dots + (-\infty) = -\infty$



## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Longest block of  $b$ 's



- Let  $w = bbab$

All runs starting in  $q$  or  $r$  have value  $-\infty + \dots + = -\infty$

All runs ending in  $p$  have value  $\dots + (-\infty) = -\infty$


There are three other runs (skipping 0's from  $I$  and  $F$ )

$$1 + 1 + 0 + 0 = 2, \quad 0 + 1 + 0 + 0 = 1, \quad 0 + 0 + 0 + 1 = 2$$

## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Longest block of  $b$ 's



- Let  $w = bbab$

All runs starting in  $q$  or  $r$  have value  $-\infty + \dots + = -\infty$

All runs ending in  $p$  have value  $\dots + (-\infty) = -\infty$

There are three other runs (skipping 0's from  $I$  and  $F$ )


$$1 + 1 + 0 + 0 = 2, \quad 0 + 1 + 0 + 0 = 1, \quad 0 + 0 + 0 + 1 = 2$$

$$\llbracket \mathcal{A} \rrbracket (bbab) = \max\{2, 1, 1, -\infty\} = 2$$

## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$


- Maximum of number of  $a$ 's and number of  $b$ 's



## Weighted automata examples

Fix the semiring  $(\mathbb{N}_{-\infty}, \max, +, -\infty, 0)$

- Maximum of number of  $a$ 's and number of  $b$ 's



There are always two runs. Consider  $bbab$

$$0 + 0 + 1 + 0 = 1 \text{ and } 1 + 1 + 0 + 1 = 3$$


$$\text{Output: } \max\{1, 3\} = 3$$

## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$

## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$




There are always two runs. Consider  $bbab$

$$0 \cdot 0 \cdot 1 \cdot 0 = 0 \text{ and } 1 \cdot 1 \cdot 0 \cdot 1 = 0$$

Output:  $0 + 0 = 0$


## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$



## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$



There are always two runs. Consider  $bbab$

$$1 \cdot 1 \cdot 2 \cdot 1 = 2 \text{ and } 2 \cdot 2 \cdot 1 \cdot 2 = 8$$


Output:  $2 + 8 = 10$



## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$

This is  $\llbracket \mathcal{A} \rrbracket (w) = \max\{2^{\#_a(w)}, 2^{\#_b(w)}\}$



There are always two runs. Consider  $bbab$


$$1 \cdot 1 \cdot 2 \cdot 1 = 2 \text{ and } 2 \cdot 2 \cdot 1 \cdot 2 = 8$$

Output:  $2 + 8 = 10$

## Weighted automata examples

Change the semiring to  $(\mathbb{Q}, +, \cdot, 0, 1)$

This is  $\llbracket \mathcal{A} \rrbracket (w) = \max\{2^{\#_a(w)}, 2^{\#_b(w)}\}$



There are always two runs. Consider  $bbab$

$$1 \cdot 1 \cdot 2 \cdot 1 = 2 \text{ and } 2 \cdot 2 \cdot 1 \cdot 2 = 8$$

Output:  $2 + 8 = 10$

It is important to write the semiring of the weighted automaton

## Weighted automata over the boolean semiring

Consider the semiring  $(\{0, 1\}, \vee, \wedge, 0, 1)$  (0 is false, 1 is true)

## Weighted automata over the boolean semiring

Consider the semiring  $(\{0, 1\}, \vee, \wedge, 0, 1)$  (0 is false, 1 is true)

- Definitions as expected

$$0 \vee 1 = 1, \quad 0 \wedge 1 = 0 \text{ etc.} \dots$$

It is a semiring

## Weighted automata over the boolean semiring

Consider the semiring  $(\{0, 1\}, \vee, \wedge, 0, 1)$  (0 is false, 1 is true)

- Definitions as expected

$$0 \vee 1 = 1, \quad 0 \wedge 1 = 0 \text{ etc.} \dots$$

It is a semiring

- Weighted automata over this semiring are finite automata

Initial, final states are states such that  $I(q) = 1$  and  $F(q) = 1$

Transitions in finite automata are transitions such that  $val(t) = 1$

## Weighted automata over the boolean semiring

Consider the semiring  $(\{0, 1\}, \vee, \wedge, 0, 1)$  (0 is false, 1 is true)

- Definitions as expected

$$0 \vee 1 = 1, \quad 0 \wedge 1 = 0 \text{ etc.} \dots$$

It is a semiring

- Weighted automata over this semiring are finite automata

Initial, final states are states such that  $I(q) = 1$  and  $F(q) = 1$

Transitions in finite automata are transitions such that  $val(t) = 1$

Then  $val(\rho) = 1 \wedge 1 \wedge 1 \dots \wedge 1 = 1$  if  $\rho$  is accepting

and  $val(\rho) = 0$  otherwise

## Weighted automata over the boolean semiring

Consider the semiring  $(\{0, 1\}, \vee, \wedge, 0, 1)$  (0 is false, 1 is true)

- Definitions as expected

$$0 \vee 1 = 1, \quad 0 \wedge 1 = 0 \text{ etc.} \dots$$

It is a semiring

- Weighted automata over this semiring are finite automata

Initial, final states are states such that  $I(q) = 1$  and  $F(q) = 1$

Transitions in finite automata are transitions such that  $val(t) = 1$

Then  $val(\rho) = 1 \wedge 1 \wedge 1 \dots \wedge 1 = 1$  if  $\rho$  is accepting

and  $val(\rho) = 0$  otherwise

The output is

$$\llbracket \mathcal{A} \rrbracket (w) = \bigvee val(\rho)$$

## Weighted automata different definition

### Definition

A weighted automaton  $\mathcal{A}$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $(d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$ , where:

- $d \in \mathbb{N}$  is the dimension;
- $\Sigma$  is a finite alphabet;
- every  $M_a$  is a  $d \times d$  matrix over  $\mathbb{S}$ ;
- $I$  and  $F$  are the initial and the final vector in  $\mathbb{S}^d$ .



## Weighted automata different definition

### Definition

A weighted automaton  $\mathcal{A}$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $(d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$ , where:

- $d \in \mathbb{N}$  is the dimension;
- $\Sigma$  is a finite alphabet;
- every  $M_a$  is a  $d \times d$  matrix over  $\mathbb{S}$ ;
- $I$  and  $F$  are the initial and the final vector in  $\mathbb{S}^d$ .

$$\llbracket \mathcal{A} \rrbracket : \Sigma^* \rightarrow \mathbb{Q}$$

$$\llbracket \mathcal{A} \rrbracket (a_1 a_2 \dots a_n) = I^\top \odot M_{a_1} M_{a_2} \dots M_{a_n} \odot F$$

## Weighted automata different definition

### Definition

A weighted automaton  $\mathcal{A}$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$  is  $(d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$ , where:

- $d \in \mathbb{N}$  is the dimension;
- $\Sigma$  is a finite alphabet;
- every  $M_a$  is a  $d \times d$  matrix over  $\mathbb{S}$ ;
- $I$  and  $F$  are the initial and the final vector in  $\mathbb{S}^d$ .

$$[\![\mathcal{A}]\!] : \Sigma^* \rightarrow \mathbb{Q}$$

$$[\![\mathcal{A}]\!](a_1 a_2 \dots a_n) = I^\top \odot M_{a_1} M_{a_2} \dots M_{a_n} \odot F$$

### Remark

It makes sense to multiply matrices over any semiring. Over  $\mathbb{N}(\max, +)$  :

$$\begin{pmatrix} 0 & -\infty \\ -\infty & 1 \end{pmatrix} \begin{pmatrix} 1 & -\infty \\ -\infty & 0 \end{pmatrix} = \begin{pmatrix} \max(0 + 1, -\infty + -\infty) & \max(0 + -\infty, -\infty + 0) \\ \max(-\infty + 1, 1 + -\infty) & \max(1 + 0, -\infty + -\infty) \end{pmatrix}$$

## Definition equivalence

Set  $\mathcal{A} = (d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$ :

## Definition equivalence

Set  $\mathcal{A} = (d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$ :

- We define  $\mathcal{A}' = (Q, \Sigma, T, I, F)$

$$Q = \{1, \dots, d\}$$

Vectors  $F, I \in \mathbb{S}^d$  are the same thing as a functions  $F, I : Q \rightarrow \mathbb{S}$

## Definition equivalence

Set  $\mathcal{A} = (d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$ :

- We define  $\mathcal{A}' = (Q, \Sigma, T, I, F)$

$$Q = \{1, \dots, d\}$$

Vectors  $F, I \in \mathbb{S}^d$  are the same thing as a functions  $F, I : Q \rightarrow \mathbb{S}$

$$T = \{(p, a, s, q) \mid p, q \in Q, a \in \Sigma, s = M_a[p, q]\}$$

## Definition equivalence

Set  $\mathcal{A} = (d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$ :

- We define  $\mathcal{A}' = (Q, \Sigma, T, I, F)$

$$Q = \{1, \dots, d\}$$

Vectors  $F, I \in \mathbb{S}^d$  are the same thing as a functions  $F, I : Q \rightarrow \mathbb{S}$

$$T = \{(p, a, s, q) \mid p, q \in Q, a \in \Sigma, s = M_a[p, q]\}$$

### Theorem

$$\llbracket \mathcal{A} \rrbracket (w) = \llbracket \mathcal{A}' \rrbracket (w) \text{ for all } w \in \Sigma^*.$$

## Definition equivalence

Set  $\mathcal{A} = (d, \Sigma, \{M_a\}_{a \in \Sigma}, I, F)$  over  $\mathbb{S}(\oplus, \odot, \mathbb{0}, \mathbb{1})$ :

- We define  $\mathcal{A}' = (Q, \Sigma, T, I, F)$

$$Q = \{1, \dots, d\}$$

Vectors  $F, I \in \mathbb{S}^d$  are the same thing as a functions  $F, I : Q \rightarrow \mathbb{S}$

$$T = \{(p, a, s, q) \mid p, q \in Q, a \in \Sigma, s = M_a[p, q]\}$$

### Theorem

$$\llbracket \mathcal{A} \rrbracket (w) = \llbracket \mathcal{A}' \rrbracket (w) \text{ for all } w \in \Sigma^*.$$

Proof.

$$\llbracket \mathcal{A} \rrbracket (\epsilon) = I^\top \odot F, \quad \llbracket \mathcal{A}' \rrbracket (\epsilon) = \bigoplus_{i=1}^d I(i) \odot F(i)$$

## Induction for $|w| > 0$

### Definition

$R_w^{p,q}$  is the set of runs in  $\mathcal{A}'$  from state  $p$  to state  $q$  over  $w$

For every  $\rho = t_1 \dots t_n$  we denote by  $trans(\rho) = val(t_1) \odot \dots \odot val(t_n)$   
( $val(\rho)$  ignoring  $I$  and  $F$ )



## Induction for $|w| > 0$

### Definition

$R_w^{p,q}$  is the set of runs in  $\mathcal{A}'$  from state  $p$  to state  $q$  over  $w$

For every  $\rho = t_1 \dots t_n$  we denote by  $trans(\rho) = val(t_1) \odot \dots \odot val(t_n)$   
( $val(\rho)$  ignoring  $I$  and  $F$ )

### Lemma

$$M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} trans(\rho)$$

## Induction for $|w| > 0$

### Definition

$R_w^{p,q}$  is the set of runs in  $\mathcal{A}'$  from state  $p$  to state  $q$  over  $w$

For every  $\rho = t_1 \dots t_n$  we denote by  $trans(\rho) = val(t_1) \odot \dots \odot val(t_n)$   
( $val(\rho)$  ignoring  $I$  and  $F$ )

### Lemma

$$M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} trans(\rho)$$

Proof of Lemma (by induction on  $|w|$ ).

if  $|w| = 1$  then there is only one run from  $p$  to  $q$

the transition from  $p$  to  $q$  whose value is  $M[p, q]$  by definition

## Induction for $|w| > 0$

### Definition

$R_w^{p,q}$  is the set of runs in  $\mathcal{A}'$  from state  $p$  to state  $q$  over  $w$

For every  $\rho = t_1 \dots t_n$  we denote by  $trans(\rho) = val(t_1) \odot \dots \odot val(t_n)$   
( $val(\rho)$  ignoring  $I$  and  $F$ )

### Lemma

$$M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} trans(\rho)$$

Proof of Lemma (by induction on  $|w|$ ).

if  $|w| = 1$  then there is only one run from  $p$  to  $q$

the transition from  $p$  to  $q$  whose value is  $M[p, q]$  by definition

if  $|w| > 1$  then write  $w = av$  for  $v \in \Sigma^+$  and  $a \in \Sigma$

## Induction continued

Notation  $t_i = (p_i, a_i, s_i, q_i) \in T$ , where  $q_i = p_{i+1}$

Then  $R_w^{p,q} = \{t_1 t_2 \dots t_n \mid p_1 = p, t_2 \dots t_n \in R_v^{p_2, q}\}$

## Induction continued

Notation  $t_i = (p_i, a_i, s_i, q_i) \in T$ , where  $q_i = p_{i+1}$

Then  $R_w^{p,q} = \{t_1 t_2 \dots t_n \mid p_1 = p, t_2 \dots t_n \in R_v^{p_2, q}\}$

$$\text{So } \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho) = \bigoplus_{q_1 = p_2 \in Q} M_{a_1}[p_1, q_1] \odot \bigoplus_{\rho' \in R_v^{p_2, q}} \text{trans}(\rho')$$

## Induction continued

Notation  $t_i = (p_i, a_i, s_i, q_i) \in T$ , where  $q_i = p_{i+1}$

Then  $R_w^{p,q} = \{t_1 t_2 \dots t_n \mid p_1 = p, t_2 \dots t_n \in R_v^{p_2, q}\}$

$$\text{So } \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho) = \bigoplus_{q_1 = p_2 \in Q} M_{a_1}[p_1, q_1] \odot \bigoplus_{\rho' \in R_v^{p_2, q}} \text{trans}(\rho')$$

$$\text{By induction} = \bigoplus_{q_1 = p_2 \in Q} M_{a_1}[p_1, q_1] \odot M_{a_2} \dots M_{a_n}[p_2, q]$$

## Induction continued

Notation  $t_i = (p_i, a_i, s_i, q_i) \in T$ , where  $q_i = p_{i+1}$

Then  $R_w^{p,q} = \{t_1 t_2 \dots t_n \mid p_1 = p, t_2 \dots t_n \in R_v^{p_2, q}\}$

$$\text{So } \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho) = \bigoplus_{q_1 = p_2 \in Q} M_{a_1}[p_1, q_1] \odot \bigoplus_{\rho' \in R_v^{p_2, q}} \text{trans}(\rho')$$

$$\text{By induction } = \bigoplus_{q_1 = p_2 \in Q} M_{a_1}[p_1, q_1] \odot M_{a_2} \dots M_{a_n}[p_2, q]$$

The lemma follows from the definition of matrix multiplication

For any matrices  $A, B$  in dimensions  $d$  we have

$$AB[p, q] = \bigoplus_{i \in \{1, \dots, d\}} A[p, i] B[i, q]$$

## Final step

We proved  $M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho)$



## Final step

We proved  $M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho)$

But  $\llbracket \mathcal{A} \rrbracket(w) = I^\top \odot M_{a_1} \dots M_{a_n} \odot F = \bigoplus_{p,q \in Q} I(p) \odot M_{a_1} \dots M_{a_n}[p, q] \odot F(q)$

## Final step

We proved  $M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho)$

But  $\llbracket \mathcal{A} \rrbracket(w) = I^T \odot M_{a_1} \dots M_{a_n} \odot F = \bigoplus_{p,q \in Q} I(p) \odot M_{a_1} \dots M_{a_n}[p, q] \odot F(q)$

By lemma  $= \bigoplus_{p,q \in Q} \bigoplus_{\rho \in R_w^{p,q}} I(p) \odot \text{trans}(\rho) \odot F(q)$

## Final step

We proved  $M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho)$

But  $\llbracket \mathcal{A} \rrbracket(w) = I^\top \odot M_{a_1} \dots M_{a_n} \odot F = \bigoplus_{p,q \in Q} I(p) \odot M_{a_1} \dots M_{a_n}[p, q] \odot F(q)$

By lemma  $= \bigoplus_{p,q \in Q} \bigoplus_{\rho \in R_w^{p,q}} I(p) \odot \text{trans}(\rho) \odot F(q)$

Which is equal to  $\bigoplus_{\rho \in R_w} \text{val}(\rho)$



## Final step

We proved  $M_{a_1} \dots M_{a_n}[p, q] = \bigoplus_{\rho \in R_w^{p,q}} \text{trans}(\rho)$

But  $\llbracket \mathcal{A} \rrbracket(w) = I^\top \odot M_{a_1} \dots M_{a_n} \odot F = \bigoplus_{p,q \in Q} I(p) \odot M_{a_1} \dots M_{a_n}[p, q] \odot F(q)$

By lemma  $= \bigoplus_{p,q \in Q} \bigoplus_{\rho \in R_w^{p,q}} I(p) \odot \text{trans}(\rho) \odot F(q)$

Which is equal to  $\bigoplus_{\rho \in R_w} \text{val}(\rho)$



The opposite translation on tutorials