

# CSCI 2011 HW 8

Fletcher Gornick

November 3, 2020

## 1 6.2 Problem 8

**For the function  $f$  defined by  $f(n) = \frac{n^2+1}{n+1}$  for each  $n \in \mathbb{N}$ , show that  $f(n) = O(n)$ .**

for  $n \geq 1$ ,  $f(n) = \frac{n^2+1}{n+1} \leq \frac{n^2+n}{n+1} < \frac{n^2+n}{n} = n+1 \leq n+n = 2n$ . Therefore  $f(n) < 2n$  for  $n \geq 1$ , and so  $f(n) = O(n)$ .

## 2 Chapter 6 Problem 12

**Let  $f : \mathbb{N} \rightarrow \mathbb{R}^+$  and  $g : \mathbb{N} \rightarrow \mathbb{R}^+$  be defined by  $f(n) = 2n^3 + n + 10$  and  $g(n) = n^3 + 4n^2 + 1$  for  $n \in \mathbb{N}$ . Show that  $f = \Theta(g)$ .**

First we show that  $f = O(g)$ .

for  $n \geq 1$ ,  $f(n) = 2n^3 + n + 10 \leq 2n^3 + n^2 + 10 < 10n^3 + 40n^2 + 10 = 10(n^3 + 4n^2 + 1)$ . Therefore  $f(n) < 10 \cdot g(n)$  for  $n \geq 1$ , and so  $f = O(g)$ .

Now we show that  $f = \Omega(g)$  or, put more simply,  $g = O(f)$ .

for  $n \geq 1$ ,  $g(n) = n^3 + 4n^2 + 1 \leq n^3 + 4n^3 + 1 = 5n^3 + 1 < 10n^3 + 50 < 10n^3 + 5n + 50 = 5(2n^3 + n + 10)$ . Therefore  $g(n) < 5 \cdot f(n)$  for  $n \geq 1$ , and so  $g = O(f)$ .

Since  $f = O(g)$  and  $g = O(f)$ , it must be the case that  $f = \Theta(g)$

## 3 6.2 Problem 14

**Let  $f : \mathbb{N} \rightarrow \mathbb{R}^+$ ,  $g : \mathbb{N} \rightarrow \mathbb{R}^+$  and  $h : \mathbb{N} \rightarrow \mathbb{R}^+$  be three functions. Prove that if  $f = \Theta(g)$  and  $g = \Theta(h)$ , then  $f = \Theta(h)$ .**

For  $n, k \in \mathbb{Z}$ ,  $n \geq k$ ,  $f = \Theta(g)$ , so there must exist  $c_1, c_2 \in \mathbb{N}$ , such that  $c_1g(n) \leq f(n) \leq c_2g(n)$ .

For  $n, k \in \mathbb{Z}$ ,  $n \geq k$ ,  $g = \Theta(h)$ , so there must exist  $d_1, d_2 \in \mathbb{N}$ , such that  $d_1h(n) \leq g(n) \leq d_2h(n)$ .

## 4 7.1 Problem 16

**Prove that  $4 \mid (3^{2n-1} + 1)$  for every positive integer  $n$ .**

## 5 7.2 Problem 8

**Prove that every prime except one has the form  $a^2 - b^2$  for some positive integers  $a$  and  $b$ .**