



Example $G = C_3 * C_2$, ($C_m := \mathbb{Z}/m\mathbb{Z}$)

What is the associated tree X ?

$$C_3 = \langle a \mid a^3 \rangle \quad C_2 = \langle b \mid b^2 \rangle$$

$$g C_3 \in X^0 \quad \text{st}(g C_3) = \{ h \{1\} \mid h \{1\} = g \{1\} \} \\ = g C_3$$

$$g C_2 \in X^1 \quad \text{st}(g C_2) = g C_2.$$



a acts on elliptic elements "rotating" around C_3 .

b acts on elliptic elements "rotating" around C_2 .

ab ? Hyperbolic, with axis passing through

$\dots ba^2C_2, bC_3, C_2, C_3, aC_2, abC_3, abaC_2, ababC_3, \dots$
translation length: 2.

EX

G group. $f: \mathbb{Z} * G \rightarrow \mathbb{Z}$ induced by id and trivial.

Then, $\text{Ker } f = \bigstar_{m \in \mathbb{Z}} t^m G t^{-m}$.

$SL_2(\mathbb{Z})$

$$SL_2(\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, \quad ad - bc = 1 \right\}.$$

Who knows a bit of hyperbolic or complex geometry, knows how $SL_2(\mathbb{Z}) < SL_2(\mathbb{R})$ acts on $\{z \in \mathbb{C} \mid \operatorname{Im} z > 0\}$.

One can find a fundamental domain and see that its translates are arranged in a tree-like fashion, from which one finds an action of $SL_2(\mathbb{Z})$ on a tree. Let us see another perspective (to obtain the same thing).

FAREY BINARY SEARCH

$$L_0, R_0 \in \mathbb{R} \quad L_0 < R_0$$

$$U \in (L_0, R_0) \quad \text{"unknown"}.$$

$$\text{Standard binary search: } M_0 := \frac{L_0 + R_0}{2}.$$

$$\text{Ask if } U = M_0 \text{ or } U \in (L_0, M_0] \text{ or } U \in (M_0, R_0);$$

$$\text{iterate with } (L_1, R_1) = (L_0, M_0) \text{ or } (M_0, R_0) \dots$$

FAREY binary search:

start with rational range $L_0 = \frac{a}{b}$, $R_0 = \frac{c}{d} \in \mathbb{Q}$, reduced fractions with $b, d > 0$.

$$\text{To split the searching range, pick } M_0 = \frac{a+c}{b+d} \in \mathbb{Q}$$

and proceed analogously.

"**MEDIAN**"

$$\text{NOTE: } \frac{a}{b} < \frac{a+c}{b+d} < \frac{c}{d}.$$

Exercise: For every $U \in \mathbb{Q} \cap (L_0, R_0)$

the procedure eventually arrives at $M_K = U$ for a certain $K \in \mathbb{N}$.

$$\left[\text{Hint: if } 0 < \frac{a}{b} < \frac{x}{y} < \frac{c}{d} \text{ (reduced fractions)}, \text{ then } (bc - ad)(x + y) \geq a + b + c + d. \right]$$

For our purposes, it is convenient to extend to $P^1\mathbb{Q} = \mathbb{Q} \cup \{\infty\}$.

Def Given $x, y \in P^1\mathbb{Q}$ distinct, they have TWO MEDIANTS:

write $x = \frac{a}{b}$, $y = \frac{c}{d}$ as reduced integer fractions. Then,
the mediant is: $\frac{a+c}{b+d}$, $\frac{a-c}{b-d}$. i.e.: $a, b \in \mathbb{Z}$, $\gcd(a, b) = 1$.

Def Given $x, y \in \mathbb{P}^1 \mathbb{Q}$, write $\Delta(x, y) = \left| \det \begin{pmatrix} a & c \\ b & d \end{pmatrix} \right|$
if $x = \frac{a}{b}$, $y = \frac{c}{d}$ as reduced integer fractions.

Lemma If $\Delta(x, y) = 1$ and z is a mediant of x and y ,
then: $\Delta(x, z) = \Delta(y, z) = 1$; x is a mediant of y and z .

Pf: $x = \frac{a}{b}$, $y = \frac{c}{d}$ reduced fractions. $z = \frac{a+c}{b+d}$.

$$\left| \det \begin{pmatrix} a & a+c \\ b & b+d \end{pmatrix} \right| = \left| \det \begin{pmatrix} a+c & a+c \\ b+d & b+d \end{pmatrix} \right| = \Delta(x, y) = 1.$$

In particular $a+c$, $b+d$ are coprime, so $\Delta(x, z) = 1$


$$\text{and } \frac{a}{b} = \frac{(-c) + (a+c)}{(-d) + (b+d)}$$

□

Procedure: Start from the triple $\{0, 1, \infty\} \subset \mathbb{P}^1 \mathbb{Q}$; notice
that $\Delta(0, \infty) = 1$ and that 1 is a mediant of 0 and ∞ ;
at each step, replace one of the elements by the other
mediant of the remaining two.

obs: at any instant we have $\{x, y, z\}$ s.t. $\underbrace{1 = \Delta(x, y) = \Delta(y, z) = \Delta(x, z)}_{(*)}$
and each member is a
mediant of the other two.

We obtain a tree
of possible outcomes!



Def $SL_2(\mathbb{Z}) \curvearrowright \mathbb{P}^1\mathbb{Q}$ as $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \frac{p}{q} = \frac{ap+cq}{bp+dq}$.

obs : If $p, q \in \mathbb{Z}$, then $\gcd(p, q) = \gcd(ap+cq, bp+dq)$

obs Let $M \in SL_2(\mathbb{Z})$, $x, y \in \mathbb{P}^1\mathbb{Q}$. Then, $\Delta(x, y) = \Delta(Mx, My)$.

Lemma If $x, y \in \mathbb{P}^1\mathbb{Q}$ and $\Delta(x, y) = 1$, then $\exists M \in SL_2(\mathbb{Z})$
s.t. $Mx = \infty$, $My = 0$.

Pf: $x = \frac{a}{b}$, $y = \frac{c}{d}$, $\det \begin{pmatrix} a & c \\ b & d \end{pmatrix} = 1$.

Take $M = \begin{pmatrix} a & c \\ b & d \end{pmatrix}^{-1}$. □

Lemma If $\Delta(x, y) = 1$, then there are exactly 2 points $z \in \mathbb{P}^1\mathbb{Q}$
such that $\Delta(x, z) = \Delta(y, z) = 1$: the mediants of
 x and y .

Pf: We know that the mediants satisfy the condition.

Enough to show: there are only two points satisfying the cond.

If $x = \infty = \frac{1}{0}$ and $y = 0 = \frac{1}{0}$, the conditions are:

$$\det \begin{pmatrix} 1 & a \\ 0 & b \end{pmatrix} = \pm 1, \quad \det \begin{pmatrix} 0 & a \\ 1 & b \end{pmatrix} = \pm 1, \quad \text{which imply}$$

$$\frac{a}{b} \in \{\pm 1\}.$$

For generic x, y : let $M \in SL_2(\mathbb{Z})$ be s.t. $Mx = \infty$, $My = 0$.

Then, z satisfies $\Delta(x, z) = \Delta(y, z) = 1$, which implies

$$\Delta(Mx, Mz) = \Delta(My, Mz) = 1,$$

and there are only 2 possibilities for Mz . □

Cor Let $x, y \in \mathbb{P}^1\mathbb{Q}$ with $\Delta(x, y) = 1$. Let $M \in SL_2(\mathbb{Z})$, $z \in \mathbb{P}^1\mathbb{Q}$.

$$\begin{array}{lcl} z \text{ is a mediant of } x \text{ and } y & \Leftrightarrow & \Delta(x, z) = \Delta(y, z) = 1 \\ Mz \text{ --- --- --- } Mx \text{ and } My & \Leftrightarrow & \Delta(Mx, Mz) = \Delta(My, Mz) = 1 \end{array}$$

Lemme Let $x, y, z \in \mathbb{P}^1 \mathbb{Q}$ satisfy $(*)$. Then, $\{x, y, z\}$ is reachable starting from $\{0, 1, \infty\}$.

Pf: First, suppose $x = \infty$. Then, $(*)$ implies that $\{\infty, y, z\} = \{\infty, n, n+1\}$ for some $n \in \mathbb{Z}$.

They are all reachable:

$$\{\infty, n-1, n\} \longleftrightarrow \{\infty, n, n+1\}$$

because the mediant of ∞ and $n \in \mathbb{Z}$ are $n-1$ and $n+1$.

Let $\{x, y, z\}$ be a triple satisfying $(*)$.

We know from the exercise that \exists reachable triple containing, e.g., x . Say, $\{x, y', z'\}$.

Let $M \in SL_2(\mathbb{Z})$ s.t. $Mx = \infty$.

Then, $\{Mx, My', Mz'\}$ and $\{Mx, My, Mz\}$ satisfy $(*)$ and contain ∞ , so they are of the form

$$\{\infty, n', n'+1\} \quad \{\infty, n, n+1\}$$

these are "connected" by a sequence of steps, so the same is true for $\{x, y, z\}$ and $\{x, y', z'\}$, by the previous corollary. \square

Conclusion: we have a tree (the FAREY TREE) X with:

$$X^0 = \{\{x, y, z\} \subseteq \mathbb{P}^1 \mathbb{Q} \text{ satisfying } (*)\}$$

$$X^1 = \{(\{x, y, z\}, z) : \{x, y, z\} \in X^0\}$$

$$\swarrow \alpha$$

$$\{x, y, z\}$$

$$\searrow \omega$$

$$\{x, y, z' \text{ the other mediant}\}$$

$SL_2(\mathbb{Z}) \curvearrowright X$ from the action on $\mathbb{P}^1 \mathbb{Q}$.

Lemma: $SL_2(\mathbb{Z}) \curvearrowright X^1$ is transitive.

Pf: Let $(\{x, y, z\}, z) \in X^1$ we can write:

$$x = \frac{a}{b}, \quad y = \frac{c}{d}, \quad z = \frac{a+c}{b+d}.$$

Up to swapping x, y , we have $M = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \in SL_2(\mathbb{Z})$.

Then, $M(\{0, \infty, 1\}, 1) = (\{x, y, z\}, z)$. \square

obs In particular, $SL_2(\mathbb{Z}) \curvearrowright X^0$ is transitive, and there are inversions!

$Z :=$ barycentric subdivision of X .

Then, $SL_2(\mathbb{Z}) \curvearrowright Z$ with segment quotient!

Fundamental domain:

$$\begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \{0, \infty, 1\} & \gamma & (\{0, \infty, 1\}, 1) \sim (\{0, \infty, -1\}, -1) \end{array}$$

Stabilizers?

• Of $\{0, \infty, 1\}$: $M \in SL_2(\mathbb{Z})$ must permute $0, \infty, 1$.

$$\left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \pm \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}, \quad \pm \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \right\} \cong \mathbb{Z}/6\mathbb{Z}$$

$\infty \xrightarrow{0 \rightarrow 1}$ $\infty \xleftarrow{0 \leftarrow 1}$

• Of $(\{0, \infty, 1\}, 1) \sim (\{0, \infty, -1\}, -1)$

$$\left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \pm \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right\} \cong \mathbb{Z}/4\mathbb{Z}$$

$0 \leftrightarrow \infty \quad 1 \leftrightarrow -1$

• Stabilizer of γ : the intersection: $\left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} \cong \mathbb{Z}/2\mathbb{Z}$.

$$\Rightarrow SL_2(\mathbb{Z}) \cong \mathbb{Z}/6\mathbb{Z} \rtimes_{\mathbb{Z}/2\mathbb{Z}} \mathbb{Z}/4\mathbb{Z}$$

presentation: $\langle x, y \mid x^6, y^4, x^3 = y^2 \rangle$.

$$x \mapsto \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}, \quad y \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$