

1. Consider the following table of data points/nodal points:

| Time (sec) t | Velocity (ms ⁻¹) v(t) |
|-----------------|--------------------------------------|
| 2 | 10 |
| 4 | 20 |
| 6 | 25 |

- a) [4+1 marks] Find an interpolating polynomial of velocity that goes through the above data points by using **Vandermonde Matrix** method. Also compute an approximate value of acceleration at Time, t=7 sec.
 b) [4 marks] Find an interpolating polynomial of velocity that goes through the above data points by using **Lagrange method**.
 c) [1 mark] If a new data point is added in the above scenario, which method you should use in finding a new interpolating polynomial. Also what will be the degree of that new polynomial?

1. a) Here, $x_0 = 2, x_1 = 4, x_2 = 6 \therefore$ Degree will be 2

$$\text{Now, } V = \begin{bmatrix} 1 & 2^1 & 2^2 \\ 1 & 4^1 & 4^2 \\ 1 & 6^1 & 6^2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 1 & 4 & 16 \\ 1 & 6 & 36 \end{bmatrix}$$

$$Y = \begin{bmatrix} 10 \\ 20 \\ 25 \end{bmatrix}$$

We Know, $V \cdot A = Y$

$$\Rightarrow A = V^{-1} \cdot Y$$

$$\Rightarrow \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 1 & 4 & 16 \\ 1 & 6 & 36 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 10 \\ 20 \\ 25 \end{bmatrix}$$

$$= \begin{bmatrix} -5 \\ 35/4 \\ -5/8 \end{bmatrix}$$

$$\text{Now, } P_2(x) = a_0 + a_1 x^1 + a_2 x^2$$

$$= -5 + \frac{35}{4}x - \frac{5}{8}x^2$$

$$\text{Now, } v(t) = P_2(t) = -5 + \frac{35t}{4} - \frac{5t^2}{8}$$

$$= 25.625 \text{ ms}^{-1}$$

$$\therefore \text{Acceleration} = \frac{d}{dx}(P_2(x)) = \frac{35}{4} - \frac{5}{4}x \Rightarrow a(t) = \frac{35}{4} - \frac{5}{4}t = 0 \text{ ms}^{-2}$$

b) Using lagrange method,

$$P_2(x) = l_0(x)f(x_0) + l_1(x)f(x_1) + l_2(x)f(x_2)$$

$$\text{Now, } l_0(x) = \frac{x-x_1}{x_0-x_1} \times \frac{x-x_2}{x_0-x_2} = \frac{x-4}{2-4} \times \frac{x-6}{2-6} = \frac{1}{8}(x^2 - 10x + 24)$$

$$l_1(x) = \frac{x-x_0}{x_1-x_0} \times \frac{x-x_2}{x_1-x_2} = \frac{x-2}{4-2} \times \frac{x-6}{4-6} = -\frac{1}{4}(x^2 - 8x + 12)$$

$$l_2(x) = \frac{x-x_0}{x_2-x_0} \times \frac{x-x_1}{x_2-x_1} = \frac{x-2}{6-2} \times \frac{x-4}{6-4} = \frac{1}{8}(x^2 - 6x + 8)$$

$$\therefore P_2(x) = \frac{1}{8}(x^2 - 10x + 24) \times 10 - \frac{1}{4}(x^2 - 8x + 12) \times 20 + \frac{1}{8}(x^2 - 6x + 8) \times 25$$

$$= -\frac{5x^2}{8} + \frac{35x}{4} - 5$$

c) If new data point is added, Newton's divided difference method should be used. Degree of polynomial will then be 3.

c) If new data point is added, Newton's divided difference method should be used. Degree of polynomial will then be 3.

2. Read the following and answer accordingly:

- (a) (4 marks) Consider the nodes $[-\pi/2, 0, \pi/2]$. Find an interpolating polynomial of appropriate degree by using Newton's divided-difference method for $f(x) = x \sin(x)$.
- (b) (2 marks) Use the interpolating polynomial to find an approximate value at $\pi/4$, and compute the percentage relative error at $\pi/4$.
- (c) (4 marks) Add a new node π to the above nodes, and find the interpolating polynomial of appropriate degree.

2.(a) Given, $f(x) = x \sin x$

$$x_0 = -\frac{\pi}{2}$$

$$f(x_0) = -\frac{\pi}{2} \times \sin(-\frac{\pi}{2}) = \frac{\pi}{2}$$

$$x_1 = 0$$

$$f(x_1) = 0 \times \sin(0) = 0$$

$$x_2 = \frac{\pi}{2}$$

$$f(x_2) = \frac{\pi}{2} \times \sin(\frac{\pi}{2}) = \frac{\pi}{2}$$

Using Newton's Divided Difference form,

$$\begin{aligned} x_0 &= -\frac{\pi}{2} & f[x_0] &= \frac{\pi}{2} \\ x_1 &= 0 & f[x_0, x_1] &= \frac{f[x_1] - f[x_0]}{x_1 - x_0} = \frac{0 - \frac{\pi}{2}}{0 - (-\frac{\pi}{2})} = \frac{-\frac{\pi}{2}}{\frac{\pi}{2}} = -1 \\ x_2 &= \frac{\pi}{2} & f[x_1] &= 0 & f[x_1, x_2] &= \frac{f[x_2] - f[x_1]}{x_2 - x_1} = \frac{\frac{\pi}{2} - 0}{\frac{\pi}{2} - 0} = 1 \\ & & f[x_2] &= \frac{\pi}{2} & f[x_0, x_1, x_2] &= \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0} \\ & & & & &= \frac{1 - (-1)}{\frac{\pi}{2} - (-\frac{\pi}{2})} \\ & & & & &= \frac{1}{\pi} \end{aligned}$$

Since 3 nodes,

$$\begin{aligned} P_2(x) &= f[x_0] + f[x_0, x_1](x - x_0) + f[x_0, x_1, x_2](x - x_0)(x - x_1) \\ &= \frac{\pi}{2} + (-1)(x + \frac{\pi}{2}) + \frac{1}{\pi}(x + \frac{\pi}{2})(x - 0) \\ &= \frac{\pi}{2} - x - \frac{\pi}{2} + \frac{2}{\pi}(x + \frac{\pi}{2}x) \\ &= -\frac{1}{2}\pi + \frac{2}{\pi}x^2 + x = \frac{2}{\pi}x^2 \end{aligned}$$

(b) At $x = \frac{\pi}{4}$,

$$P_2(\frac{\pi}{4}) = \frac{2}{\pi} \times (\frac{\pi}{4})^2 = \frac{2}{\pi} \times \frac{\pi^2}{16} = \frac{\pi}{8}$$

$$\text{Now, } f(\frac{\pi}{4}) = \frac{\pi}{4} \times \sin\left(\frac{\pi}{4}\right) = \frac{\pi}{4} \times \frac{1}{\sqrt{2}} = \frac{\pi}{4\sqrt{2}}$$

$$\begin{aligned} \text{Percentage relative error} &= \left| \frac{f(\frac{\pi}{4}) - P(\frac{\pi}{4})}{f(\frac{\pi}{4})} \right| \times 100\% \\ &= 29.29\% \end{aligned}$$

(c) Adding new node x_3 ,

$$\begin{aligned} x_0 &= -\frac{\pi}{2} & f[x_0] &= \frac{\pi}{2} \\ x_1 &= 0 & f[x_0, x_1] &= \frac{f[x_1] - f[x_0]}{x_1 - x_0} = \frac{0 - \frac{\pi}{2}}{0 - (-\frac{\pi}{2})} = \frac{-\frac{\pi}{2}}{\frac{\pi}{2}} = -1 \\ x_2 &= \frac{\pi}{2} & f[x_1] &= 0 & f[x_0, x_1, x_2] &= \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0} = \frac{\frac{1}{\pi} - (-1)}{\frac{\pi}{2} - (-\frac{\pi}{2})} = \frac{2}{\pi} \\ x_3 &= \pi & f[x_2] &= \frac{\pi}{2} & f[x_1, x_2] &= \frac{f[x_2] - f[x_1]}{x_2 - x_1} = \frac{\frac{\pi}{2} - 0}{\frac{\pi}{2} - 0} = 1 \\ & & f[x_3] &= 0 & f[x_0, x_1, x_2, x_3] &= \frac{f[x_1, x_2, x_3] - f[x_0, x_1, x_2]}{x_3 - x_0} = \frac{-1 - \frac{2}{\pi}}{\pi - (-\frac{\pi}{2})} = \frac{-2}{\pi} \end{aligned}$$

$$f[x_0, x_1, x_2] = \frac{2}{\pi} \quad f[x_0, x_1, x_2, x_3] = -\frac{2}{\pi}$$

$$f[x_0, x_1, x_2, x_3] = \frac{f[x_1, x_2, x_3] - f[x_0, x_1, x_2]}{x_3 - x_0}$$

$$= \frac{-\frac{2}{\pi} - \frac{2}{\pi}}{\pi + \frac{\pi}{2}} = \frac{-\frac{4}{\pi}}{\frac{3\pi}{2}} = -\frac{8}{3\pi^2}$$

Since there are 4 nodes,

$$\begin{aligned} P_3(x) &= P_2(x) + f[x_0, x_1, x_2, x_3](x-x_0)(x-x_1)(x-x_2) \\ &= \frac{2}{\pi}x^2 + \left(-\frac{8}{3\pi^2}\right)(x+\frac{\pi}{2})(x-\theta)(x-\frac{\pi}{2}) \\ &= \frac{2}{\pi}x^2 - \frac{8}{3\pi^2}x\left(x^2 - \frac{\pi^2}{4}\right) \\ &= \frac{2}{\pi}x^2 - \frac{8}{3\pi^2}x^3 + \frac{8\pi^2}{12\pi^2}x \\ &= \frac{2}{\pi}x^2 - \frac{8}{3\pi^2}x^3 + \frac{2}{3}x \end{aligned}$$

