


## Nodes from a goal set

$$G_n \approx \{\text{nodes with distance } n \text{ from } G\}$$




$$G_0 := G$$

$$\{V\}_0 = \{V\}$$

## Nodes from a goal set

$$G_n \approx \{\text{nodes with distance } n \text{ from } G\}$$

$$G_{n+1} := \{s \mid (\exists s' \in G_n) \text{ arc}(s, s')\} - \bigcup_{i=1}^n G_i$$



$$G_0 := G$$


$$\{V\}_0 = \{V\}$$

$$\{V\}_1 = \{SA, NSW\}$$

## Nodes from a goal set

$$G_n \approx \{\text{nodes with distance } n \text{ from } G\}$$

$$G_{n+1} := \{s \mid (\exists s' \in G_n) \text{ arc}(s, s')\} - \bigcup_{i=1}^n G_i$$



$$G_0 := G$$

$$\{V\}_0 = \{V\}$$


$$\{V\}_1 = \{SA, NSW\}$$

$$\{V\}_2 = \{WA, NT, Q\}$$

## Nodes from a goal set

$$G_n \approx \{\text{nodes with distance } n \text{ from } G\}$$

$$G_{n+1} := \{s \mid (\exists s' \in G_n) \text{ arc}(s, s')\} - \bigcup_{i=1}^n G_i$$



$$G_0 := G$$

$$\{V\}_0 = \{V\}$$

$$\{V\}_1 = \{SA, NSW\}$$

$$\{V\}_2 = \{WA, NT, Q\}$$

$$\{V\}_\infty = \{T\}$$

## Distance $d_G$ to minimize

$$d_G(s) := \begin{cases} n & \text{if } s \in G_n \\ \infty & \text{otherwise} \end{cases}$$

Refine

$$\delta_G(s) := \begin{cases} 1 & \text{if } s \in G \\ 0 & \text{otherwise} \end{cases}$$

to reward from 1 to 0 ( $\approx$  distance from 0 to  $\infty$ )

Distance  $d_G$  to minimize  $\rightsquigarrow$  reward  $r_G$  to maximize

$$d_G(s) := \begin{cases} n & \text{if } s \in G_n \\ \infty & \text{otherwise} \end{cases}$$

Refine

$$\delta_G(s) := \begin{cases} 1 & \text{if } s \in G \\ 0 & \text{otherwise} \end{cases}$$

to reward from 1 to 0 ( $\approx$  distance from 0 to  $\infty$ )

$$r_G(s) := \begin{cases} 2^{-n} & \text{if } s \in G_n \\ 0 & \text{otherwise} \end{cases}$$

halving the reward as we step back (starting at  $G$ )

Distance  $d_G$  to minimize  $\rightsquigarrow$  reward  $r_G$  to maximize

$$d_G(s) := \begin{cases} n & \text{if } s \in G_n \\ \infty & \text{otherwise} \end{cases}$$

Refine

$$\delta_G(s) := \begin{cases} 1 & \text{if } s \in G \\ 0 & \text{otherwise} \end{cases}$$

to reward from 1 to 0 ( $\approx$  distance from 0 to  $\infty$ )

$$r_G(s) := \begin{cases} 2^{-n} & \text{if } s \in G_n \\ 0 & \text{otherwise} \end{cases}$$

halving the reward as we step back (starting at  $G$ )

$$r_G(s) = \frac{1}{2} r_G(s') \text{ if } \text{arc}(s, s') \text{ and } d_G(s') < d_G(s).$$

## Rewards looking ahead

$$H_0(s) := \delta_G(s)$$

$$H_{n+1}(s) := \delta_G(s) + \frac{1}{2} \max\{H_n(s') \mid arc_{=}(s, s')\}$$

where

$$arc_{=}(s, s') \iff arc(s, s') \text{ or } s = s'.$$



## Rewards looking ahead

$$H_0(s) := \delta_G(s)$$

$$H_{n+1}(s) := \delta_G(s) + \frac{1}{2} \max\{H_n(s') \mid \text{arc}_=(s, s')\}$$

where

$$\text{arc}_=(s, s') \iff \text{arc}(s, s') \text{ or } s = s'.$$

For  $s \in G_0$ ,

$$H_{n+1}(s) = 1 + \frac{1}{2} H_n(s) = a_{n+1}$$

where

$$a_n := \sum_{k=0}^n 2^{-k} = 2(1 - 2^{-(n+1)})$$

## Rewards looking ahead

$$H_0(s) := \delta_G(s)$$

$$H_{n+1}(s) := \delta_G(s) + \frac{1}{2} \max\{H_n(s') \mid \text{arc}_=(s, s')\}$$

where

$$\text{arc}_=(s, s') \iff \text{arc}(s, s') \text{ or } s = s'.$$

For  $s \in G_0$ ,

$$H_{n+1}(s) = 1 + \frac{1}{2}H_n(s) = a_{n+1}$$

where

$$a_n := \sum_{k=0}^n 2^{-k} = 2(1 - 2^{-(n+1)})$$

and for  $s \in G_{k+1}$ ,  $s' \in G_k$  and  $\text{arc}(s, s')$ ,

$$H_{k+n+1}(s) = \frac{1}{2}H_{k+n}(s')$$

Rewards looking ahead:  $\lim_{n \rightarrow \infty} H_n(s) = 2r_G(s)$

$$H_0(s) := \delta_G(s)$$

$$H_{n+1}(s) := \delta_G(s) + \frac{1}{2} \max\{H_n(s') \mid \text{arc}_=(s, s')\}$$

where

$$\text{arc}_=(s, s') \iff \text{arc}(s, s') \text{ or } s = s'.$$

For  $s \in G_0$ ,

$$H_{n+1}(s) = 1 + \frac{1}{2}H_n(s) = a_{n+1}$$

where

$$a_n := \sum_{k=0}^n 2^{-k} = 2(1 - 2^{-(n+1)})$$

and for  $s \in G_{k+1}$ ,  $s' \in G_k$  and  $\text{arc}(s, s')$ ,

$$H_{k+n+1}(s) = \frac{1}{2}H_{k+n}(s')$$

$$H = \lim_{n \rightarrow \infty} H_n$$

$$H(s) = \delta_G(s) + \frac{1}{2} \max\{H(s') \mid \text{arc}=(s, s')\}$$

a foolproof heuristic for the shortest solution

Frontier = [Hd|Tl] with  $H(\text{Hd}) \geq H(s)$  for all  $s$  in Tl.

$$H = \lim_{n \rightarrow \infty} H_n$$

$$H(s) = \delta_G(s) + \frac{1}{2} \max\{H(s') \mid \text{arc}=(s, s')\}$$

a foolproof heuristic for the shortest solution

Frontier = [Hd|Tl] with  $H(\text{Hd}) \geq H(s)$  for all  $s$  in Tl.

What if arcs have different costs?

$$H = \lim_{n \rightarrow \infty} H_n$$

$$H(s) = \delta_G(s) + \frac{1}{2} \max\{H(s') \mid \text{arc}=(s, s')\}$$

a foolproof heuristic for the shortest solution

Frontier = [Hd|Tl] with  $H(\text{Hd}) \geq H(s)$  for all  $s$  in Tl.

What if arcs have different costs?

Modify  $\delta_G(s)$  to

$$Q_0(s, s') := \begin{cases} 1 & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \\ -\max_{s_1, s_2} \text{cost}(s_1, s_2) & \text{otherwise} \end{cases}$$

and  $H_{n+1}(s)$  to

$$Q_{n+1}(s, s') := Q_0(s, s') + \frac{1}{2} \max\{Q_n(s', s'') \mid \text{arc}=(s', s'')\}$$

## Discounted rewards ( $0 \leq \gamma < 1$ )

(immediate) rewards  $r_1, r_2, r_3, \dots$  at times  $1, 2, 3, \dots$  give a  $\gamma$ -discounted value of

$$V := r_1 + \gamma r_2 + \gamma^2 r_3 + \dots = \sum_{i \geq 1} \gamma^{i-1} r_i$$

## Discounted rewards ( $0 \leq \gamma < 1$ )

(immediate) rewards  $r_1, r_2, r_3, \dots$  at times  $1, 2, 3, \dots$  give a  $\gamma$ -discounted value of

$$\begin{aligned} V &:= r_1 + \gamma r_2 + \gamma^2 r_3 + \dots = \sum_{i \geq 1} \gamma^{i-1} r_i \\ &= \left( \sum_{i=1}^t \gamma^{i-1} r_i \right) + \gamma^t V_{t+1} \end{aligned}$$

where  $V_t$  is the value from time step  $t$  on ( $V_1 = V$ )

$$V_t := \sum_{i \geq t} \gamma^{i-t} r_i = r_t + \gamma V_{t+1}$$



## Discounted rewards ( $0 \leq \gamma < 1$ )

(immediate) rewards  $r_1, r_2, r_3, \dots$  at times  $1, 2, 3, \dots$  give a  $\gamma$ -discounted value of

$$\begin{aligned} V &:= r_1 + \gamma r_2 + \gamma^2 r_3 + \dots = \sum_{i \geq 1} \gamma^{i-1} r_i \\ &= \left( \sum_{i=1}^t \gamma^{i-1} r_i \right) + \gamma^t V_{t+1} \end{aligned}$$

where  $V_t$  is the value from time step  $t$  on ( $V_1 = V$ )

$$V_t := \sum_{i \geq t} \gamma^{i-t} r_i = r_t + \gamma V_{t+1}$$

which is bound by bounds on  $r_i$

$$m \leq r_i \leq M \text{ for each } i \geq t \text{ implies } \frac{m}{1-\gamma} \leq V_t \leq \frac{M}{1-\gamma}$$

since  $\sum_{i \geq 0} \gamma^i = (1-\gamma)^{-1}$

## Discounted rewards ( $0 \leq \gamma < 1$ )

(immediate) rewards  $r_1, r_2, r_3, \dots$  at times  $1, 2, 3, \dots$  give a  $\gamma$ -discounted value of

$$\begin{aligned} V &:= r_1 + \gamma r_2 + \gamma^2 r_3 + \dots = \sum_{i \geq 1} \gamma^{i-1} r_i \\ &= \left( \sum_{i=1}^t \gamma^{i-1} r_i \right) + \gamma^t V_{t+1} \end{aligned}$$

where  $V_t$  is the value from time step  $t$  on ( $V_1 = V$ )

$$V_t := \sum_{i \geq t} \gamma^{i-t} r_i = r_t + \gamma V_{t+1} \quad (\text{backward induction})$$

which is bound by bounds on  $r_i$


$$m \leq r_i \leq M \quad \text{for each } i \geq t \quad \text{implies} \quad \frac{m}{1-\gamma} \leq V_t \leq \frac{M}{1-\gamma}$$

since  $\sum_{i \geq 0} \gamma^i = (1-\gamma)^{-1}$

$$Q = \lim_{n \rightarrow \infty} Q_n$$

$$Q(s, s') := Q_0(s, s') + \frac{1}{2} \max\{Q(s', s'') \mid \text{arc}=(s', s'')\}$$

$$Q_0(s, s') := \begin{cases} 1 & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \end{cases}$$



soln not chosen


$$Q(s, g) = -3$$

$$Q(s, a) = -2 = Q(a, b) = Q(b, s)$$

$$Q = \lim_{n \rightarrow \infty} Q_n$$

$$Q(s, s') := Q_0(s, s') + \frac{1}{2} \max\{Q(s', s'') \mid \text{arc}=(s', s'')\}$$


$$Q_0(s, s') := \begin{cases} 1 & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \end{cases}$$



soln not chosen

$$Q(s, g) = -3$$

$$Q(s, a) = -2 = Q(a, b) = Q(b, s)$$



costlier soln chosen

$$Q(s, g) = -5$$

$$Q(a, g) = -4 = Q(s, a)$$

## Upping the reward

Adjust  $Q_0(s, s')$  to

$$R(s, s') := \begin{cases} r & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \end{cases}$$

for some reward  $r$  high enough to offset costs of reaching a goal

## Upping the reward

Adjust  $Q_0(s, s')$  to

$$R(s, s') := \begin{cases} r & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \end{cases}$$

for some reward  $r$  high enough to offset costs of reaching a goal

$$\text{e.g. } r \geq 2^n nc$$

for solutions up to  $n - 1$  arcs with arcs costing  $\leq c$

e.g.  $n$  states, and  $c$  is max arc cost.

## Upping the reward

Adjust  $Q_0(s, s')$  to

$$R(s, s') := \begin{cases} r & \text{if } s = s' \in G \\ -\text{cost}(s, s') & \text{else if } \text{arc}(s, s') \end{cases}$$

for some reward  $r$  high enough to offset costs of reaching a goal

$$\text{e.g. } r \geq 2^n nc$$

for solutions up to  $n - 1$  arcs with arcs costing  $\leq c$

e.g.  $n$  states, and  $c$  is max arc cost.

Let

$$V(s) := \max\{Q(s, s') \mid \text{arc}=(s, s')\}$$

so for  $0 \leq i < n$ ,  $s' \in G_i$  and  $\text{arc}(s, s')$ ,

$$V(s') \geq 2^{n-i}(n-i)c$$

$$V(s) \geq -c + \frac{1}{2}V(s') \geq 2^{n-(i+1)}(n-(i+1))c \geq 2c.$$

## Recap

From node  $s$ , find path to goal via  $s'$  maximizing

$$Q(s, s') := R(s, s') + \frac{1}{2} V(s')$$

with discount  $\frac{1}{2}$  on future  $V(s')$ , contra

$$\begin{aligned} \text{cost}(s_1 \cdots s_k) &= \sum_{i=1}^{k-1} \text{cost}(s_i, s_{i+1}) \\ &\neq \sum_{i=1}^{k-1} 2^{1-i} \text{cost}(s_i, s_{i+1}) \end{aligned}$$



## Recap

From node  $s$ , find path to goal via  $s'$  maximizing

$$Q(s, s') := R(s, s') + \frac{1}{2} V(s')$$

with discount  $\frac{1}{2}$  on future  $V(s')$ , contra

$$\begin{aligned} \text{cost}(s_1 \cdots s_k) &= \sum_{i=1}^{k-1} \text{cost}(s_i, s_{i+1}) \\ &\neq \sum_{i=1}^{k-1} 2^{1-i} \text{cost}(s_i, s_{i+1}) \end{aligned}$$

Trade min cost guarantee for cost-benefit analysis with chance  $\frac{1}{2}$  of survival/doom.

## Recap

From node  $s$ , find path to goal via  $s'$  maximizing

$$Q(s, s') := R(s, s') + \frac{1}{2} V(s')$$

with discount  $\frac{1}{2}$  on future  $V(s')$ , contra

$$\begin{aligned} \text{cost}(s_1 \cdots s_k) &= \sum_{i=1}^{k-1} \text{cost}(s_i, s_{i+1}) \\ &\neq \sum_{i=1}^{k-1} 2^{1-i} \text{cost}(s_i, s_{i+1}) \end{aligned}$$

Trade min cost guarantee for cost-benefit analysis with chance  $\frac{1}{2}$  of survival/doom.

NEXT: more uncertainty, approached via approximations like

$$\begin{aligned} Q_n(s, s') &\approx Q(s, s') \text{ up to look ahead } n \\ Q(s, s') &= \lim_{n \rightarrow \infty} Q_n(s, s'). \end{aligned}$$