# Predictive Analysis of Malaysian Dengue Hemorrrhagic Fever Data from 2010 - 2017 using BigML

Foong M. Wong

December 17, 2019

#### Introduction


- Dengue Fever (DF)
- More than  $\frac{1}{3}$  of population affected by dengue (World Health Organization, 2017)



Infected Aedes mosquitoes

# Dengue Symptoms

- "Break-bone fever"
- Severe joint and muscle pain, rash, fever



Symptoms of Dengue Fever


#### Dengue Fever in Malaysia

- 2013: 43,436 dengue cases
- 2014: 108,698 dengue cases
- 2015: 120,836 dengue cases item Reported by Malaysian Ministry of Health in 2017

#### Difference between DF and DHF

- Dengue Fever (DF) V.S. Dengue Hemorrhagic Fever (DHF)
  - ► There is a slight difference.
  - ightharpoonup DF ightharpoonup DHF

#### Evolution of DF to DHF



#### Dengue Fever to Dengue Hemorrhagic Fever

Severe abdominal pain

Convulsions, bruising

Uncontrolled bleeding

Vomiting, diarrhea

Sudden high fever

Severe headache

Skin rash

Pain behind the eyes

Muscle, joint, bone pain

Circulatory system failure,

shock

Can be fatal

#### Research Goals

- Identify:
  - Infected areas/states
  - Age groups of dengue hemorrhagic fever
- Run time-series analysis:
  - Epidemic pattern
  - Future outbreaks

# Research Methodologies

- Data Collection
- ② Data Pre-processing
- Oata Loading
- Current data analysis
- Sun time-series analysis

#### Data Collection

#### Malaysian Open Data Portal



Dengue Hemorrhagic Dever data from 2010 - 2017 (Source: http://www.data.gov.my/)

# Data Pre-processing

Use tool for data cleaning and formatting




Illustration on Data Cleaning process

#### Data Pre-processing (cont.)


- Translate the variables names into English
- Transform to the right data format!

		BILANGAI	N KES PEN	YAKIT DEN	GUE HAEM	IORRHAG	C FEVER T	AHUNAN N	MENGIKUT	NEGERI D	AN KUMP	ULAN UMI	JR, TAHUN	2010		
KUMPULAN UMUR	JOHOR	КЕДАН	KELANTAN	MELAKA	NEGERI SEMBILAN	PAHANG	PERAK	PERLIS	PULAU PINANG	SABAH	SARAWAK	SELANGOR	TERENGGANU	WP KUALA LUMPUR	WP LABUAN	MALAYSIA
0-4	10	0	4	1	0	0	0	0	3	1	5	32	2	5	0	63
5-9	16	0	11	2	2	4	4	0	1	3	12	63	4	21	0	143
10-14	63	0	25	29	9	12	8	0	5	2	10	182	9	37	0	391
15-19	67	0	55	38	15	16	4	0	1	2	12	243	14	38	0	505
20-24	102	0	32	52	18	10	2	0	0	0	22	348	13	44	0	643
25-29	115	0	31	41	18	13	2	0	3	2	21	330	14	48	0	638
30-34	78	0	20	28	14	13	4	0	2	2	12	246	5	33	0	457
35-39	63	0	23	28	13	10	1	0	4	1	17	166	6	20	0	352
40-44	35	1	14	27	9	7	2	0	1	3	20	132	5	15	0	271
45-49	29	1	22	18	13	12	0	0	1	1	8	83	3	6	0	197
50-54	18	0	21	15	9	11	1	0	0	0	11	61	3	11	0	161
55-59	12	1	6	16	6	4	3	0	0	0	4	34	4	10	0	100
60-64	12	0	5	5	2	2	0	0	0	0	3	16	0	5	0	50
65-69	7	0	8	3	0	1	0	0	0	0	3	10	0	2	0	34
70-74	3	0	0	4	0	0	1	0	0	0	0	7	0	2	0	17
>75	0	0	2	1	0	1	0	0	0	0	2	2	1	0	0	9
Grand Total	630	3	279	308	128	116	32	0	21	17	162	1955	83	297	0	4031

Dengue Hemorrhagic Fever Dataset in 2010

### Data Pre-processing (cont.)


- Include cases from 13 states, remove cases from 2 federal territories
- Avoid duplicated cases (since 2 federal territories are part of the 13 states)



Dengue Hemorrhagic Fever 2010 Dataset


# Data Pre-processing (cont.)

- Remove the last column and last row
- Rename 'KUMPULAN UMUR' to 'AGE GROUP'



Dengue Hemorrhagic Fever 2010 Dataset

### Data Formatting (cont.)



Python script to create formatted datasets to be read into BigML


#### **Data Loading**

- Online machine learning service
- Easy to use




BigML: A machine learning platform

# Data Loading (cont.)



#### **Data Merging**



Merging 2010 - 2017 individual datasets

### Data Overview - Age Groups




18 / 30

# Data Overview - Age Groups (cont.)

- In South-East Asia, severe DHF cases are predominant among paediatric patients aged between 2 and 15 years (R. Bhatia, A.P. Dash, T. Sunyoto, 2013)
- In Malaysia, majority of the affected community are the age group of 13–35 years old (Sam et al, 2013)




#### Data Overview - 13 states



### Why Selangor has the highest number of DHF cases?

Selangor - Highest number of dengue fever cases



Hotspot for Dengue in Malaysia based on Website IDengue for community (Government Portal, 2016)

# Why Selangor has the highest number of DHF cases? (cont.)

- Selangor state recorded the highest number of dengue cases in Malaysia.
- Selangor is a state with a high population density in Malaysia compared to other states (Ghani et al, 2019)
  - Household cleanliness
  - Behavior of its citizens

# Comparison of Knowledge, Attitude, and Practice among Communities Living in Hotspot and Non-Hotspot Areas of Dengue (Ghani et al, 2019)

- Communities living in non-hotspot areas of dengue had better knowledge and attitude about dengue
- People are not clear about dengue treatment and people's behavior not keeping household clean



Filthy water in household area


# Comparison of Knowledge, Attitude, and Practice among Communities Living in Hotspot and Non-Hotspot Areas of Dengue (cont.)

- Low education, socioeconomic status and knowledge on dengue prevention
- Educational campaigns for those with lower income and education,
   Government prevention



#### Conclusion

- Used BigML to run data analysis to describe Dengue Hemorrhagic Fever Pattern over the past 8 years (2010-2017)
  - ► Age group (20-24)(Highest), 25-29 (Second Highest), >75 (Lowest)
  - ► Selangor (Highest), Johor (Second highest), Perlis (Lowest)



#### Future Improvements

- Use BigML/other software to forecast dengue hemorrhagic outbreaks using time series analysis
- Run predictive analysis on Dengue Death Cases Data

#### References

- Datasets of Number of Annual Dengue Hemorrhagic Fever (DHF) Cases by State and Age 2010 2017.
- BigML.com: A comprehensive Machine Learning platform for all predictive use cases. https://bigml.com/. 2019.
- Data Formatting Python Script https://github.com/foongminwong/dengue-analysis/blob/master/dengue-data.ipynb. 2019.
- R. Bhatia, A.P. Dash, T. Sunyoto Changing epidemiology of dengue in South-East Asia WHO South-East Asia J Public Health, 2 (1) (2013), pp. 23-27
- S.S. Sam, S.F. Omar, B.T. Teoh, J. Abd-Jamil, S. AbuBakar Review of dengue hemorrhagic fever fatal cases seen among adults: a retrospective study PLoS Negl Trop Dis, 7 (5)=(2013), p. e2194

# References (cont.)

- Jesse Alaniz. Most common types of mosquitoes in north america. "www.thebugmaster.com/ most-common-types-of-mosquitoes/", 2019.
- N.A. Muhammad Azami, S.A. Salleh, H.M. Neoh, S.Z. Syed Zakaria, R. Jamal Dengue epidemic in Malaysia: not a predominantly urban disease anymore BMC Res Notes, 4 (2011), p. 216
- Nurul Akmar Ghani ,Shamarina Shohaimi,Alvin Kah-Wei Hee,Hui-Yee Chee ,Oguntade Emmanuel and Lamidi Sarumoh Alaba Ajibola Comparison of Knowledge, Attitude, and Practice among Communities Living in Hotspot and Non-Hotspot Areas of Dengue in Selangor, Malaysia Trop. Med. Infect. Dis. (2019), p. 37
  - World Health Organization. World health organization. dengue and severe dengue.

# References (cont.)



- Malaysia Ministry of Health (MMOH). Hotspot locality list. idengue portal.

  "http://idengue.remotesensing.gov.my/idengue/page2.php?kandungan-
  - "http://idengue.remotesensing.gov.my/idengue/page2.php?kandungan=e2017.
- Rajesh Bhatia, Aditya P Dash, Temmy Sunyoto, et al. Changing epidemiology of dengue in south-east asia. WHO South-East Asia Journal of Public Health, 2(1):23, 2013.

Thank You!

Questions?