Supplement for: Carbon cycling in mature and regrowth forests globally: a macroecological synthesis based on the global Forest Carbon (ForC) database

Contents

6	Appendix S1. Duplicates and Conflicting Records within ForC	2
7	Table S1. Numbers of records by biome and age class	3
8	Table S2. Model parameter estimates for age trends and biome differences in young forests \dots	4
9	Figure S1. Age trends and biome differences for NEP	8
10	Figure S2. Age trends and biome differences for GPP	E
11	Figure S3. Age trends and biome differences for NPP	10
12	Figure S4. Age trends and biome differences for $ANPP$	11
13	Figure S5. Age trends and biome differences for $ANPP_{woody}$	12
14	Figure S6. Age trends and biome differences for $ANPP_{stem}$	13
15	Figure S7. Age trends and biome differences for $ANPP_{foliage}$	14
16	Figure S8. Age trends and biome differences for $ANPP_{litterfall}$	15
17	Figure S9. Age trends and biome differences for $BNPP$	16
18	Figure S10. Age trends and biome differences for $BNPP_{coarse}$	17
19	Figure S11. Age trends and biome differences for $BNPP_{fine}$	18
20	Figure S12. Age trends and biome differences for R_{eco}	19
21	Figure S13. Age trends and biome differences for R_{root}	20
22	Figure S14. Age trends and biome differences for R_{soil}	21
23	Figure S15. Age trends and biome differences for $R_{het-soil}$	22
24	Figure S16. Age trends and biome differences for B_{tot}	23
25	Figure S17. Age trends and biome differences for B_{ag}	24
26	Figure S18. Age trends and biome differences for $B_{ag-wood}$	25
27	Figure S19. Age trends and biome differences for $B_{foliage}$	26
28	Figure S20. Age trends and biome differences for B_{root}	27
29	Figure S21. Age trends and biome differences for $B_{root-coarse}$	28
30	Figure S22. Age trends and biome differences for $B_{root-fine}$	29
31	Figure S23. Age trends and biome differences for DW_{tot}	30
32	Figure S24. Age trends and biome differences for $DW_{standing}$	31
33	Figure S25. Age trends and biome differences for DW_{down}	
34	Figure S26. Age trends and biome differences for OL	33

35 Appendix S1. Duplicates and Conflicting Records within ForC

- 36 Status of duplicates and conflicting records within ForC
- 37 Generating ForC_simplified
- Replicate measurements (i.e., replicates from within a single study) were averaged. Records that subsumed
- $_{39}$ others— i.e., the time period included that of ≥ 2 other records or dates were unknown and therefore
- 40 conflicted with ≥ 2 other records—were removed. For each group of duplicate records—i.e., measurements of
- 41 the same variable in the same plot at the same time—one record was assigned precedence (recorded in
- 42 D.precedence field). When measurement periods overlapped or were not specified, precedence was given first
- 43 to records representing longer measurement periods (i.e., end.date start.date) and then to more recently
- 44 published values. We manually reviewed duplicates that differed only in methodology, assigning precedence
- to the record employing a more comprehensive approach (e.g., inclusion of understory, lianas, or bamboo as
- opposed to just trees) or using a favored methodology.

Table S1. Numbers of records by biome and age class

	n records		
Biome	Mature	Young	
Boreal climate zones			
Boreal broadleaf	3	420	
Boreal conifer	691	1548	
Boreal Other	53	335	
Excluded climate zones			
Other broadleaf	8	142	
Other conifer	42	89	
Other Other	6	41	
Temperate climate zones	,		
Temperate broadleaf	1015	3662	
Temperate conifer	878	3519	
Temperate Other	241	2761	
Tropical climate zones			
Tropical broadleaf	1337	1455	
Tropical conifer	3	18	
Tropical Other	8	2045	

⁴⁸ Sample sizes refer to data set after merging of duplicates, removal of stands with no age or history data, and

removal of managed and disturbed stands. For vegetation type, "Other" refers to stands that are mixed

 $_{50}$ broadleaf/ conifer or that have not been classified. Focal biomes are indicated in bold.

$_{\scriptscriptstyle{51}}$ Table S2. Model parameter estimates for age trends and biome differences in $_{\scriptscriptstyle{52}}$ young forests

Variable	Parameter	Estimate	SE	t_{value}
NEP				
NEP	log10(stand.age)	2.12	0.35	6.00
NEP	BiomeTropical broadleaf	1.56	1.60	0.97
NEP	BiomeTemperate broadleaf	-1.01	0.67	-1.51
NEP	BiomeTemperate conifer	-0.20	0.60	-0.33
NEP	BiomeBoreal conifer	-2.40	0.75	-3.22
GPP				
GPP	log10(stand.age)	4.27	1.00	4.28
GPP	BiomeTropical broadleaf	26.47	3.70	7.16
GPP	BiomeTemperate broadleaf	7.96	1.91	4.16
GPP	BiomeTemperate conifer	9.99	1.54	6.47
GPP	BiomeBoreal conifer	1.48	2.02	0.73
NPP				
NPP	$\log 10(\text{stand.age})$	0.82	0.48	1.71
NPP	BiomeTropical broadleaf	10.37	1.53	6.78
NPP	BiomeTemperate broadleaf	5.62	0.96	5.84
NPP	BiomeTemperate conifer	4.49	0.95	4.73
NPP	BiomeBoreal conifer	2.35	1.20	1.95
ANPP				
ANPP	log10(stand.age)	5.58	0.93	6.01
ANPP	BiomeTropical broadleaf	-1.42	1.53	-0.92
ANPP	BiomeTemperate broadleaf	2.28	0.98	2.34
ANPP	BiomeTemperate conifer	1.72	0.91	1.89
ANPP	BiomeBoreal conifer	0.90	1.21	0.74
ANPP	log10(stand.age):BiomeTemperate broadleaf	-4.09	1.09	-3.75
ANPP	log10(stand.age):BiomeTemperate conifer	-4.55	1.09	-4.19
ANPP	log10(stand.age):BiomeBoreal conifer	-4.68	1.16	-4.04
ANPP_woody				
$ANPP_{woody}$	$\log 10 ({\rm stand.age})$	1.62	0.53	3.07
$ANPP_{woody}$	BiomeTropical broadleaf	0.54	1.55	0.35
$ANPP_{woody}$	BiomeTemperate broadleaf	0.03	1.06	0.03
$ANPP_{woody}$	BiomeTemperate conifer	0.36	0.79	0.45
$ANPP_{woody}$	BiomeBoreal conifer	-1.03	0.97	-1.06
ANPP_stem				
$ANPP_{stem}$	$\log 10 ({ m stand.age})$	-0.03	0.80	-0.03
$ANPP_{stem}$	BiomeTropical broadleaf	2.38	1.24	1.92
$ANPP_{stem}$	BiomeTemperate broadleaf	2.86	0.66	4.33
$ANPP_{stem}$	BiomeTemperate conifer	-0.83	0.75	-1.10
$ANPP_{stem}$	BiomeBoreal conifer	-0.37	1.64	-0.23
$ANPP_{stem}$	$\log 10 ({\rm stand.age}) : {\rm BiomeTemperate~broadleaf}$	-0.03	0.90	-0.04
$ANPP_{stem}$	$\log 10 ({\rm stand.age})$:BiomeTemperate conifer	2.28	0.93	2.45
$ANPP_{stem}$	$\log 10 ({\rm stand.age})$:BiomeBoreal conifer	0.78	1.28	0.61
ANPP_branch				
$ANPP_{branch}$	$\log 10 ({ m stand.age})$	0.16	0.17	0.93
$ANPP_{branch}$	BiomeTemperate broadleaf	0.40	0.34	1.17
$ANPP_{branch}$	BiomeTemperate conifer	0.06	0.28	0.22
$ANPP_{branch}$	BiomeBoreal conifer	0.10	0.40	0.24

Variable	Parameter	Estimate	SE	t_{value}
$ANPP_{foliage}$	$\log 10 ({\rm stand.age})$	1.33	0.16	8.40
$ANPP_{foliage}$	BiomeTropical broadleaf	0.67	0.54	1.23
$ANPP_{foliage}$	BiomeTemperate broadleaf	-0.14	0.32	-0.43
$ANPP_{foliage}$	BiomeTemperate conifer	-0.48	0.30	-1.61
$ANPP_{foliage}$	BiomeBoreal conifer	-1.52	0.38	-3.98
$ANPP_{litterfall}$	log10(stand.age)	1.42	0.86	1.64
$ANPP_{litterfall}$	BiomeTropical broadleaf	1.85	1.34	1.37
$ANPP_{litterfall}$	BiomeTemperate broadleaf	-0.51	1.60	-0.32
$ANPP_{litterfall}$	BiomeTemperate conifer	-0.66	1.93	-0.34
$ANPP_{litterfall}$	BiomeBoreal conifer	-2.08	1.89	-1.10
$ANPP_{repro}$	$\log 10 ({\rm stand.age})$	1.42	0.86	1.64
$ANPP_{repro}$	BiomeTropical broadleaf	1.85	1.34	1.37
$ANPP_{repro}$	BiomeTemperate broadleaf	-0.51	1.60	-0.32
$ANPP_{repro}$	BiomeTemperate conifer	-0.66	1.93	-0.34
$ANPP_{repro}$	BiomeBoreal conifer	-2.08	1.89	-1.10
$ANPP_{folivory}$	log10(stand.age)	1.42	0.86	1.64
$ANPP_{folivory}$	BiomeTropical broadleaf	1.85	1.34	1.37
$ANPP_{folivory}$	BiomeTemperate broadleaf	-0.51	1.60	-0.32
$ANPP_{folivory}$	BiomeTemperate conifer	-0.66	1.93	-0.34
$ANPP_{folivory}$	BiomeBoreal conifer	-2.08	1.89	-1.10
M_{woody}	log10(stand.age)	1.42	0.86	1.64
M_{woody}	BiomeTropical broadleaf	1.85	1.34	1.37
M_{woody}	BiomeTemperate broadleaf	-0.51	1.60	-0.32
M_{woody}	BiomeTemperate conifer	-0.66	1.93	-0.34
M_{woody}	BiomeBoreal conifer	-2.08	1.89	-1.10
BNPP	$\log 10 ({\rm stand.age})$	0.84	0.25	3.41
BNPP	BiomeTropical broadleaf	3.05	0.70	4.37
BNPP	BiomeTemperate broadleaf	0.90	0.46	1.95
BNPP	BiomeTemperate conifer	0.84	0.45	1.85
BNPP	BiomeBoreal conifer	0.09	0.60	0.15
$BNPP_{coarse}$	$\log 10 ({\rm stand.age})$	0.09	0.09	1.00
$BNPP_{coarse}$	BiomeTropical broadleaf	0.31	0.39	0.80
$BNPP_{coarse}$	BiomeTemperate broadleaf	0.31	0.16	1.99
$BNPP_{coarse}$	BiomeTemperate conifer	0.59	0.15	3.92
$BNPP_{coarse}$	BiomeBoreal conifer	0.09	0.18	0.50
$BNPP_{fine}$	$\log 10 ({\rm stand.age})$	0.90	0.24	3.77
$BNPP_{fine}$	BiomeTropical broadleaf	3.24	0.54	5.97
$BNPP_{fine}$	BiomeTemperate broadleaf	0.04	0.40	0.10
$BNPP_{fine}$	BiomeTemperate conifer	0.21	0.39	0.54
$BNPP_{fine}$	BiomeBoreal conifer	-0.06	0.51	-0.12
R_{eco}	$\log 10 ({\rm stand.age})$	1.40	0.56	2.49
R_{eco}	BiomeTropical broadleaf	25.78	2.62	9.84
R_{eco}	BiomeTemperate broadleaf	9.63	1.19	8.08
R_{eco}	BiomeTemperate conifer	10.01	0.99	10.14
R_{eco}	BiomeBoreal conifer	5.09	1.37	3.71
R_{auto}	$\log 10 ({ m stand.age})$	1.40	0.56	2.49
R_{auto}	BiomeTropical broadleaf	25.78	2.62	9.84
R_{auto}	BiomeTemperate broadleaf	9.63	1.19	8.08
R_{auto}	BiomeTemperate conifer	10.01	0.99	10.14
R_{auto}	BiomeBoreal conifer	5.09	1.37	3.71
$R_{auto-ag}$	$\log 10 ({\rm stand.age})$	1.40	0.56	2.49
	BiomeTropical broadleaf	25.78	2.62	9.84
$R_{auto-ag}$	Bioine fropicar broadicar			
$R_{auto-ag}$ $R_{auto-ag}$	BiomeTemperate broadleaf	9.63	1.19	8.08
_		9.63 10.01	1.19 0.99	8.08 10.14

Variable	Parameter	Estimate	SE	t_{value}
R_{root}	$\log 10 ({ m stand.age})$	4.47	1.42	3.15
R_{root}	BiomeTropical broadleaf	2.98	1.68	1.78
R_{root}	BiomeTemperate broadleaf	2.60	1.21	2.15
R_{root}	BiomeTemperate conifer	2.91	0.92	3.14
R_{root}	BiomeBoreal conifer	0.89	1.81	0.49
R_{root}	$\log 10 ({\rm stand.age}) : {\rm BiomeTemperate~broadleaf}$	-3.65	1.61	-2.26
R_{root}	$\log 10 ({\rm stand.age})$:BiomeTemperate conifer	-4.03	1.54	-2.61
R_{root}	$\log 10 ({\rm stand.age})$:BiomeBoreal conifer	-4.07	1.75	-2.32
R_{soil}	$\log 10 ({ m stand.age})$	2.84	1.03	2.77
R_{soil}	BiomeTropical broadleaf	10.11	1.31	7.71
R_{soil}	BiomeTemperate broadleaf	7.23	0.73	9.85
R_{soil}	BiomeTemperate conifer	8.93	0.62	14.34
R_{soil}	BiomeBoreal conifer	4.56	1.17	3.89
R_{soil}	$\log 10 ({\rm stand.age}) : {\rm BiomeTemperate~broadleaf}$	-1.41	1.11	-1.27
R_{soil}	$\log 10 ({\rm stand.age}) : Biome Temperate\ conifer$	-3.33	1.09	-3.06
R_{soil}	$\log 10 ({\rm stand.age})$:BiomeBoreal conifer	-2.59	1.19	-2.17
$R_{het-soil}$	$\log 10 ({ m stand.age})$	0.32	0.36	0.90
$R_{het-soil}$	BiomeTropical broadleaf	5.45	0.69	7.95
$R_{het-soil}$	BiomeTemperate broadleaf	3.92	0.63	6.25
$R_{het-soil}$	BiomeTemperate conifer	3.73	0.58	6.44
$R_{het-soil}$	BiomeBoreal conifer	2.51	0.77	3.28
B_{tot}	$\log 10 ({ m stand.age})$	45.83	9.46	4.85
B_{tot}	BiomeTropical broadleaf	-0.49	19.16	-0.03
B_{tot}	BiomeTemperate broadleaf	2.83	35.92	0.08
B_{tot}	BiomeTemperate conifer	-18.87	23.78	-0.79
B_{tot}	BiomeBoreal conifer	-116.99	36.60	-3.20
B_{tot}	$\log 10 ({\rm stand.age})$: Biome Temperate broadleaf	17.12	22.24	0.77
B_{tot}	log10(stand.age):BiomeTemperate conifer	24.21	16.52	1.47
B_{tot}	log10(stand.age):BiomeBoreal conifer	52.70	21.34	2.47
B_{ag}	$\log 10 ({\rm stand.age})$	53.45	2.71	19.69
B_{ag}	BiomeTropical broadleaf	-13.27	5.57	-2.38
B_{ag}	BiomeTemperate broadleaf	-48.27	5.60	-8.62
B_{ag}	BiomeTemperate conifer	-58.38	5.56	-10.51
B_{ag}	BiomeBoreal conifer	-46.06	12.55	-3.67
B_{ag}	log10(stand.age):BiomeTemperate broadleaf	11.42	4.29	2.66
B_{ag}	log10(stand.age):BiomeTemperate conifer	13.66	4.31	3.17
B_{ag}	log10(stand.age):BiomeBoreal conifer	-3.97	7.37	-0.54
$B_{ag-wood}$	log10(stand.age)	43.04	19.12	2.25
$B_{ag-wood}$	BiomeTropical broadleaf	-9.96	31.58	-0.32
$B_{ag-wood}$	BiomeTemperate broadleaf	-49.19	40.47	-1.22
$B_{ag-wood}$	BiomeTemperate conifer	-180.05	30.71	-5.86
$B_{ag-wood}$	BiomeBoreal conifer	-89.11	62.50	-1.43
$B_{ag-wood}$	log10(stand.age):BiomeTemperate broadleaf	37.51	32.42	1.16
$B_{ag-wood}$	log10(stand.age):BiomeTemperate conifer	131.73	27.05	4.87
$B_{ag-wood}$	log10(stand.age):BiomeBoreal conifer	29.80	39.34	0.76
$B_{foliage}$	log10(stand.age)	2.42	0.34	7.23
$B_{foliage}$	BiomeTropical broadleaf	1.45	0.81	1.79
$B_{foliage}$	BiomeTemperate broadleaf	0.93	0.77	1.22
$B_{foliage}$	BiomeTemperate conifer	-4.39	1.16	-3.77
	BiomeBoreal conifer	-2.10	2.04	-1.03
		2.10	2.01	
$B_{foliage}$		-1.69	0.51	_3 31
$B_{foliage} \ B_{foliage}$	$\log 10 ({\rm stand.age}) : {\rm BiomeTemperate~broadleaf}$	-1.69 3.79	0.51	
$B_{foliage} \ B_{foliage} \ B_{foliage}$	$\label{eq:condition} $\log 10({\rm stand.age})$: BiomeTemperate broadleaf \\ \log 10({\rm stand.age})$: BiomeTemperate conifer$	3.79	0.77	4.94
$B_{foliage} \ B_{foliage}$	$\log 10 ({\rm stand.age}) : {\rm BiomeTemperate~broadleaf}$			-3.31 4.94 0.11 9.62

Variable	Parameter	Estimate	SE	t_{value}
B_{root}	BiomeTemperate broadleaf	-6.25	1.95	-3.21
B_{root}	BiomeTemperate conifer	-9.75	1.74	-5.61
B_{root}	BiomeBoreal conifer	-10.91	4.19	-2.60
B_{root}	$\log 10 ({\rm stand.age}) : Biome Temperate\ broadleaf$	1.33	1.62	0.82
B_{root}	$\log 10 ({\rm stand.age})$:BiomeTemperate conifer	1.08	1.54	0.70
B_{root}	log10(stand.age):BiomeBoreal conifer	2.08	2.56	0.81
$B_{root-coarse}$	log10(stand.age)	9.76	5.25	1.86
$B_{root-coarse}$	BiomeTropical broadleaf	-4.86	8.62	-0.56
$B_{root-coarse}$	BiomeTemperate broadleaf	0.13	6.68	0.02
$B_{root-coarse}$	BiomeTemperate conifer	-39.75	7.29	-5.45
$B_{root-coarse}$	BiomeBoreal conifer	-20.38	12.40	-1.64
$B_{root-coarse}$	log10(stand.age):BiomeTemperate broadleaf	-1.29	6.69	-0.19
$B_{root-coarse}$	log10(stand.age):BiomeTemperate conifer	28.58	6.96	4.11
$B_{root-coarse}$	log10(stand.age):BiomeBoreal conifer	5.93	8.08	0.73
$B_{root-fine}$	log10(stand.age)	-0.07	0.19	-0.38
$B_{root-fine}$	BiomeTropical broadleaf	2.80	2.44	1.15
$B_{root-fine}$	BiomeTemperate broadleaf	2.67	1.11	2.41
$B_{root-fine}$	BiomeTemperate conifer	2.72	1.12	2.43
$B_{root-fine}$	BiomeBoreal conifer	10.63	2.69	3.96
DW_{tot}	log10(stand.age)	1.89	0.97	1.95
DW_{tot}	BiomeTropical broadleaf	2.68	3.33	0.80
DW_{tot}	BiomeTemperate broadleaf	13.88	6.53	2.13
DW_{tot}	BiomeTemperate conifer	26.69	7.77	3.44
DW_{tot}	BiomeBoreal conifer	4.87	13.29	0.37
$DW_{standing}$	log10(stand.age)	5.07	1.10	4.62
$DW_{standing}$	BiomeTropical broadleaf	-5.09	46.76	-0.11
$DW_{standing}$	BiomeTemperate broadleaf	21.96	20.49	1.07
$DW_{standing}$	BiomeTemperate conifer	-5.64	24.65	-0.23
$DW_{standing}$	BiomeBoreal conifer	-6.64	33.11	-0.20
DW_{down}	log10(stand.age)	6.03	2.38	2.53
DW_{down}	BiomeTropical broadleaf	-2.22	5.48	-0.41
DW_{down}	BiomeTemperate broadleaf	15.16	12.00	1.26
DW_{down}	BiomeTemperate conifer	45.18	6.77	6.67
DW_{down}	BiomeBoreal conifer	45.21	11.04	4.10
DW_{down}	log10(stand.age):BiomeTemperate broadleaf	-12.19	7.93	-1.54
DW_{down}	log10(stand.age):BiomeTemperate conifer	-16.97	3.89	-4.36
DW_{down}	log10(stand.age):BiomeBoreal conifer	-25.53	7.25	-3.52
OL	$\log 10(\text{stand.age})$	0.58	1.05	0.55
OL	BiomeTropical broadleaf	3.24	3.42	0.95
OL	BiomeTemperate broadleaf	14.75	3.88	3.80
OL	BiomeTemperate conifer	1.49	3.23	0.46
OL	BiomeBoreal conifer	37.24	6.84	5.45
OL	log10(stand.age):BiomeTemperate broadleaf	-3.45	2.45	-1.41
OL	log10(stand.age):BiomeTemperate conifer	5.85	1.88	3.11
OL	log10(stand.age):BiomeBoreal conifer	1.34	4.57	0.29

Figure S1. Age trends and biome differences for NEP




Figure S1 | Age trends and biome differences for NEP. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

$_{54}$ Figure S2. Age trends and biome differences for GPP




Figure S2 | Age trends and biome differences for GPP. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S3. Age trends and biome differences for NPP




Figure S3 | Age trends and biome differences for NPP. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

$_{56}$ Figure S4. Age trends and biome differences for ANPP




Figure S4 | Age trends and biome differences for ANPP. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S5. Age trends and biome differences for $ANPP_{woody}$




Figure S5 | Age trends and biome differences for $ANPP_{woody}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S6. Age trends and biome differences for $ANPP_{stem}$




Figure S6 | Age trends and biome differences for $ANPP_{stem}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S7. Age trends and biome differences for $ANPP_{foliage}$




Figure S7 | Age trends and biome differences for $ANPP_{foliage}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifiant differences between biomes.

60 Figure S8. Age trends and biome differences for ANPP_{litterfall}




Figure S8 | Age trends and biome differences for $ANPP_{litterfall}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

$_{61}$ Figure S9. Age trends and biome differences for BNPP




Figure S9 | Age trends and biome differences for BNPP. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S10. Age trends and biome differences for $BNPP_{coarse}$




Figure S10 | Age trends and biome differences for $BNPP_{coarse}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

$_{63}$ Figure S11. Age trends and biome differences for $BNPP_{fine}$




Figure S11 | Age trends and biome differences for $BNPP_{fine}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifiant differences between biomes.

Figure S12. Age trends and biome differences for R_{eco}




Figure S12 | Age trends and biome differences for R_{eco} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S13. Age trends and biome differences for R_{root}




Figure S13 | Age trends and biome differences for R_{root} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S14. Age trends and biome differences for R_{soil}




Figure S14 | Age trends and biome differences for R_{soil} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S15. Age trends and biome differences for $R_{het-soil}$




Figure S15 | Age trends and biome differences for $R_{het-soil}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S16. Age trends and biome differences for B_{tot}




Figure S16 | Age trends and biome differences for B_{tot} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifiant differences between biomes.

Figure S17. Age trends and biome differences for B_{ag}




Figure S17 | Age trends and biome differences for B_{ag} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S18. Age trends and biome differences for $B_{ag-wood}$




Figure S18 | Age trends and biome differences for $B_{ag-wood}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S19. Age trends and biome differences for $B_{foliage}$




Figure S19 | Age trends and biome differences for $B_{foliage}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S20. Age trends and biome differences for B_{root}




Figure S20 | Age trends and biome differences for B_{root} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S21. Age trends and biome differences for $B_{root-coarse}$




Figure S21 | Age trends and biome differences for $B_{root-coarse}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S22. Age trends and biome differences for $B_{root-fine}$




Figure S22 | Age trends and biome differences for $B_{root-fine}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S23. Age trends and biome differences for DW_{tot}




Figure S23 | Age trends and biome differences for DW_{tot} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S24. Age trends and biome differences for $DW_{standing}$




Figure S24 | Age trends and biome differences for $DW_{standing}$. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S25. Age trends and biome differences for DW_{down}




Figure S25 | Age trends and biome differences for DW_{down} . Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.

Figure S26. Age trends and biome differences for OL




Figure S26 | Age trends and biome differences for OL. Map shows data sources (x and o indicate young and mature stands, respectively). Left plot shows age trends in forests up to 100 years old, as characterized by a linear mixed effects model with fixed effects of age and biome. Solid lines indicate significant effect of age, non-pareallel lines indicate a significant age x biome interaction. Boxplot illustrates distribution across mature forests, with different letters indicating signifant differences between biomes.