

Supplementary Information: Global patterns of forest autotrophic carbon fluxes

Rebecca Banbury Morgan Valentine Herrmann Norbert Kunert
Ben Bond-Lamberty Helene C. Muller-Landau Kristina J. Anderson-Teixeira

Contents

Table S1. Climate variable definitions, sources, and abbreviations	2
Table S2. Model summaries for each climate variable as a single fixed effect in models for each C flux	3
Table S3. Significance of relationships of forest C fluxes to MAT alone and in combination with MAP.	4
Table S4. Comparison of growing season length and MAT as predictors of forest C fluxes	5
Table S5. Best (lowest AIC) single-climate variable models by C flux.	6
Table S6. Pairwise comparisons of correlations with climate variables between C fluxes, with all analyses conducted on a set of sites common to each pair.	7
Figure S1. Sample distribution maps for the nine forest C fluxes analyzed	8
Figure S2. Correlations among climate variables and latitude	9
Figure S3. Ratios among forest C fluxes as a function of latitude and climate variables	10
Figure S4. Individual plots of forest C fluxes in relation to mean annual climate, part 1	11
Figure S5. Individual plots of forest C fluxes in relation to mean annual climate, part 2	12
Figure S6. Individual plots of forest C fluxes in relation to mean climate seasonality, part 1	13
Figure S7. Individual plots of forest C fluxes in relation to mean climate seasonality, part 2	14
Figure S8. Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 1	15
Figure S9. Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 2	16
References	17

Table S1. Climate variable definitions, sources, and abbreviations

Abbreviation	Climate variable	Units	Definition	Time span	Source
MAT	Mean annual temperature	°C	Annual mean temperature, from primary literature or WorldClim if not reported	1970 - 2000	Primary literature; WorldClim ¹
MAP	Mean annual precipitation	mm yr ⁻¹	Annual mean precipitation, from primary literature or WorldClim if not reported	1970 - 2000	Primary literature; WorldClim ¹
T Seas	Temperature seasonality	°C	Standard deviation (variation) of monthly temperature averages	1970 - 2000	WorldClim ¹
P Seas	Precipitation seasonality	%	Coefficient of variation of mean monthly precipitation x 100	1970 - 2000	WorldClim ¹
ATR	Annual temperature range	°C	Maximum temperature of warmest month - minimum temperature of coldest month	1970 - 2000	WorldClim ¹
Solar R	Solar radiation	kJ m ⁻² yr ⁻¹	Solar radiation	1970 - 2000	WorldClim2 ²
Cloud	Cloud cover	%	Cloud percentage cover	1901 - 2014	CRU time-series dataset v 4.03 ³
AFD	Annual frost days	days yr ⁻¹	Number of freeze days annually	1901 - 2014	CRU time-series dataset v 4.03 ³
AWD	Annual wet days	days yr ⁻¹	Number of days with precipitation >0.1 mm annually	1901 - 2014	CRU time-series dataset v 4.03 ³
PET	Potential evapotranspiration	mm yr ⁻¹	Mean annual potential evapotranspiration	1950 - 2000	Global Aridity Index and Potential Evapotranspiration Climate Database ⁴
AI	Aridity		MAP/mean annual PET	1950 - 2000	Global Aridity Index and Potential Evapotranspiration Climate Database ⁴
VPD	Vapour pressure deficit	kPa	Mean monthly vapour pressure deficit	1958 - 2015	TerraClimate ⁵
Max VPD	Maximum vapour pressure deficit	kPa	Maximum monthly vapour pressure deficit	1958 - 2017	Derived from TerraClimate data
WSM	Water stress months	months yr ⁻¹	Number of months annually with MAP < PET	1970 - 2000	Derived from WorldClim data
LGS	Length of growing season	months yr ⁻¹	Number of months annually with mean minimum temperature > 0.5°C	1901 - 2014	Derived from CRU data
gsT	Growing season temperature	°C	Mean growing season temperature	1901 - 2014	Derived from CRU data
gsP	Growing season precipitation	mm month ⁻¹	Mean monthly precipitation during growing season months	1901 - 2014	Derived from CRU data
gsPET	Growing season PET	mm month ⁻¹	Mean monthly potential evapotranspiration during growing season months	1901 - 2014	Derived from CRU data
gsR	Growing season solar radiation	mm month ⁻¹	Mean monthly solar radiation during growing season	1901 - 2014	Derived from WorldClim2 data

* The WorldClim version used was the most recent available at the time of analysis

¹ Hijmans et al. (2005) ² Fick and Hijmans. (2017) ³ Harris et al. (2017) ⁴ Trabucco and Zomer (2019) ⁵ Abatzoglou et al. (2018)

Table S2. Model summaries for each climate variable as a single fixed effect in models for each C flux

Carbon Flux	Latitude			MAT			MAP			T Seas			P Seas			ATR			Solar R			AI		
	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R-sq	dAIC
GPP	Lin	0.64	54.9	Lin	0.61	52.5	Lin	0.18	33.3	Poly	0.71	69.5	-	-	-	Poly	0.69	63.0	Log	0.16	8.9	-	-	-
NPP	Log	0.50	44.3	Lin	0.42	41.5	Poly	0.21	16.7	Log	0.52	44.3	-	-	-	Log	0.49	42.3	Poly	0.16	12.5	Lin	0.04	2.8
ANPP	Lin	0.44	63.4	Lin	0.44	80.5	Poly	0.16	19.7	Log	0.41	58.7	-	-	-	Log	0.37	51.9	Lin	0.11	12.3	Lin	0.05	2.1
ANPP stem	Lin	0.18	22.2	Lin	0.24	38.5	Log	0.05	7.3	Lin	0.14	17.6	Poly	0.05	5	Lin	0.12	13.6	Log	0.06	6.8	Lin	0.07	4
ANPP foliage	Lin	0.50	37.7	Lin	0.58	52.9	Poly	0.25	13.3	Lin	0.48	34.1	-	-	-	Lin	0.50	36.1	Log	0.17	10.1	Lin	0.11	6.8
BNPP root	Lin	0.34	22.9	Log	0.31	21.0	Poly	0.15	6.2	Log	0.36	26.6	-	-	-	Log	0.33	23.6	Poly	0.29	18.8	-	-	-
BNPP fine root	Lin	0.17	8.0	Lin	0.15	7.2	Log	0.11	5.4	Lin	0.17	8.4	-	-	-	Log	0.19	10.9	Log	0.14	7.2	Log	0.06	2.4
R auto	Lin	0.65	13.1	Lin	0.59	10.9	Poly	0.60	8.6	Log	0.65	13.1	-	-	-	Log	0.60	11.5	Log	0.27	2.4	Poly	0.48	3.7
R root	Log	0.22	8.8	Lin	0.24	8.3	Lin	0.15	6.8	Log	0.24	9.5	-	-	-	Log	0.22	8.8	-	-	-	Lin	0.16	7.3

Carbon Flux	Cloud			AFD			AWD			PET			VPD			Max VPD			WSM			LGS		
	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC	Model	R ²	Δ AIC
GPP	-	-	-	Log	0.54	50.0	Lin	0.11	5.7	Poly	0.36	19.7	Poly	0.31	15.9	-	-	-	-	-	-	Lin	0.53	38.2
NPP	Lin	0.06	3.6	Lin	0.40	38.5	Lin	0.11	7.3	Poly	0.32	24.3	Poly	0.18	15.3	-	-	-	Lin	0.04	4	Lin	0.38	28.4
ANPP	Poly	0.09	7.1	Log	0.41	61.6	Lin	0.17	18.7	Poly	0.27	24.5	Poly	0.23	21.4	Poly	0.06	2.2	Poly	0.06	3	Lin	0.34	44.0
ANPP stem	Poly	0.09	5.4	Log	0.17	22.3	-	-	-	Poly	0.20	14.0	Poly	0.21	17.7	Log	0.14	7.5	-	-	-	Log	0.11	12.6
ANPP foliage	-	-	-	Lin	0.53	43.4	Lin	0.15	7	Log	0.32	24.2	Log	0.35	30.0	Poly	0.07	4.9	Poly	0.17	7.8	Log	0.46	32.9
BNPP root	-	-	-	Lin	0.28	19.1	Poly	0.11	3.4	Poly	0.36	23.2	Poly	0.26	13.9	-	-	-	-	-	-	Lin	0.26	14.7
BNPP fine root	-	-	-	Lin	0.16	9.2	Lin	0.08	2.7	Log	0.14	7.1	Log	0.06	1.9	-	-	-	-	-	-	Lin	0.13	5.8
R auto	-	-	-	Log	0.57	9.4	Null	0.26	0.6	Log	0.36	4.8	Log	0.35	4.3	-	-	-	Null	0.3	1.5	Lin	0.47	5.8
R root	Log	0.16	1.9	Log	0.19	7.3	Lin	0.17	3.5	Poly	0.19	1.7	Poly	0.27	6.7	-	-	-	Lin	0.14	6.1	Lin	0.19	5.9

Model forms tested include first-order linear (Lin), second-order polynomial (Poly), and logarithmic (Log). Shown are models with lowest AIC value. ΔAIC value is calculated with respect to the null.

Table S3. Significance of relationships of forest C fluxes to MAT alone and in combination with MAP.

Carbon flux	MAT	MAT + MAP	MAT x MAP	Best model	R ²
GPP	<0.0001	<0.0001	NS	MAT + MAP	0.66
NPP	<0.0001	NS	0.018	MAT x MAP	0.48
ANPP	<0.0001	0.035	NS	MAT + MAP	0.45
ANPP stem	<0.0001	NS	0.021	MAT x MAP	0.26
ANPP foliage	<0.0001	NS	NS	MAT	0.59
BNPP root	<0.0001	NS	NS	MAT	0.29
BNPP fine root	0.0021	NS	NS	MAT	0.15
R auto	0.00016	0.041	NS	MAT + MAP	0.71
R root	0.0011	NS	NS	MAT	0.25

Table S4. Comparison of growing season length and MAT as predictors of forest C fluxes

Fixed effect	AIC value	Δ AIC relative to best model	Marginal R ²
GPP			
MAT	126.43	0.00	0.62
Growing season length	140.81	14.38	0.54
None	178.96	52.54	0.00
NPP			
MAT	174.88	0.00	0.52
Growing season length	191.54	16.65	0.40
None	216.17	41.29	0.00
ANPP			
MAT	249.51	0.00	0.29
Growing season length	254.21	4.70	0.26
None	268.94	19.43	0.00
ANPP stem			
MAT	235.96	0.00	0.15
Growing season length	237.29	1.33	0.14
None	243.14	7.18	0.00
ANPP foliage			
MAT	484.88	0.00	0.45
Growing season length	520.96	36.09	0.35
None	560.35	75.47	0.00
BNPP root			
MAT	184.54	0.00	0.59
Growing season length	204.93	20.38	0.46
None	237.47	52.92	0.00
BNPP fine root			
MAT	540.19	0.00	0.24
Growing season length	566.37	26.18	0.11
None	578.66	38.46	0.00
R auto			
MAT	45.26	0.00	0.63
Growing season length	50.36	5.10	0.50
None	56.17	10.91	0.00
R root			
MAT	133.54	0.00	0.25
Growing season length	135.93	2.39	0.20
None	141.79	8.25	0.00

Table S5. Best (lowest AIC) single-climate variable models by C flux.

Carbon flux	Climate variable	Model type	Δ AIC relative to null model	Δ AIC relative to next best model	R ²
GPP	T Seas	Poly	69.5	6.55	0.71
NPP	MAT	Lin	41.5	0.21	0.42
	T Seas	Log	44.3	-	0.52
ANPP	MAT	Lin	80.5	21.4	0.44
ANPP stem	MAT	Lin	38.5	15.87	0.24
ANPP foliage	MAT	Lin	52.9	11.05	0.58
BNPP root	T Seas	Log	26.6	3.01	0.36
BNPP fine root	ATR	Log	10.9	2.11	0.19
R auto	T Seas	Log	13.1	1.62	0.65
	ATR	Log	11.5	-	0.60
R root	T Seas	Log	9.5	0.76	0.24
	ATR	Log	8.8	-	0.22
	MAT	Lin	8.3	-	0.24

Table includes all models within Δ AIC ≤ 2.0 of the best model

Table S6. Pairwise comparisons of correlations with climate variables between C fluxes, with all analyses conducted on a set of sites common to each pair.

C flux variable 1	C flux variable 2	Climate variable	R ² variable 1	R ² variable 2	Model type variable 1	Model type variable 2	Number of plots	Variable with higher R ²
GPP	NPP	Latitude	0.62	0.66	Lin	Lin	37	NPP
		MAT	0.62	0.70	Log	Lin	37	NPP
		T Seas	0.65	0.70	Log	Log	37	NPP
NPP	ANPP	Latitude	0.52	0.48	Log	Log	158	NPP
		MAT	0.30	0.44	Log	Lin	158	ANPP
		T Seas	0.47	0.43	Lin	Lin	158	NPP
	BNPP	Latitude	0.49	0.34	Log	Lin	116	NPP
		MAT	0.41	0.22	Log	Log	116	NPP
		T Seas	0.49	0.41	Log	Log	116	NPP
ANPP	ANPP stem	Latitude	0.35	0.13	Lin	Lin	176	ANPP
		MAT	0.42	0.17	Lin	Lin	176	ANPP
		T Seas	0.29	0.09	Lin	Lin	176	ANPP
	ANPP foliage	Latitude	0.32	0.45	Log	Log	96	ANPP foliage
		MAT	0.36	0.50	Lin	Lin	96	ANPP foliage
		T Seas	0.27	0.42	Lin	Lin	96	ANPP foliage
GPP	R auto	Latitude	0.64	0.34	Null	Null	11	GPP
		MAT	0.69	0.34	Null	Null	11	GPP
		T Seas	0.64	0.32	Null	Null	11	GPP
BNPP	R root	Latitude	0.01	0.39	Null	Null	9	R root
		MAT	0.08	0.35	Null	Null	9	R root
		T Seas	0.01	0.63	Null	Null	9	R root

Figure S1. Sample distribution maps for the nine forest C fluxes analyzed




Figure S1: Maps showing distribution of samples for the nine forest C fluxes analyzed

Figure S2. Correlations among climate variables and latitude




Figure S2: Scatterplots and Pearson's R values for relationships among climate variables and latitude

Figure S3. Ratios among forest C fluxes as a function of latitude and climate variables




Figure S3: Ratios among forest C fluxes as a function of latitude and climate variables. Regressions test variation in allocation to component fluxes for three relationships: (1) $GPP = NPP + R_{auto}$, (2) $NPP = ANPP + BNPP$, and (3) $ANPP = ANPP_{foliage} + ANPP_{stem}$. We interpret analysis of variation in $NPP : R_{auto}$ in the context of CUE (GPP/NPP), as variation in CUE will be directly related to variation in $NPP : R_{auto}$.

Figure S4. Individual plots of forest C fluxes in relation to mean annual climate, part 1




Figure S4: Individual plots of forest C fluxes in relation to mean annual climate, part 1.

Figure S5. Individual plots of forest C fluxes in relation to mean annual climate, part 2




Figure S5: Individual plots of forest C fluxes in relation to mean annual climate, part 2.

Figure S6. Individual plots of forest C fluxes in relation to mean climate seasonality, part 1




Figure S6: Individual plots of forest C fluxes in relation to mean climate seasonality, part 1.

Figure S7. Individual plots of forest C fluxes in relation to mean climate seasonality, part 2




Figure S7: Individual plots of forest C fluxes in relation to mean climate seasonality, part 2.

Figure S8. Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 1




Figure S8: Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 1.

Figure S9. Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 2




Figure S9: Growing season length-standardized forest C fluxes in relation to mean growing season climate, part 2.

References

- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data*, 5(1), 170191. <https://doi.org/10.1038/sdata.2017.191>
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas: New climate surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302–4315. <https://doi.org/10.1002/joc.5086>
- Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset: Updated high-resolution grids of monthly climatic observations. *International Journal of Climatology*, 34(3), 623–642. <https://doi.org/10.1002/joc.3711>
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25(15), 1965–1978. <https://doi.org/10.1002/joc.1276>
- Trabucco, A., & Zomer, R. J. (2019). *Global Aridity Index and Potential Evapo-Transpiration (ET0) Climate Database v2*. 10. <https://doi.org/10.6084/m9.figshare.7504448.v3>