

TITULO

MARIA CAMILA REMOLINA-GUTIERREZ, JAIME E. FORERO-ROMERO, JUAN N. GARAVITO-CAMARGO
 Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia
Draft version November 15, 2014

ABSTRACT

MISSING: The abstract

Subject headings: Lyman Alpha Emission,MISSING....

1. INTRODUCTION

The Lyman-Alpha emission line is the spectral line produced when an electron goes from the second energy level to the first, loosing energy and emitting it as light. When this energy decays are seen in Hydrogen atoms, the wavelength of the Ly- α line is 121.567 nm. The detection of this emission line is a main technique in extragalactic astronomy because galaxies with a strong emission of this (called Lyman Alpha Emitters - LAEs) are used to study the evolution of the universe in big scale.

LAEs are young and far away galaxies with most of its matter composed by Hydrogen atoms, being the rest heavier elements, like Iron, produced by growing stars burning their material. Then, this spectral line tells very important information about the galaxy, most of it actually, which creates a motivation to find a way to model the Ly- α line according to certain free parameters that vary from a galaxy to another and that can be obtained from observations.

Our purpose in this paper is then to make a theoretical model, based on radiative transfer simulations, that use MonteCarlo computational methods to emulate how the line profile is going to come out of the galaxy if it is rotating and has a surrounding outflow, both of this phenomena characterized by key parameters that are going to be described further on the paper.

MISSING: What has been done related to this?

As it has been described, the main purpose of this project is to mix a rotating galaxy with an external outflow surrounding it and analyze the resulting spectrum. The rotation model for the LAE consists on a sphere with an homogeneous mixture of dust and hydrogen at a constant temperature undergoing a solid-body rotation. The spectrum then depends on the maximum velocity of the sphere, the neutral hydrogen optical depths and the viewing angle. What happens is that photons are

emitted from a central distribution of gas and escape the sphere after a radiative transfer process that ends at the border.

In this model, rotation does not induce any spatial anisotropy in the integrated line flux, the escape fraction or the average number of scatterings, which led to the creation of an analytic approximation for the galaxy spectrum. This mathematical expression helps to minimize the computational costs and lets try with several parameters without spending great amounts of time running the code.

After this photons escape the galaxy to empty space, they encounter a thin spherical shell of matter at a certain radius from the center. That material is the outflow of the LAE, created mainly by starbursts which composition is characterized with the parameters of Hydrogen column density and metallicity. Besides its components, the galactic outflow is also defined by an expanding constant velocity that affects the redshift of the photons.

As there is a lot of empty space inside the shell, it acts as a reflector of photons. This means, that when one wants to comes out it is most likely to bounce inside the sphere several times before finally escaping. However it is not likely that the photon comes back again inside the galaxy, which helps that the first process does not have to be repeated, and both stages can be treated as consecutive, not simultaneous.

2. THEORETICAL BACKGROUND

In this section we describe the two different models that are used in this project, that are able to reproduce a real and consistent Ly- α profile. As first we implement the rotation followed by an outflow.

MISSING:
 - Express prettier the intro of the section

We use the simplified rotation model developed by [garavito-camargo et al] in which a rotating galaxy is modeled as a solid rotating sphere, with a homogeneous mixture of hydrogen and dust. Photons can be initially at the center or can be homogeneously distributed inside the sphere. The equations governing this solid-body rotation sphere in which the axis of rotation is defined to be align with the z -axis are:

$$v_x = -\frac{y}{R}V_{\max}, \quad (1)$$

$$v_y = \frac{x}{R}V_{\max}, \quad (2)$$

$$v_z = 0, \quad (3)$$

Where R is the radius of the sphere and V_{\max} is the linear velocity at the sphere's surface. The minus/plus sign in the x/y -component of the velocity indicates the direction of rotation. In this case we take the angular velocity in the same direction as the \hat{k} unit vector.

In this work we use the analytical expression for rotation derived in [garavito-camargo et al] where a rotating sphere can be seen as a static sphere in the laboratory frame with a bulk velocity difference in each surface element with respect to a distant observer. With the previous analysis the outgoing spectra can be expressed as:

$$J(x, i) \approx 2\pi \int_0^R db \int_0^{2\pi} d\phi J(x, b, \phi, i), \quad (4)$$

Where $J(x, b, \phi, i)$ is the spectrum of the flux emerging from the surface at point (b, ϕ) and is expressed as:

$$J(x, b, \phi, i) = \frac{\sqrt{\pi}}{\sqrt{24}a\tau_0} \left(\frac{(x - x_b)^2}{1 + \cosh \left[\sqrt{\frac{2\pi^3}{27}} \frac{|(x - x_b)^3|}{a\tau_0} \right]} \right). \quad (5)$$

2.2. Outflow Model: ThinShell

MISSING:

- Important equations and description of the parameters
- Why we use ThinShell instead of Wind. Its advantages (explains redshift, avoids re-diffusion, bounces).
- Ref: Alvaro and Julian

2.3. Joint Model

MISSING:

- Full explanation.
- How exactly are this 2 mixed
- Tell and make a detailed explanation of the free parameters that we are going to explore.
- State the fixed parameters.




FIG. 1.— **Influence of Galaxy Viewing Angle:** The values of the fixed parameters are $v_{out} = 100 \text{ km s}^{-1}$ and $\log nH = 21.3125$. The 3 possible angles are shown in the plot with different colors. The increase is visible as well as its small enlargement factor.

3. RESULTS

Here goes the results....

MISSING:

- What are the compact results.
- Some plots.
- Not a physical analysis yet.
- Write better (prettier)

3.1. Influences of the Free Parameters

In order to study the influence of each of the three free parameters, we fix two of them and see how the final spectrum varies along the other one left. In each case we will state these changes.

3.1.1. Influence of the Galaxy Viewing Angle: θ_{gal}

If one sets fixed outflow v_{out} and $\log nH$ in each case the viewing angle has the same effect: it increases proportionally the intensity. However this change is not that significant. The resulting spectrums are completely the same, but enlarged vertically by a small factor. Fig. 1 helps visualize this effect in a better way.

3.1.2. Influence of the Hydrogen Column Density: $\log nH$

The effect of the $\log nH$ is the creation of 2 peaks: the left one very thin, tall and pronounced, and the right one very wide, small and soften. When the $\log nH$ is increased, the left peak starts to decrease while mixing with the right one, decreasing their height ratio until the left peak completely disappears. The resulting spectrum, with high column density, is a wide single mountain with intensity significantly less than at the beginning. Fig. 2 helps visualize this effect in a better way.

3.1.3. Influence of the Outflow Expanding Velocity: v_{out}

MISSING:

- Influence Outflow Velocity.




FIG. 2.— **Influence of Hydrogen Column Density:** The values of the fixed parameters are $v_{out} = 100 \text{ km s}^{-1}$ and $\theta_{gal} = 90^\circ$. There are three stages of the $\log nH$ value shown: initial, intermediate and final, with the values shown on the plot.

4. DISCUSSION

MISSING:

- Comparison with some other result (probably observations).
- Why is this result useful?
- What possible implications can this model have?

Ideas:

REFERENCES

Hunter, J. D. 2007, Computing In Science & Engineering, 9, 90

Pérez, F., & Granger, B. E. 2007, Computing in Science and Engineering, 9, 21

The decrease of the intensity while increasing the column density is caused because the column density is proportional to the absorption of light in the gas. The more $\log nH$, the less photons get out of the outflow.

5. CONCLUSIONS

Here goes the conclusions....

ACKNOWLEDGMENTS

To ...

The data, source code and instructions to replicate the results of this paper can be found here <https://github.com/mariacamilaremolinagutierrez/LymanAlpha/>. Most of our code benefits from the work of the IPython and Matplotlib communities (Pérez & Granger 2007; Hunter 2007).

MISSING:

- Help from Alvaro and Julian: data, explanations, advice and collaboration.
- Soooo many more acknowledgments.

MISSING IN REFERENCES:

- A looooot of references not here yet.
- Alvaro's paper.(Can galactic outflows explain the properties of Ly emitters? Alvaro Orsi, Cedric G. Lacey and Carlton M. Baugh)
- Use the correct citation method.