Supplementary Information: Tropical tree drought sensitivity is jointly shaped by drought characteristics, species adaptations, and individual microenvironments

Contents

Climate data and correlations
Data cleaning additional methods
Growth timeseries and anomalies
Drought year growth of species and individuals
Variables and their distributions
Conditional dependencies

Authors

Krishna Anujan 1,2* ORCiD: 0000-0003-3604-5895 Sean McMahon 2,3 ORCiD: 0000-0001-8302-6908 Sarayudh Bunyavejchewin 4 ORCiD: 0000-0002-1976-5041

Stuart J. Davies³ ORCiD: 0000-0002-8596-7522

 Helene C. Muller-Landau 3 ORCiD : 0000-0002-3526-9021 Nantachai Pongpattananurak 5 ORCiD: 0000-0001-8687-6182 Kristina Anderson-Teixeira 1,3 ORCiD: 0000-0001-8461-9713

Climate data and correlations




Figure S1: Raw climate data timeseries from 2009 to 2020 from the Forest Fire Research Station near HKK. Measurements represent daily rolling means over a 30-day time window for precipitation, max temperature and VPD and rolling sum for number of dry days.




Figure S2: Climate anomalies for all variables from 2009 to 2020from the Forest Fire Research Station near HKK. Values represent number of standard deviations from the long-term mean for daily rolling mean values.

${\bf Data\ cleaning\ additional\ methods}$

- Data cleaning stepsCalculating DBH equation
- Corroboration with tape measurements
- ullet error distribution

Growth timeseries and anomalies




Figure S3: Growth anomalies across the timeseries. Growth anomalies for each year calculated as the number of standard deviations in growth each year from mean growth across all years and summarised for plot and species.

Drought year growth of species and individuals




Figure S4: Species sensitivities in drought years 2010 and 2015. Boxplots represent 25th, 50th (median) and 75th percentile of sensitivity for each species in each year and whiskers represent 95% CIs. All individual sensitivities represented by jittered points.




Figure S5: Correlations of species sensitivities across the two years. Panel a shows correlation of species median sensitivities in the two years and Pearson's correlation coefficient. Colour gradient represents deciduousness scale and dotted line is the 1:1 line. Panel b shows difference in magnitude of species median sensitivites among the two years




Figure S6: Correlation of individual sensitivities across the two years along with Pearson's correlation coefficient and the 1:1 line.

Variables and their distributions

Map of trees





Figure S7: Map of trees with dendrobands in HKK, included in analyses in the 2010 and 2015 datasets



 $\textbf{Figure S8:} \ \, \textbf{Calculated Topographic Wetness Index (TWI) values for the HKK plot}$




Figure S9: Distribution of DBH with Crown Illumination Index (CII) in 2014.

Conditional dependencies

To analyse the influence of microenvironmental variables on growth sensitivity, we first created a Directed Acyclic Graph describing the relationships. To be able to test the DAG with the dataset, we used dagitty to derive the conditional dependencies that need to be tested, namely CII \perp TWI and DBH \perp TWI.




Figure S10: Testing correlations of variables across all individuals

There is low correlation between these two variable pairs across all individuals. We then tested conditional dependencies at the species level.

Most species (barring a few) had low correlation between these variables in our dataset, allowing us to proceed with analysis and interpretation.




Figure S11: Testing correlations for DBH \perp TWI by species




Figure S12: Testing correlations for CII $\perp\!\!\!\perp$ TWI by species