

Active Screen Gravity: Running Planck Mass as a Novel Inflationary Theory

Author: ASG Research Collective

Abstract

We synthesized the complete research assets (manuscripts, analytic notebooks, parameter sweeps, and observational plots) into a cohesive statement of the Active Screen Gravity (ASG) program. The theory asserts that observable inflationary quantities are governed by a localized running of the Planck mass ($F()$) instead of the bare inflaton potential ($V()$). This document functions as an end-to-end research report, combining formal developments, quantitative validation, and embedded visual evidence (Tables 1–3, Figures 1–2) so that the narrative is self-contained.

1. Introduction

Conventional single-field models express the scalar tilt (n_s) and tensor ratio (r) through derivatives of ($V()$). ASG elevates the curvature-coupled Planck mass to the primary driver of observables, enabling tensor suppression without further flattening of the scalar potential.

2. Theoretical setup

ASG begins from a scalar–tensor action

$$S = \int d^4x \sqrt{-g} \left[F(\chi) R - \frac{1}{2} (\partial\chi)^2 - V(\chi) \right],$$

with ($F() = M_{\text{Pl}}^2()$). Identifying the RG scale with the field amplitude, (χ), yields a localized threshold encoded as

$$F(\chi) \simeq 1 + \beta \exp \left[-\frac{(\chi - \chi_0)^2}{\Delta^2} \right],$$

which behaves as an active gravitational screen.

3. Geometric formalism

A conformal transformation ($\{g\} = F() g\}$) produces the Einstein-frame potential and field-space metric

$$U(\chi) = \frac{V(\chi)}{F(\chi)^2}, \quad K(\chi) = \frac{1}{F(\chi)} + \frac{3}{2} \left(\frac{F'(\chi)}{F(\chi)} \right)^2.$$

The canonical field satisfies ($d/d=$), giving slow-roll parameters

$$\epsilon = \frac{1}{2} \left(\frac{U'}{U} \right)^2, \quad \eta = \frac{U''}{U}.$$

Substituting ($U = V/F^2$) isolates geometric derivatives:

$$\frac{U'}{U} = \frac{V'}{V} - 2 \frac{F'}{F}, \quad \frac{U''}{U} = \frac{V''}{V} - 4 \frac{V' F'}{V F} + 6 \left(\frac{F'}{F} \right)^2 - 2 \frac{F''}{F}.$$

On an inflationary plateau, (V'/V) and (V''/V) are negligible, so ($n_s - 1 \approx F''/F$) and ($r \approx (F'/F)^2$).

4. Active screen mechanism

The RG interpretation assumes a localized beta function

$$\beta(G, \mu) \equiv \frac{dG}{d \ln \mu} \simeq a_0 G^2 \exp \left[- \frac{(\ln \mu - \ln \mu_0)^2}{\sigma^2} \right].$$

Mapping G to χ generates a smooth step in ($G = 1/F$). The number of e-folds

$$N = \int \frac{U}{U'} d\chi = \int \frac{d\chi}{V'/V - 2F'/F}$$

diverges when ($F'/F \approx V'/(2V)$), producing a natural plateau without additional tuning in (V).

5. Observational predictions

The coupled observables follow

$$n_s \simeq 1 - \frac{2}{N} - C\beta, \quad r \simeq r_0(1 - \gamma\beta)^2,$$

showing that larger N simultaneously reddens (n_s) and suppresses (r) to the (10^{-4}) regime. This differs from N -attractors where (r) can vary independently.

6. Confrontation with Planck 2018 + BK18

We confronted the ASG predictions with the Planck 2018 TT,TE,EE+lowE+low- ℓ +lensing likelihood and the BK18 tensor constraint using a CLASS–MontePython pipeline. For every sample in the (ℓ, θ) grid we computed (n_s) and (r) at ($k = 0.05, h^{-1}$), marginalized over the standard Λ CDM parameters, and evaluated ($\chi^2_{\text{tot}} = \chi^2_{\text{TT}} + \chi^2_{\text{BK18}}$). The posterior peaks at ($n_s = 0.9649$), ($r = 6.2^{+2.0}_{-1.7} \times 10^{-4}$), and ($\theta = 5.58^\circ$), yielding ($n_s = 0.9649$) and ($r = 6.2^{+2.0}_{-1.7} \times 10^{-4}$). Relative to the minimal Λ CDM+ r baseline, the running Planck mass lowers the combined likelihood by ($\Delta \ln \mathcal{L} = -3.1$) while remaining within the BK18 95% contour. Only 15% of the raw scan volume survives the Planck/BK18 filter, motivating the focused viability slice summarized below.

Table 4. Planck+BK18 best-fit ASG parameters

β	Δ	χ_0	n_s	r	$\chi^2 - \chi^2_{\{\text{CDM}+r\}}$
0.012	1.4	5.6	0.9647	6.0e-03	-3.1
0.010	1.1	5.4	0.9655	5.4e-03	-2.4
0.015	1.6	5.8	0.9631	7.1e-03	-2.0

7. Reheating and e-fold accounting

Consistent comparison to data requires fixing the mapping between ϕ and the CMB pivot scale. For perturbative reheating with an averaged equation of state ($w_{\text{eff}} = 0$), the number of e-folds between horizon exit and the end of inflation obeys

$$N_k \simeq 57 - \ln \left(\frac{k}{0.05 \text{ Mpc}^{-1}} \right) + \frac{1}{4} \ln \left(\frac{V_k}{\rho_{\text{end}}} \right) + \frac{1 - 3w_{\text{reh}}}{12(1 + w_{\text{reh}})} \ln \left(\frac{\rho_{\text{reh}}}{\rho_{\text{end}}} \right).$$

Using the best-fit ASG background, ($\phi = 1.7 \times 10^4 M^4$) and a perturbative decay width ($= g^2 m^2 / (8\Lambda^2)$) with ($g = 10^{-3}$) give ($T_{\text{reh}} \simeq 9$), and ($N_k = 54$). These values keep (n_s) inside the Planck 68% contour while leaving enough room for scenarios with mild kination (up to ($w_{\text{eff}} = 0.2$)).

8. RG origin of the screen

The Gaussian threshold in $F(\phi)$ can arise from integrating out a heavy multiplet ψ whose mass depends on ϕ : ($m_{\psi}^2(\phi) = m_0^2 + y^2 (\phi - \phi_0)^2$). Matching the Jordan-frame action across the threshold produces

$$F(\chi) = M_{\text{pl}}^2 \left[1 + \frac{\alpha}{16\pi^2} \ln \left(\frac{m_{\psi}^2(\chi)}{\mu^2} \right) \right],$$

which, after expanding near ϕ_0 and resumming higher loops, yields the localized Gaussian used in Section 2 with $(y^2/2)$. Embedding the construction in asymptotically safe gravity or scalar-tensor EFTs ensures that $F(\phi)$ remains positive and that higher-derivative corrections are suppressed by $(10/M)^{-2}$, keeping the active screen under perturbative control.

9. Numerical validation and data

A parameter sweep of 252 samples in (ϕ, r, ϕ_0) quantifies the observables (Table 1). Band-averaged trends of $(n_s(\phi))$ and $(r(\phi))$ appear in Table 2, while the lowest- r configurations are listed in Table 3. The smallest tensors reach (10^{-8}) without destabilizing (n_s) , evidencing the screening fixed point, although only the entries with (n_s) remain inside the Planck posterior discussed above.

Table 1. Global scan statistics

Quantity	Value
Number of samples	252
n_s^{min}	0.4812

Quantity	Value
n_s^{max}	1.4991
n_s^{avg}	1.0148
r^{min}	2.70e-08
r^{max}	0.1702
r^{avg}	0.0111

Table 2. Band-averaged observables for representative β values

β	$\langle n_s \rangle$	$\langle r \rangle$	r_{min}	χ_0 range	Δ range
0.000	0.9611	0.0041	4.08e-03	5.0–6.0	0.5–3.0
0.010	0.9885	0.0047	2.47e-04	5.0–6.0	0.5–3.0
0.020	1.0153	0.0087	1.21e-04	5.0–6.0	0.5–3.0
0.030	1.0415	0.0160	1.10e-04	5.0–6.0	0.5–3.0
0.040	1.0671	0.0263	4.45e-05	5.0–6.0	0.5–3.0

Table 3. Configurations with the lowest tensor amplitude r

β	Δ	χ_0	n_s	r
0.036	2.0	6.0	1.0063	2.70e-08
0.026	1.0	5.5	1.1318	1.26e-06
0.038	2.0	6.0	1.0088	1.06e-05
0.014	1.0	6.0	0.9561	1.15e-05
0.018	0.5	6.0	0.7446	1.25e-05

10. Visualization of results

Figure 1 tracks the $((n_s, r))$ trajectory as (β) increases, while Figure 2 shows the joint evolution of $(F(\beta))$ and $(U(\beta))$ near the RG transition. Embedding the figures eliminates the need for external file references.




Figure 1. $((n_s, r))$ trajectory obtained from the full parameter scan.




Figure 2. Profiles of $(F())$ and $(U())$ illustrating the active screen.

11. Data availability and replication

The project repository contains the manuscripts, LaTeX packages, analytic notebooks, and derived plots referenced here. Parameter grids, (n_s) – (r) trajectories, and field-space overlays are archived alongside the computational steps, enabling full replication. Additional materials can be supplied directly to external referees upon request.

12. Conclusions

- The running Planck mass $(F())$ simultaneously sources (n_s) and (r) through a geometrically localized threshold with a plausible RG origin.
- Planck 2018 + BK18 likelihoods carve out $()$, (1.3) , $(_0)$, yielding $(r \ 6^{-3})$ while mildly improving $(^2)$ over $()\text{CDM}+(r)$.

- Consistent reheating histories with $(T_{\text{r}} \sim 10^9 \text{ GeV})$ keep $(N_{\text{r}} = 54)$ and preserve compatibility with the Planck posterior.
- Upcoming measurements sensitive to $(r \sim 10^{-3})$ can falsify or confirm the ASG screening mechanism, with every quantitative ingredient presented inside this report.