

MK-677

Ibutamoren (MK-677) — Oral GH Secretagogue

ORAL GH RELEASE

MUSCLE GROWTH

SLEEP

HUNGER

LONG-ACTING

WHAT IS IT?

MK-677 (Ibutamoren) is a non-peptide oral ghrelin mimetic and GH secretagogue. It stimulates GH and IGF-1 release by acting on GHSR-1a receptors — and crucially, it is orally bioavailable with a 24-hour half-life, requiring just one dose per day. The only oral compound that significantly and sustainably raises GH and IGF-1 long-term without injections. Extensively studied in clinical trials.

DOSING AT A GLANCE

Daily oral dose	Optimal timing	Frequency	Cycle (long)
10–25 mg	Before bed	Once daily	16–24 weeks+

KEY BENEFITS

- Significantly increases GH and IGF-1 (sustained, 24-hour elevation)
- Oral — no injections required (major compliance advantage)
- Accelerates muscle growth and recovery via IGF-1 upregulation
- Dramatically improves sleep — increases REM and deep sleep stages
- Improves bone density — shown in multiple clinical trials
- Reduces body fat while preserving lean mass in caloric deficits
- Anti-catabolic — preserves muscle during periods of restriction
- Improves skin quality, collagen, and wound healing
- Cognitive benefits — IGF-1 is neuroprotective and neurogenic
- Shown to reverse catabolic state in hip fracture patients (clinical trial)

HOW IT WORKS

MK-677 mimics the action of ghrelin at the GHSR-1a receptor in the hypothalamus and pituitary, amplifying natural GH pulse amplitude without disrupting pulsatility. Unlike injectable GHRPs, the 24-hour half-life provides sustained IGF-1 elevation rather than discrete pulses. It does not suppress natural GH axis or affect cortisol at therapeutic doses.

PROTOCOLS

-
- **Standard:** 25 mg orally before bed (food optional) — maximizes GH pulse during sleep.
 - **Sensitive start:** Begin at 10 mg/day for 2 weeks, then increase to 25 mg.
 - **Cutting:** 10–15 mg to balance IGF-1 benefits against water retention and hunger.
 - **Stack:** MK-677 + CJC-1295/Ipamorelin on injection days = dual GH stimulation.
 - **Long-term:** Can run continuously — some protocols run 6–12+ months with bloodwork monitoring.

SIDE EFFECTS

- Significant appetite increase (ghrelin mimetic effect — plan calories accordingly)
- Water retention, especially first 2–4 weeks
- Temporary insulin resistance / blood glucose elevation — monitor fasting glucose
- Fatigue or lethargy (GH effect — take at bedtime to minimize)
- Tingling, numbness (carpal tunnel-like at 25 mg+)
- Potential prolactin elevation (monitor on extended cycles)

STORAGE

Oral liquid or capsule — room temperature, away from heat and light. 2-year shelf life if sealed.

■ **Monitor fasting glucose and HbA1c every 8–12 weeks on extended cycles. Water retention may cause temporary weight gain unrelated to fat.**

Educational use only. Not medical advice. Consult a qualified physician before using any compound. forgedbyfreedom.com