

Melanotan II

Melanotan II (MT-II)

TANNING

LIBIDO

FAT LOSS

APPETITE
SUPPRESSION

SPONTANEOUS
ERECTIONS

WHAT IS IT?

Melanotan II is a synthetic cyclic analogue of alpha-MSH (melanocyte-stimulating hormone). It activates multiple melanocortin receptors: MC1R (skin pigmentation), MC3R/MC4R (libido, appetite, erection), and MC5R (exocrine glands). Originally developed as a sunless tanning drug, it became popular for its libido-enhancing and appetite-suppressing side effects.

DOSING AT A GLANCE

Starting Dose	Maintenance	Timing	Protocol
0.25–0.5 mg	0.5–1 mg	Before sun/bed	Load then taper

KEY BENEFITS

- Accelerated skin tanning — deeper tan with less UV exposure required
- Significantly increased libido and sexual arousal in men and women
- Spontaneous erections in men (even without sexual stimulation at higher doses)
- Appetite suppression — meaningful fat loss aid
- Stimulates lipolysis via MC4R activation
- Enhanced sensitivity and arousal in women
- UV-protective pigmentation reduces sunburn risk

HOW IT WORKS

MT-II is a cyclic peptide analogue that binds MC1R, MC3R, MC4R, and MC5R with high affinity. MC1R activation on melanocytes stimulates eumelanin (dark pigment) production. MC4R activation in the hypothalamus drives sexual arousal, appetite suppression, and erection via neurological pathways.

PROTOCOLS

- **Loading:** 0.25 mg SC daily, gradually increase by 0.25 mg every 2–3 days to 0.5–1 mg.
- **Tanning maintenance:** 0.5 mg 2–3x/week with UV exposure.

- **Libido:** 0.5–1 mg SC 1–2 hrs before sexual activity.
- **Tip:** Start low — nausea is common at higher doses and diminishes with frequency.

SIDE EFFECTS

- Nausea and facial flushing (very common, especially on loading doses)
- Spontaneous erections (unwanted in inappropriate situations)
- Darkening of existing moles — monitor carefully
- Appetite suppression (can be profound)
- Fatigue within 30–60 min post-injection
- Pigmentation of skin (irreversible — lightens slowly when stopped)

STORAGE

Lyophilized: room temp. Reconstituted: refrigerate, use within 30 days. Light sensitive.

■ ***Mole darkening and new nevus formation reported — dermatologist monitoring recommended with regular use. Not FDA approved.***

This document is for educational purposes only. Not medical advice. Consult a qualified physician before using any peptide or performance compound. forGEDbyfreedom.com